

33rd International Symposium on Essential Oils

4 – 7 September 2002
Lisboa • Portugal

**Program, Book of Abstracts
and Participants List**



33rd International Symposium on Essential Oils

Faculty of Sciences of Lisbon, 4 to 7 September 2002, Lisbon, Portugal

Program,
Book of Abstracts
and Participants List

Editors

Ana Cristina Figueiredo
José Gonçalves Barroso
Luis Gaspar Pedro

Publisher

Centro de Biotecnologia Vegetal - Fundação da Faculdade de Ciências de Lisboa

Printer

Quinta Dimensão

Cover

TVMdesigners

ISBN: 972-9348-10-3

Additional copies can be obtained from the Editors at Faculdade de Ciências de Lisboa, C2, Piso
1, Campo Grande, 1749-016 Lisboa, Portugal

No responsibility is assumed by the Organizing and Scientific Committees on the content of the Abstracts

Table of Contents

Welcome Address	1
Acknowledgements	2
Scholarships	4
Organisation	5
Organising Committee	5
Symposium Venue.....	5
Organising Committee Contact	5
Secretariat Address	5
WWW Information.....	5
Scientific Committee	6
General Information	7
Symposium Office	7
Symposium Language	7
Badges	7
Oral Communications / Slide Preview.....	7
Posters.....	7
Poster Awards.....	7
Meals	8
Currency	8
Means of conveyance to the Symposium venue.....	8
Insurance / Liability.....	8
Scientific Program	9
Wednesday, September 4, 2002	9
Thursday, September 5, 2002	9
Friday, September 6, 2002.....	10
Saturday, September 7, 2002	11
Social Program	12
Get-together Party.....	12
Symposium Dinner	12
Half Day Tour to Lisbon	12
Accompanying Persons' Program	13
Half Day Tour to Sintra	13
Full Day Tour to Évora / Arrábida	13
Plenary Lectures Abstracts	15
Scents from Rain Forests – New Results	17
Essential oils: sample preparation and analysis.....	18
Essential oil biodiversity	20
Volatile signals: chemical structures and ecological aspects	21
New trends in intellectual property relating to perfumery materials.....	22
Biological activities of the essential oils	23
Oral Communications Abstracts	25
Posters Abstracts	43
Index	195
Participants List	205

Welcome Address

On behalf of the *Faculdade de Ciências da Universidade de Lisboa* I welcome you to our University and to the *33rd International Symposium on Essential Oils*. It is the first time that such an important meeting takes place in Portugal and we are very honoured to host it.

Looking at your programme for the next four days I was particularly impressed by the existence of a workshop devoted to the topic of “Essential oils and public opinion”. In fact, nowadays, it is extremely important that scientists, besides communicating among themselves, dedicate a special attention to the communication with the public. It is crucial that the society at large understands the importance of science and the need to support a robust research program.

Despite the fact that you have to attend the interesting talks of your full programme, I hope that you still have some spare time to visit our beautiful city of Lisbon. I wish you all a very pleasant stay in Portugal.

Prof. Dr. Augusto Barroso
The Dean of the Faculty of Sciences of Lisbon

Acknowledgements

With the high patronage of His Excellency the President of the Republic, Dr Jorge Sampaio

The 33rd International Symposium on Essential Oils was generously sponsored by:

- ❖ Faculdade de Ciências da Universidade de Lisboa
- ❖ Fundação para a Ciência e a Tecnologia: Apoio do Programa Operacional Ciência, Tecnologia, Inovação do Quadro Comunitário de Apoio III
- ❖ Reitoria da Universidade de Lisboa
- ❖ Departamento de Biologia Vegetal
- ❖ Firmenich SA
- ❖ Fundação Calouste Gulbenkian
- ❖ Ministério da Agricultura, do Desenvolvimento Rural e Pescas – Secretaria de Estado da Agricultura
- ❖ International Federation of Essential Oils and Aroma Trades (IFEAT)
- ❖ Givaudan Dubendorf Lda
- ❖ Fundação Luso-Americana
- ❖ Fundação Oriente
- ❖ Socer – Comércio e Indústrias de Resinas S.A.
- ❖ Moellhausen S.p.A
- ❖ Sociedade Portuguesa de Bioquímica
- ❖ Estética Viva

The generous obligingness of the following entities is gratefully acknowledged:

- ❖ Instituto do Vinho do Porto
- ❖ UNICER
- ❖ Câmara Municipal de Lisboa
- ❖ HA
- ❖ Banco Pinto & Sotto Mayor
- ❖ Lisboa Convention Bureau
- ❖ El Corte Inglés
- ❖ TAP Air Portugal
- ❖ Adega Cooperativa de Favaios
- ❖ Caves Neto Costa
- ❖ Comissão de Viticultura da Região dos Vinhos Verdes
- ❖ Sandeman
- ❖ Instituto de Conservação da Natureza
- ❖ Parque Natural de Sintra-Cascais
- ❖ Direcção Geral de Florestas
- ❖ Marconi
- ❖ Região de Turismo do Alto Minho
- ❖ Região de Turismo do Algarve
- ❖ Região de Turismo da Madeira
- ❖ ICEP Portugal
- ❖ Nestlé
- ❖ Imperial
- ❖ Avis

Gratitude is given to all those that contributed to make this event public:

- ❖ Divisão de Informação da Faculdade de Ciências de Lisboa
- ❖ Associação Industrial Portuguesa
- ❖ Associação Portuguesa de Horticultura
- ❖ Câmara Municipal de Lisboa
- ❖ Estética Viva
- ❖ Liga dos Amigos de Conimbriga
- ❖ Lisboa Convention Bureau
- ❖ Ordem dos Biólogos
- ❖ Ordem dos Engenheiros
- ❖ Sociedade Portuguesa de Bioquímica
- ❖ Sociedade Portuguesa de Biotecnologia
- ❖ Allured
- ❖ Elsevier
- ❖ Fitoterapia
- ❖ Herbnet
- ❖ MediConf
- ❖ Society for Medicinal Plant Research (GA)
- ❖ The American Society of Pharmacognosy
- ❖ The Phytochemical Society of Europe
- ❖ Unicamp
- ❖ Wiley

Scholarships

- ❖ The International Federation of Essential Oils and Aroma Trades (IFEAT) supported the Registration fees of the following students:

Anamarija Partl - Zavod za farmaceutsku botaniku FBFa, Croatia

Biljana Božin - Faculty of Natural Sciences, University of Novi Sad, Yugoslavia

Carlos Ribeiro - Escola Superior Agrária de Beja, Portugal

Eloisa Helena A Andrade – Coordenação de Botânica, Museu Emílio Goeldi, Brazil

Gökalp İşcan - TBAM, Anadolu University, Turkey

K. Termentzi – Aristotle University of Thessaloniki, Greece

Lin C. Ming - Department of Vegetable Production - FCA/Unesp / Botucatu, Brazil

M. C. Kalita - Dept. of Biotechnology, Gauhati University, India

Mohamed Kardali - Facultad de Biología, Universidad Complutense, Spain

Pal Apáti - Faculty of Pharmacy, Semmelweis University, Hungary

Sandy F. van Vuuren – Faculty of Health Sciences, University of Witwatersrand, South Africa

Stéphanie Cozzani - Université de Corse, France

Tiago Cunha Luis – Faculdade de Ciências da Universidade de Lisboa, Portugal

Wellington de A. Gonzaga - Universidade Federal de Santa Maria, Brazil

- ❖ Estética Viva supported the Registration fee of the following student:

Pedro António Santos – Faculdade de Ciências de Lisboa, Portugal

Organisation

Organising Committee

Prof. Dr Ana Cristina Figueiredo
Prof. Dr José Gonçalves Barroso
Prof. Dr Luis Gaspar Pedro

Symposium Venue

The Faculty of Sciences of Lisbon hosts the *33rd International Symposium on Essential Oils*.

Organising Committee Contact

33rd International Symposium on Essential Oils
Faculdade de Ciências de Lisboa
Centro de Biotecnologia Vegetal
Departamento de Biologia Vegetal
C2, Piso 1, Campo Grande 1749-016 Lisboa, Portugal
Tel.: +351217500069
FAX: +351217500048
email: acsf@fc.ul.pt

Secretariat Address

Veranatura / 33rd ISEO
Rua Abel Botelho 17-A
1500-007 Lisboa, Portugal
Tel. +351217786205 (7)
FAX: +351217786199
email: veratura@mail.telepac.pt

WWW Information

<http://biologia.fc.ul.pt/ISEO2002.htm>

Scientific Committee

Prof. Dr Ana Cristina Figueiredo, Faculty of Sciences of Lisbon, Portugal
Prof. Dr José Gonçalves Barroso, Faculty of Sciences of Lisbon, Portugal
Prof. Dr Luis Gaspar Pedro, Faculty of Sciences of Lisbon, Portugal
Prof. Dr Proença da Cunha, Faculty of Pharmacy of Coimbra, Portugal
Dr Teresa Nogueira, INETI, Portugal
Prof. Dr A. Baerheim-Svendsen, University of Oslo, Norway
Prof. Dr K.-H. Kubeczka, Wurzburg, Germany
Prof. Dr Edward Davis, Washington State University, USA
Prof. Dr Elisabeth Stahl-Biskup, University of Hamburg, Germany
Prof. Dr Gerhard Buchbauer, University of Vienna, Austria
Prof. Dr Patrizia Rubiolo, University of Torino, Italy
Dr Roman Kaiser, Givaudan Dübendorf Ltd., Switzerland
Dr Salvaterra-Garcia, Firmenich, Switzerland
Prof. Dr Wittko Francke, University of Hamburg, Germany
Prof. Dr Carlo Bicchi, University of Torino, Italy
Prof. Dr Chlodwig Franz, University of Veterinary Medicine of Vienna, Austria
Prof. Dr Johannes J. C. Scheffer, Leiden University, The Netherlands
Prof. Dr Stavros Katsiotis, University of Aristotle, Greece

General Information

Symposium Office

The Symposium Office will be located in the Foyer of the Lecture Halls in the building C3 of the Faculty of Sciences of Lisbon.

Registration and Opening Hours: Wednesday, September 4: 14:00 – 20:00
Thursday, September 5 to Saturday, September 7: 09:00 – 18:00

Symposium Language

English will be the official language of the *33rd ISEO*. There is no simultaneous translation.

Badges

Participants and accompanying persons are requested to wear their Symposium badges for identification and admittance to Scientific and Social venues.

Oral Communications / Slide Preview

The Plenary Lectures are limited to 50min, Oral Communications to 30min and Short Oral Communications to 10min, all with plus 5min for discussion. A room for slide preview is available. Overhead and slide projectors as well as MS-Power Point presentation facilities are available. Slides should be handed in at the Symposium Office, in the Foyer of the Lecture Halls, with the corresponding number of the presentation and the name of the presenting author at least 1h prior to the beginning of the respective session.

Posters

Two poster sessions are scheduled, odd numbers on Thursday and even numbers on Friday. Poster boards measuring 90cm wide and 120cm height are available for poster presentation. The Registration Office provides fixation facilities. All posters will remain exhibited throughout the symposium and the authors are recommended to be present at their posters during posters sessions.

Poster Awards

Poster awards will be granted for Layout and Scientific Content.

Meals


Owing to the short time allotted to lunch, we would highly recommend you to take lunch in the Canteen. The Canteen serves meals from 12:00 to 14:00. Lunch vouchers are required at the Canteen. You can either find the lunch vouchers in your bag if you ordered them with your registration fee, or you can purchase them at the Symposium Office. Neither the Canteen nor the Campus coffee shops accept foreign currency.


There are also several restaurants, snack bars and coffee shops in the area around the University Campus.


Currency

All payments should be done in Euro.

Means of conveyance to the Symposium venue

 1•3•7•31•31-A•35•36•47•50•67•68•83•85

 Cidade Universitária • Campo Grande

 A few parking places will be available at the Faculty parking

Insurance / Liability

The Organisers are in no respect responsible for any accommodation problem, nor for any accident, injury, theft, loss and property damage to any person during the Symposium.

Scientific Program

Wednesday, September 4, 2002

14.00 - 19.00	Registration at the Symposium Office	Foyer of the Lecture Halls
17.00 - 17.45	Opening Ceremony <i>Welcome Addresses</i> Prof. Dr Augusto Barroso, Dean of the Faculty of Sciences of Lisbon	Main Lecture Hall
17.45 – 18.00	Coffee Break	Foyer of the Lecture Halls
18.00 - 19.00	Plenary Lecture Chair: Prof. Dr Karl-Heinz Kubeczka <i>Scents from rain forests – new results</i> Dr Roman Kaiser, Givaudan Dübendorf Ltd., Switzerland	Main Lecture Hall
19.00 - 21.00	Get-together Party	

Thursday, September 5, 2002

09.00 – 10.00	Plenary Lecture Chair: Prof. Dr Johannes Scheffer & Prof. Dr Stahl-Biskup <i>Essential oils: sample preparation and analysis</i> Prof. Dr Patrizia Rubiolo, University of Torino, Italy	Main Lecture Hall
10.00 – 10.30	Oral Communications	Main Lecture Hall
10.00 – 10.15	<i>Orange concentrated essences – manufacture by fractional distillation and pervaporation</i> Dr Mércia Bettini, Flavor Tec., Brazil	
10.15 – 10.30	<i>Hydroponics - a novel way to create new aroma in living aromatic plants</i> Dr Braja Mookherjee, International Flavour & Fragrances, USA	
10.30 – 11.00	Coffee Break	Foyer of the Lecture Halls
11.00 – 12.15	Oral Communications Chair: Prof. Dr Czeslaw Wawrzenczyk & Prof. Dr Wittko Francke	Main Lecture Hall
11.00 – 11.30	<i>Rapid and reliable analysis of essential oils applying various vibrational spectroscopy methods</i> Prof. Dr Hartwig Schulz, Institute for Plant Analysis, Germany	
11.30 – 12.00	<i>Highly enantioselective cyclodextrin derivatives and conventional inner diameter short capillary columns: an approach to improve separation of medium to low volatility compounds</i> Prof. Dr Carlo Bicchi, University of Torino, Italy	
12.00 – 12.15	<i>Supercritical carbon dioxide extraction of essential oil from <i>Satureja fruticosa</i> Béguinot</i> Dr José Coelho, Technical University of Lisbon, Portugal	
12.15 – 14.00	Lunch	Canteen/Local Restaurants

14.00 – 15.30	Oral Communications	Main Lecture Hall
	Chair: Dr Brian Lawrence & Prof. Dr Hüsnu Can Başer	
14.00 – 14.30	<i>Volatile components of southern hemispheric epiphytic liverworts <i>Frullania</i> and <i>Schusterella</i> species (Frullaniaceae)</i>	
	Prof. Dr Yoshinori Asakawa, Tokushima Bunri University, Japan	
14.30 – 14.45	<i>Repeated measurement analysis as tool in essential oil research</i>	
	Dr Johannes Novak, University of Veterinary Medicine, Austria	
14.45 – 15.00	<i>Biotransformation of cuparene, plagiocilide, aristolene and nootkatone by some microorganisms</i>	
	Prof. Dr Yoshiaki Noma, Tokushima Bunri University, Japan	
15.00 – 15.30	Coffee Break	Foyer of the Lecture Halls
15.30 – 16.30	Plenary Lecture	Main Lecture Hall
	Chair: Prof. Dr Raimo Hiltunen	
	<i>Biosynthesis of essential oils</i>	
	Dr Edward Davis, Washington State University, USA	
16.30 – 18.30	Poster Session (Odd Numbers)	Foyers of the Lecture Halls

Friday, September 6, 2002

09.00 – 09.50	Plenary Lecture	Main Lecture Hall
	Chair: Prof. Dr Chlodwig Franz & Prof. Dr Gerhard Buchbauer	
	<i>Essential oil biodiversity</i>	
	Prof. Dr E. Stahl-Biskup, University of Hamburg, Germany	
09.50 – 10.30	Oral Communications	Main Lecture Hall
09.50 – 10.15	<i>Essential oils of <i>Sideritis</i> species of Turkey</i>	
	Prof. Dr Hüsnu Can Başer, Anadolu University, Turkey	
10.15 - 10.30	<i>Data bank of the Amazon aromatic plants and their essential oils</i>	
	Prof. Dr José Guilherme Maia, Museu Emílio Goeldi, Brazil	
10.30 – 11.00	Coffee Break	Foyer of the Lecture Halls
11.00 – 12.00	Plenary Lecture	Main Lecture Hall
	Chair: Prof. Dr Patrizia Rubiolo	
	<i>Volatile signals: chemical structures and ecological aspects</i>	
	Prof. Dr Wittko Francke, University of Hamburg, Germany	
12.00 – 14.00	Lunch	Canteen/Local Restaurants
14.00 – 15.00	Plenary Lecture	Main Lecture Hall
	Chair: Prof. Dr Carlo Bicchi	
	<i>New trends in intellectual property relating to perfumery materials</i>	
	Dr Maria Salvaterra-Garcia, Firmenich SA, Switzerland	
15.00 – 15.30	Coffee Break	Foyer of the Lecture Halls

15.30 – 17.00	Workshop	Main Lecture Hall
	Chair: Prof. Dr Carlo Bicchi Composition of essential oils and public opinion: Problems and solutions	
17.00 - 19.00	Poster Session (Even Numbers)	Foyers of the Lecture Halls
19.30 – 23.00	Symposium Dinner	

Saturday, September 7, 2002

09.00 – 10.00	Plenary Lecture	Main Lecture Hall
	Chair: Prof. Dr Yoshinori Asakawa & Prof. Dr Stanislaw Lochynski Biological activities of the essential oils Prof. Dr Gerhard Buchbauer, University of Vienna, Austria	
10.00 – 10.30	Oral Communications	Main Lecture Hall
10.00 – 10.15	Influence of Ylang-ylang oil on mental, emotional and human physiological parameters Dr Tapanee Hongratanaworakit, Srinakharinwirot University, Thailand	
10.15 – 10.30	Adverse effects of essential oils and components: fact and fiction Dr Maria Lis-Balchin, South Bank University, UK	
10.30 – 11.00	Coffee Break	Foyer of the Lecture Halls
11.00 – 11.45	Oral Communications	Main Lecture Hall
	Chair: Prof. Dr Karl-Heinz Kubeczka & Dr Edward Davis	
11.00 – 11.15	The essential oil of Australian Sandalwood (<i>Santalum spicatum</i>) – effects of different samples on Human physiology and subjective evaluation Dr Eva Heuberger, University of Vienna, Austria	
10.15 – 11.30	Exploring the biological activity and phytochemistry of South African medicinal aromatic plants used in traditional healing rites Dr A. M. Viljoen, University of the Witwatersrand, South Africa	
10.30 – 11.45	Composition and biological activities of essential oils of <i>Fagara zanthoxyloides</i> (dried fruits) from Cameroon as well as of <i>Syzygium cuminii</i> and <i>Syzygium travancoricum</i> (fresh leaves) from India Dr Leopold Jirovetz, University of Vienna, Austria	
11.45 – 12.00	Closing Ceremony	Main Lecture Hall
12.00 – 14.00	Lunch	Canteen/Local Restaurants
14.00 – 19.00	Half-day excursion	Half Day Tour to Lisbon

Social Program

Get-together Party

Wednesday – 4 September

The Get-together Party is included in the registration fee.

Symposium Dinner

Friday – 6 September

The Symposium dinner will take place at a local restaurant.

Half Day Tour to Lisbon

Saturday – 7 September

Departure from Symposium venue. Panoramic tour: the main avenues and squares of Lisbon. Walking tour in the typical old town to S. Jorge Castle to have a magnificent panoramic view over Lisbon downtown and the Tagus river.

The Belém district with the monuments connected with the Discoveries period: visit of the Jerónimos Monastery, masterpiece of the Portuguese Manuelin Style; visit to the Church and the Cloister, considered to be one of the most interesting ones in the world. Stop by the Belém Tower and the Discoverers' Monument.

Accompanying Persons' Program

Half Day Tour to Sintra

Thursday – 5 September

Departure from the Symposium venue. Sintra – recognized by the UNESCO as World Heritage – visit to the former Royal Palace of the 15th and 16th century: a beautiful collection of old glazed tiles, wooden painted ceilings and furniture. The picturesque old town: shopping possibility. Drive to the seaside resorts Cascais and Estoril and back to Lisbon, along the Atlantic Ocean and the Tagus river.

Full Day Tour to Évora / Arrábida

Friday – 6 September

Departure from the Symposium venue. Drive on the new Vasco da Gama bridge, one of the longest in Europe (ca. 17Km long), to the South bank of the Tagus river. The Alentejo Province – large plains with wheat and oak trees. Visit of the Museum-Town Évora: walking tour in the main streets and squares with nice marble fountains – the ruins of the Roman Temple, visit of the Cathedral (the Church, the Cloister and the Religious Art Museum) and the S. Francisco Church.

Time free for lunch. Panoramic tour on the Arrábida Mountains, with beautiful panoramic views: the Mediterranean vegetation and the Atlantic. Drive back on the Lisbon suspension bridge overlooking the town.

Plenary Lectures Abstracts



Scents from Rain Forests – New Results

Roman Kaiser

roman.kaiser@givaudan.com

Givaudan Dubendorf Ltd, Ueberlandstrasse 138, CH-8600 Dubendorf / Switzerland

Since the dawn of time, the secretive and fragile kingdom of Rain Forests, which covers 11 -13 % of all emerged land and shelters eight to nine tenths of all forms of known and unknown life, has harboured extraordinary biological treasures that deserve to be treated with the utmost care. According to a further estimate, 70 – 80 % of these species, i. e. around 60 % of the entire biodiversity, is located for ecological reason in the canopy region of the Rain Forests.

Convinced that this richness would also be reflected on the olfactory level, for the past years we have searched in the understory as well as in the canopy of Rain Forests for new attractive scents.

This communication continues the discussion of chemical, biological and olfactory aspects of such scents, which we encountered in the American, African, or Indoaustralian tropics, respectively.

The respective micro-samples have been trapped by applying our non-destructive headspace trapping method which we adapted to field experiments under extreme conditions. Complementary, a field-adapted “solid phase extraction” method (SPE) was applied if fragrant compounds had to be trapped from aqueous solutions.

Discussed among others are new highly unsaturated dodecan-5-olides identified in the flower scent of *Coryanthes* species (Orchidaceae), 3-methyl-4-alcanolides identified in that of *Passiflora chocoensis* (Passifloraceae) and *Lycaste deppei* (Orchidaceae), a new methoxy-methyl acetophenone identified as a main constituent of the flower scent of *Cyphomandra divaricata* (Solanaceae), new carotenoid-derived constituents in the flower scents of *Anthurium salvadorensense* (Araceae) and *Palisota* species GHS06 (Commelinaceae) as well as unusual esters of 2-pentanol in that of *Hoya* vs. *sussuela* (Asclepiadaceae).

Essential oils: sample preparation and analysis

Patrizia Rubiolo

patrizia.rubiolo@unito.it

Dipartimento di Scienza e Tecnologia del Farmaco – Via P.Giuria 9 – I - 10125 Torino, Italy

The definition of essential oil, extremely precise and internationally accepted, is “An essential oil is the product obtained by hydrodistillation, steam distillation or expression (for *Citrus* fruit) of a plant”.

Since essential oil components are generally medium to highly volatile compounds with medium to low polarity (terpenoids, phenyl propanoids and so on), GC is the technique of choice for their analysis.

In this definition, sample preparation is considered very important and fundamental although hydrodistillation or steam distillation are not the only techniques used to obtain the so-called “volatile fraction” of a plant. Therefore it is necessary to use a correct terminology to indicate the volatile or semi-volatile fraction of a plant, in particular it would be better to talk about the “volatile fraction” and to use the definition “essential oil” only when the sample preparation is distillation or expression.

This lecture will deal with the following main topics:

1. Alternative techniques to distillation: this part will mainly deal with advantages and disadvantages of SFE as an alternative technique to distillation to *extract* the volatile fraction in comparison to the classical steam- or hydrodistillation.

2. High concentration capacity static headspace sampling techniques: this part will be focussed on recently introduced techniques such as headspace-solid phase microextraction (HS-SPME) and headspace sorptive extraction (HSSE) that have opened new perspectives in the analysis of the volatile fraction allowing us to run a static headspace sampling but at the same time concentrating trace components.

3. Recent improvements in GC analysis of essential oils: this part will concern the recent achievement in GC analysis of essential oils in particular with the so called “high speed (or fast) GC” with both narrow bore and conventional inner diameter short columns.

Biosynthesis of essential oils

Edward M. Davis and Rodney Croteau

edd@mail.wsu.edu

Institute of Biological Chemistry, Washington State University, Pullman, WA, USA

Monoterpenes comprise structurally diverse, major components of many essential oils, and the biosynthesis of the monoterpenes of peppermint (*Mentha x piperita*) have provided a model system for their study. Recent advances in the isolation of intact epidermal oil glands (peltate glandular trichomes) have permitted investigations of the biosynthesis and developmental regulation of metabolism of these C₃ oxygenated *p*-menthane monoterpenes, and have provided an enriched source of mRNA for the generation of an expressed sequence tag (EST) library. The identification and heterologous overexpression of genes responsible for the biosynthesis of the universal monoterpene precursor, geranyl diphosphate, and for downstream steps leading to the primary peppermint monoterpene, menthol, have allowed detailed characterization of these catalysts and the transgenic manipulation of peppermint oil composition and yield.

Essential oil biodiversity

Elisabeth Stahl-Biskup

elisabeth.stahl-biskup@uni-hamburg.de

Universität Hamburg, Institut für Pharmazie, Abt. Pharmazeutische Biologie and Mikrobiologie, Bundesstrasse 45, D-20146 Hamburg

The expression "biodiversity" was coined in the nineties of the last century by the biologist Edward O. Wilson who tried to describe the total variety of the world's living organisms including all genes, species and ecosystems, and the ecological processes of which they are part. No wonder that an internet search "biodiversity" provides about 1,200,000 hits, a fact which reflects its broad meaning in biological contexts. Discussing essential oil biodiversity we focus on the chemical polymorphism in aromatic plants, which means the infraspecific variation of their essential oil patterns in qualitative and quantitative respects. The levels of examination can be represented by species, populations or even individuals. Essential oil polymorphism is due to the fact that the biogenesis of terpenoids as well as of phenylpropane derivatives are genetically controlled by several steps. Furthermore ontogenetic variability and environmental factors can be considered to influence the biosynthesis. Essential oil polymorphism is widespread in the plant kingdom; in 1975 Tétényi estimated 360 species of 36 families which were known to be chemically polymorphous.

As a result of the understanding that it is a fundamental requirement to analyse infraspecific differences, we encounter a flood of reports on polychemism. Therefore, nowadays the compilation of all species which show essential oil polymorphism would be a worthwhile but arduous undertaking. It is a fact that the Lamiaceae have always been an interesting object of polychemism research (Lawrence, 1980), the genus *Thymus* being one of the most frequently investigated and the object of the most detailed research regarding this phenomenon (Sáez and Stahl-Biskup, 2002). Since the early 1960s, one species, *Thymus vulgaris*, has been at the heart of ecological and genetic research on the evolutionary dynamics of genetic polymorphism. Its essential oil variation has a genetic basis and the presence of six distinct genetically-based forms has thus provided an attractive system to explore the ecological genetics of secondary compound variation and the population structure and spatial dynamics of genetic polymorphism in thyme (Thompson, 2002).

Nowadays taxonomists agree that the analysis of the chemical polymorphism can only be completed analysing a huge number of individuals producing an immense amount of quantitative data which can only be categorized and classified by computer-based multivariate statistical analyses. In this respect the genus *Thymus* again is a good example for demonstrating the methods to discover and to document essential oil polychemism. Special focus will be put on *Thymus* in south-eastern Spain, western Iberia and northern Europe. As a résumé it can be anticipated that in *Thymus* two forms of polymorphism are manifested: some species occur with only few chemotypes, other species show more than 7 or even an uncertain number of chemotypes. The highest chemical variability seems to be concentrated in species from the sections *Serpyllum* and *Thymus*.

Lawrence BM (1980) VIIe Congrès International des Huiles Essentielles, Octobre, Cannes-Grasse, 118-131.

Sáez F, Stahl-Biskup E (2002) In: *Thyme –The Genus Thymus* (eds: Stahl-Biskup E, Sáez F), Vol. 24, Series "Medicinal and Aromatic Plants – Industrial Profiles" (ed. Hardman R), Taylor & Francis, London, pp 125 (in press).

Tétényi P (1975) *Planta Med.* **28**: 244-256.

Thompson JD (2002) In: *Thyme –The Genus Thymus*, pp. 44; Vol. 24, Series "Medicinal and Aromatic Plants – Industrial Profiles" (ed. Hardman R) Taylor & Francis, London, pp 125 (in press).

Volatile signals: chemical structures and ecological aspects

Wittko Francke

francke@chemie.uni-hamburg.de

Institut für Organische Chemie der Universität Hamburg, Martin-Luther-King-Platz 6, D-20146 Hamburg

Molecular recognition is prerequisite to the beginning of life, and thus, “chemical signalling“ is the oldest means for the transmission of information.

During evolution, principles of chemical communication may have been developed several times and for different reasons, typical elements being made up by secondary metabolites.

Most of the relevant compounds are represented by acetogenins, polyketides and terpenoids, the biosynthesis of which is not restricted to animals, but is also valid in plants and microorganisms, in terrestrial as well as in aquatic ecosystems.

Typical sex pheromones of moths are *de novo* produced medium chain unbranched, unsaturated, even numbered acetogenins with an oxygen containing functional group at one end of the chain. There are, however, secondary alcohols and polyenes, including the corresponding epoxides, showing uneven numbers of carbon atoms. This latter group represents degradation products of fatty acids, i.e. β -oxidation/decarboxylation or epoxidation/decarboxylation, respectively.

Replacement of acetate units by propanoate during the synthesis of a typical acetogenin will result in methyl branching which adds to the variability of principle structures since the position of the methyl group and the configuration of the stereogenic center may account for the species specificity of the compound and for its biological activity. The same is true for polyketides which are produced from propanoates, exclusively. Relevant substances are made up of 2-5 propanoate units showing several stereocenters at the branching points as well as at the hydroxy groups originating from the aldol-type biogenesis of the compounds. Stereochemical compositions play a particularly important role in the biological activity of such structures. The possible role of (endo)symbionts in their production and transformation is yet unknown.

A third large group of semiochemicals is represented by terpenes. They may be sequestered and used unchanged or oxidised and/or rearranged, however, they may also be produced *de novo*. The involvement of propanoate units in the principle biosynthesis of terpenes results in the formation of homomevalonate, which finally yields homoterpenes. Again, the additional methyl group gives rise to stereoisomers which enhance the specificity of the produced signal.

Sequestered food constituents which animals use as chemical signals, provide information about the quality of the environment. The more such compounds are altered through metabolic activities in the animal (finally leading to a totally *de novo* synthesis), the more they reflect the physiological state of the emitter. During biosynthesis of semiochemicals, the action of less specific enzymes will lead to the production of „byproducts“ (stereoisomers, chain length) of the “actual signal“ which may add a fine tuning of the bouquet or facilitate evolution through the introduction of a „disposable variant“. In many cases, stereochemistry plays a major role in chemical communication: A well defined (correct) composition of positional and geometrical isomers or of enantiomers and diastereomers of attractants are decisive for an optimal biological activity. Closely related species may simply generate specific blends by producing different blends (including different stereochemical proportions) of the same compounds – which may even be mutually deterrent.

Concerning aspects of coevolution between a producer (biosynthesis and release) and a receiver (perception and transduction) the problem strongly resembles the story of the hen and the egg.

New trends in intellectual property relating to perfumery materials

Maria Salvaterra-Garcia

maria.garcia@firmenich.com

Firmenich SA, Corporate R&D Division, P. O. Box 239, 1211 Geneva 8, Switzerland

It has been traditional in all industries, and that of Perfumery is no exception, to follow innovation through the patent literature since, if patent law functions properly, it enhances the incentive to innovate.

The trend already observed a few years ago in the flavor and fragrance industry towards the patenting of more complex "systems", rather than just "molecules" as such, seems to have accelerated, although a few of the industry's players have remained strong in the research and protection of the latter.

Other recent patent trends relate to the protection of new products for manufacturing odorants through molecular biology methods.

Finally, outsiders to the industry are actively pursuing research and patenting of olfactory receptors, the use of which could in the future complement the traditional creation and formulation of fragrances and flavors.

These different tendencies will be discussed through some examples.

Biological activities of the essential oils

Gerhard Buchbauer

gerhard.buchbauer@univie.ac.at

Institute of Pharmaceutical Chemistry, University of Vienna, A-1090 Vienna, Austria

A review is given on scientifically proven activities of essential oils. Essential oils (EO) as natural mixtures (sometimes up to more than 250 single volatile compounds) are very valuable natural products and used in many fields, e.g. fragrance and perfumes, cosmetics, aromatherapy and phytotherapy, spices and nutrition, sources of raw materials. Above all, EO are used as therapeutic agents against various ailments, disorders and psychosomatic complaints. Contrary to esoteric and holistic believes, the rather small and lipophilic fragrance compounds of an EO exhibit distinct physiological activities (besides their sensorial qualities) which can be shown in animal and human experiments.

The psychological effects, e.g. the creation of a sort of well feeling, happiness, etc. and especially the physiological/pharmacological activities are topic of this review. Own research results obtained with healthy subjects as well as experimental data found by other scientists are presented. In addition also some exemplary and interesting animal experiments are included in this report. Finally, some pros and cons regarding the therapeutically use of EO are mentioned.

Oral Communications Abstracts



Orange concentrated essences – Manufacture by fractional distillation and pervaporation

Mércia de Fátima M. Bettini^a, Loudes Maria Corrêa Cabral^b

flavortec@flavortec.com.br; lcabral@ctaa.embrapa.br

^a Flavor Tec – Aromas de Frutas Ltda- Av.Bela Vista-971, Pque Ind., Pindorama, SP, Brazil, CEP- 15830-00

^b Embrapa Agroindústria de Alimentos – Av. Américas, 29501, Bairro Guaratiba, Rio de Janeiro, Brazil, CEP-23020-470

The citrus juice processing generates large amounts of wastes. Almost 50% of the fruit weight finishes as waste in the form of peel, segment membranes, rags and seeds. Besides that, during juice concentration, volatile constituents contained on the vapour phase are condensed by the aroma recovery systems. The missing of these aromatic constituents results in a decrease of juice quality, unless these compounds are being reincorporated on it. The aroma recovery systems get two separated phases: essence oil (oil phase) and aqueous essence (water phase). The average yield of these products is very low. It's necessary around 2,000 kg of oranges to produce 1 kg of water phase and 8,000 Kg of oranges for 1 Kg of oil phase. These two products are rich in aldehydes, esters and other volatile compounds. Oil phase has a high concentration of terpenes and water phase is rich in alcohol. As the storage and transportation of these products are expensive, the evaluation of concentration processes of these essences is of high interest.

The aim of this work is to study the special fractions of these aromatic volatile compounds from the oil phase and water phase, obtained by fractional distillation and pervaporation. The fractional distillation is far the most used process, mainly after the development of efficient columns, which can reach 100 or more theoretical plates. Nowadays, it is possible to conduct efficient fractional processes. Relative to alternative treating techniques, pervaporation provides a system which is simple to operate and easy to maintain. The high reliability and versatility of pervaporation systems make pervaporation an attractive choice for applications involving the recovery or removal of volatile organics from water streams.

By fractional distillation of the oil phase, concentrated fractions with high contents of fragrant compounds as ethyl butyrate and valencene were obtained. For the concentration of the water phase, the two processes, fractional distillation and pervaporation, were carried out. Essences with ethanol content higher than 80% were obtained. Their smell and taste, proceeding from the original esters and aldehydes of the raw materials, such as acetaldehyde, seems to be very good.

The evaluation of these concentrated essences was performed by physicochemical and chromatographic analysis. Different amounts of these concentrated products were added to juices and soft drinks, showing wide acceptance by sensory evaluations.

Hydroponics - a novel way to create new aroma in living aromatic plants

Braja D. Mookherjee, Subha M. Patel, Baoping Jin and Robert W. Trenkle

BD.Mookherjee@iff.com

Research & Development, IFF, 1515 Highway #36, Union Beach, NJ, USA, 07735

Both in ancient and present time, various people have grown plants hydroponically. It is known in scientific literature that hydroponically grown vegetables have different tastes due to their change in protein, carbohydrate and fat of the aroma contents. But to our knowledge, nobody has ever studied the fragrance composition of the hydroponically grown aromatic plants. We have grown various flowers, viz. Hyacinth, Rose, Jasmin, fruits, like lime, lemon, strawberry and many herbs and spice plants and have studied the composition of these hydroponically grown living flowering plants with those grown under the conventional soil method. It was found that these hydroponically grown flower, fruits, herbs and spices not only have superior growth patterns, but they also produce different aroma constituents than the conventionally grown plants.



As for example, in the case of Hyacinth, phenyl ethyl alcohol in conventionally grown Hyacinth is 3.3%, which has increased to 23% in hydroponically grown Hyacinth. Benzyl Tiglate, which is 12% in conventionally grown Hyacinth, is not detected in the hydroponically grown hyacinth. In the case of hybrid tea rose, the green note components, viz. *cis*-3-hexenyl acetate and hexyl acetate have decreased from 7% in conventionally grown rose to 0.3% in hydroponically grown rose. At the same time, Citral content, which is 10% in conventionally grown rose, decreases to 5% in the hydroponically grown rose. In the case of strawberry, ethyl acetate content which is 12% in conventional grown fruit decreases to 0.1% in hydroponically grown strawberry. Ethyl caproate, which is 35.9% in conventionally grown strawberry, has decreased to 9.7% in hydroponically grown strawberry. On the other hand, 2-nonanone, which is 3.6% in conventionally grown strawberry, has increased to 19% in hydroponically grown strawberry. In the case of Persian Lime, limonene content has increased from 11.6% in conventionally grown fruit to 23% in the hydroponic condition, whereas Citral, which is 22% in conventional has reduced to 7% under hydroponic condition.

Present paper will describe in details the difference between the aroma chemistry of the hydroponically grown flowers, fruits, herbs and spices to those of the soil grown plants. In addition, the significance of the aroma of hydroponically grown flowers in the fragrance industry will be illustrated.

Rapid and reliable analysis of essential oils applying various vibrational spectroscopy methods

Hartwig Schulz

H.Schulz@bafz.de

Federal Centre of Breeding Research on Cultivated Plants, Institute for Plant Analysis, Neuer Weg 22-23, D-06484 Quedlinburg, Germany

During the last years near-infrared spectroscopy (NIRS) has been successfully introduced as an efficient tool for rapid and reliable analyses of various essential oils [Schulz *et al.*, 1999; Steuer *et al.*, 2001]. Usually the NIR spectra (range: 1100 to 2500 nm) are registered in the transfection mode using quartz cuvettes, equipped with a diffuse gold reflector (path length 0.2 mm). Since NIRS is an indirect analysis method, at first it is necessary to establish calibration equations for the individual essential oil substances (analytes). In this context standard GC methods are applied to obtain the reference data. In most cases statistical errors of the NIR predicted data (SECV) are not significantly higher than the relating standard deviations of the GC method. The individually developed calibration statistics can be used not only for rapid quality control purposes but they can be also successfully introduced to monitor the enrichment of valuable essential oil components during distillation or extraction processes.

The recent availability of attenuated total reflectance (ATR) technology for mid-infrared spectroscopy has made the handling of liquid samples very quick and simple. Whereas in the past MIR spectroscopy was frequently used as a qualitative technique for the identification and verification of pure essential oil substances, up to now there exist only a few applications in this area which use multivariate calibration algorithms to extract the quantitative data from the rich in information and well-structured MIR signals. Today portable Fourier-Transform (FT) IR spectrometers are available which need only sample amounts of several microliters. The simultaneous analysis of several oil components can be performed in some seconds [Schulz *et al.*, 2002].

Also NIR-FT-Raman spectroscopy, which is to be seen as a complementary method to MIR, has been successfully applied for quality analysis purposes of various essential oils [Schulz *et al.*, 2002]. The spectra were processed with a PLS algorithm and the accuracy of the individual calibration models is generally characterized by the statistical parameters R^2 and SECV. Compared to the NIR calibration results generally a lower detection limit is reached due to the larger signals of individual fundamental vibration modes. All spectroscopic techniques described here have the potential to replace existing analytical procedures because they are capable of faster quality control measurements, and they allow on-line analyses during distillation or extraction processes to be performed. In addition to rapid analysis of isolated essential oils, partly the described spectroscopic techniques have also potential to determine non-destructively the amount and composition of essential oils directly in the fresh plant or in the relating drug without performing any clean-up procedures. Furthermore there exists the option to connect a Raman spectrometer with a microscope which offers the opportunity to map in-vivo the occurrence of selected essential oil substances in the plant matrix.

Schulz H, HH Drews, H Krüger (1999) *J. Essent. Oil Res.* **11**, 185-190.

Schulz H, B Schrader, R Quilitzsch, B Steuer (2002) *Appl. Spectrosc.* **56**, 117-124.

Steuer B, H Schulz, E Läger (2001) *Food Chemistry* **72**, 113-117.

Highly enantioselective cyclodextrin derivatives and conventional inner diameter short capillary columns: an approach to improve separation of medium to low volatility compounds

Carlo Bicchi, Claudio Brunelli, Chiara Cordero, Patrizia Rubiolo

carlo.bicchi@unito.it

Dipartimento di Scienza e Tecnologia del Farmaco, Via P. Giuria 9, i-10125 Torino, Italy

This communication reports the results obtained by combining highly enantioselective cyclodextrin derivatives with conventional inner diameter short capillary columns in the enantiomer separation of medium to low volatility compounds.

Cyclodextrins (CDs) are successful enantioselective chiral selector used in the gas chromatographic (GC) separation of underivatized volatile enantiomers. The CD ring size and the substituents of the sugar units at C-2, C-3 and C-6 hydroxylated positions influence the CD enantioselectivity and chemical and physical properties.

6-tert-Butyl-Di-Methyl-Silyl (TBDMS)- β -CDs substituted in positions 2 and 3 with methyl (ME) or ethyl (ET) groups are among the most effective chiral selector for enantioselective GC and show enantioselectivities that very often are complementary. To increase enantioselectivity completely asymmetric ME-ET-6-TBDMS- β -CD were synthesised and tested as GC stationary phases, with the aim of developing CDs of as universal as possible application, and of evaluating how the simultaneous introduction of ME and ET groups in positions 2 and 3 of CD ring influences enantioselectivity.

The increased enantioselectivity of the new derivatives affords to analyse several racemates with shorter columns still obtaining enantiomer base line separation. Shorter columns not only contribute to shorten analysis time thus overcoming one of the main drawbacks of enantioselective GC with CD derivatives, but also concur to reduce the enantiomer elution temperature. This favours separation of enantiomers (in particular of those of medium-to-low-volatility racemates) because of the very low difference in energy involved in the enantiomer host-guest interaction whose discrimination is favoured by a lower temperature.

Supercritical carbon dioxide extraction of essential oil from *Satureja fruticosa* Béguinot

A. P. Pereira^a, J. S. Urieta^b, J. Burillo^b, A. C. Figueiredo^c, J. G. Barroso^c, R. L. Mendes^a, J. A. Coelho^a, A. M. F. Palavra^a

jcoelho@deq.isel.ipl.pt

^aCentro de Química Estrutural, IST, Lisbon, Portugal

^bDep. Química-Física, Universidad - Servicio de Investigación Agroalimentario, Zaragoza, Spain

^cCentro de Biotecnologia Vegetal, Dep. de Biología Vegetal, FCL, C2, Campo Grande, 1749-017 Lisbon, Portugal

Aromatic plants produce essential oils, which have important biological functions in plant metabolism. The quality of these oils is also very important in human consumption. The main health applications of *Satureja fruticosa* Béguinot are due to its digestive and gastric properties.

Supercritical fluid extraction (SFE) is a separation process, based on the solvent power of supercritical fluids, which can be controlled by changing the pressure and temperature (Palavra *et al.*, 1993, 1995). With this technique it is possible to obtain solvent-free extracts and avoid the degradation of thermally labile components existing in the oils. Therefore, the natural odour and flavour of the plant are maintained, thus making SFE a very promising process for the separation of essential oils from herbaceous matrices (Palavra *et al.*, 1997).

Flowers and leaves of *Satureja fruticosa* Béguinot, cultivated at Servicio de Investigación Agroalimentario, Zaragoza, Spain, were used as matrices in supercritical CO₂ extraction followed by a two stage fractional separation. The process was carried out in a flow apparatus, provided with an extraction vessel (1 L) and two separators (0.27 L, each) at temperatures of 40 and 50 °C and pressures of 90 and 100 bar.

The isolated essential oils were analysed by GC-MS and GC. The oxygen-containing monoterpenes, pulegone (46-56 %), piperitenone (10-13 %), piperitenone oxide (9-14 %) and isomenthone (7-9 %) were the major oil constituents.

The effect of the plant particle size, extraction time and solvent flow rate on the extraction yield and quality of the obtained oil is discussed.

Bruno T, CAN Castro, JF Hammel, AMF Palavra (1993) Supercritical Fluid Extraction of Biological Products. In Recovery Processes for Biological Materials, eds. JF Kennedy, JMS Cabral, J. Wiley & Sons, Chichester.

Coelho JP, RL Mendes, MC Provost, JMS Cabral, JM Novais, AMF Palavra (1997) "Supercritical CO₂ Extraction of Volatile Compounds from Rosemary", ACS Symposium Series, 670: 101-109, in Supercritical Fluids Extraction and Pollution Prevent, ed. A Martin, A Abraham and AK Sunal.

Mendes RL, JP Coelho, HL Fernandes, IJ Marrucho, JMS Cabral, JM Novais, AF Palavra (1995) J. Chem. Tech. Biotechnol., 62: 53-59.

Volatile components of southern hemispheric epiphytic liverworts *Frullania* and *Schusterella* species (Frullaniaceae)


Y. Asakawa^a, M. Toyota^a, A. Bardon^b, N. I. Kamiya^b, M. von Konrat^c, J. Braggins^c

asakawa@ph.bunri-u.ac.jp

^aFaculty of Pharmaceutical Sciences, Tokushima Bunri University, Yamashiro-cho, 180, Tokushima 770-8514, Japan; ^bInstituto de Química Organica, Facultad de Bioquímica, Química y Farmacia, Universidad Nacional de Tucuman, Ayacucho 471, Tucuman 4000, Argentina; ^cSchool of Biological Sciences, The University of Auckland, Private Bag, 92019, New Zealand

A number of liverworts and mosses are medicinal plants and are said to possess certain biological activity (cytotoxic, antimicrobial, antifungal, antidotal, antipyretic, anticancer, sedative, diuretic and antiseptic activity etc.) and effect (for cardiopathy, hemostatis, pulmonary tuberculosis and neurasthenia etc.) (Asakawa 1982, 1995 & 1999). Many liverworts also show characteristic fragrant odour and hot-taste.

There are ca. 300 of *Frullania* species (Frullaniaceae). Some sesquiterpene lactones isolated from *Frullania* species display potent allergenic contact dermatitis, piscicidal activity and cytotoxicity against certain cancer cell lines (Asakawa 1995). We are continuing to study the chemical constituents of the southern hemispheric epiphytic *Frullania* and *Schusterella* species from view-point of search on the biologically active substances and chemosystematics of the Frullaniaceae.



The volatile components of *Frullania* and *S. chevalierii* have been analyzed by GC-MS. The major components were isolated by HPLC and their absolute configurations established by spectroscopy and X-ray crystallographic analysis. The present *Frullania* species are chemically divided into five groups: 1) Sesquiterpene lactone, 2) Sesquiterpene lactone-bibenzyl, 3) Bibenzyl, 4) 2-Alkanone and 5) the others. *S. chevalierii* is closely related chemically to sesquiterpene lactone group of the *Frullania* species since it elaborates two eudesmanolides, β -cyclocostunolide and its dihydro derivative, as the major components.

The Argentine *F. brasiliensis* produces both new eremophilanolides (**1,2**) and eudesmanolides (**3-6**), along with a new 3,3',4-trimethoxybibenzyl. This species resembles chemically the European *Frullania dilatata* (Asakawa *et al.* 1976, Nagashima *et al.* 1999).

Asakawa Y, Muller J-C, Ourisson G, Foussereau J, Ducombs G. (1976) *Bull. Soc. Chim. France (Chim. Mol.)* 1465.

Asakawa Y. (1982) *Progress in the Chemistry of Organic Natural Products*, **42**, 1-285. Springer, Wien.

Asakawa Y. (1995) *Progress in the Chemistry of Organic Natural Products*, **65**, 1-618. Springer, Wien.

Asakawa Y. (1999) *Recent Advance in Phytochemistry* **33**, 319-342. Kluwer Academic/Plenum Publishers, N. York.

Nagashima F, Takaoka S, Huneck S, Asakawa Y (1999) *Phytochemistry* **37**, 1317-1321.

Repeated measurement analysis as tool in essential oil research

Johannes Novak, Chlodwig M. Franz

Johannes.Novak@vu-wien.ac.at

Institute for Applied Botany, University of Veterinary Medicine, Veterinärplatz 1, A-1210 Wien, Austria

‘Repeated measurement analysis’ is a special statistical procedure to analyse groups of dependent variables, which are measurements of the same trait. By repeating essential oil analysis of individuals in time or space, the accuracy of the determination of either environmental or ontogenetical influences (repetition in time) or plant organ differences (repetition in space) is enhanced.

To demonstrate the capability of this method, two model experiments are presented. In one experiment, a population of marjoram (*Origanum majorana*) was sampled twice at the same stage of development but grown under different environmental conditions. In the second example, different plant organs of monks-pepper (*Vitex agnus-castus*) from individual plants of several accessions were analysed.

Biotransformation of cuparene, plagiocbilide, aristolene and nootkatone by some microorganisms

T. Hashimoto^a, Y. Noma^b, N. Nishimatsu^a, S. Onishi^a, M. Furusawa^b, Y. Asakawa^a


ynoma@tokushima.bunri-u.ac.jp

^aFaculty of Pharmaceutical Sciences, Tokushima Bunri University, Yamashiro-cho, Tokushima 770-8514 Japan

^bFaculty of Domestic Sciences, Tokushima Bunri University, Yamashiro-cho, Tokushima 770-8514 Japan

We are continuing to study the biotransformation of sesquiterpenoids by several microorganisms and mammals to obtain functional substances (Noma & Asakawa 1995; Hashimoto, Noma & Asakawa 2001).

The present paper concerns the biotransformation of four sesquiterpenoids, (-)-cuparene (**1**) and plagiocbilide (**2**) which are the major sesquiterpenes isolated from the liverworts *Marchantia polymorpha* and *Plagiochila fruticosa*, respectively, and aristolene (**3**) from the crude drug *Nardostachys chinensis* and nootkatone (**4**) from grape fruit by *Aspergillus niger*, *Bortyosphaeria dothidea* or *Chorella pyrenoidosa*.



(-)-Cuparene (**1**) was biotransformed to compounds **5-7** by *A. niger* as the major products. Gem-dimethyls of **1** was not oxidized at all. On the other hand, one of the gem-dimethyls of plagiocbilide (**2**) and aristolene (**3**) was oxidized by the same fungus to give **8** and **9** from **2**, and **10-11** from **3**. Aristolene (**3**) was also converted to **12** by *C. pyrenoidosa* as the major component. 7-Hydronootkatone (**13**) was mainly obtained from nootkatone (**4**) by *Bortyosphaeria dothidea*. On the other hand, 11,12-dihydroxynootkatone (**14**) and 12-hydroxynootkatone (**15**) were major products from **13** by *A. niger*.

Hashimoto T, Noma Y, Asakawa Y. (2001) *Heterocycles* 54, 529-559.

Noma Y, Asakawa Y. (1995) In: *Biotechnology in agriculture and forestry. medicinal and aromatic plants VIII* (Bajaj, YPS ed.) Vol. 33, pp. 62-96 (1996) Springer Berlin, Heidelberg.

Essential oils of *Sideritis* species of Turkey

K. Hüsnu Can Başer^a, Neşe Kırimer^a, Hayri Duman^b, Betül Demirci^a

khcbaser@anadolu.edu.tr

^a Medicinal and Aromatic Plant and Drug Research Centre (TBAM), Anadolu University, 26470-Eskişehir, Turkey.

^b Gazi University, Faculty of Science and Letters, 06500-Ankara, Turkey.

Anatolian peninsula is one of the two main gene-centres of the genus *Sideritis* L. (Lamiaceae) together with the Iberic peninsula. While Turkey contains *Sideritis* species belonging to Sect. *Empedoclia* (Rafin.) Benth and Sect. *Hesiodia* Benth, the Spanish *Sideritis* species belong to Sect. *Sideritis*. Species in Sect. *Empedoclia* are used for making tea in Turkey and Greece.

We have collected all the *Sideritis* species growing wild in Turkey and investigated their essential oil compositions by GC and GC/MS. The paper will present a comprehensive view of the *Sideritis* species of Turkey as regards their essential oil composition.

Data Bank of the Amazon aromatic plants and their essential oils

José Guilherme S. Maia, Eloisa Helena A. Andrade, Milton Helio L. da Silva, Maria das Graças B. Zoghbi, Léa Maria M. Carreira

gmaia@museu-goeldi.br

Museu Emilio Goeldi, Departamento de Botânica, P.O. Box 399, 66040-170 Belém, Brazil

As part of the inventory of the Amazon odoriferous flora we created a data bank based on the aromatic plants and their essential oils. Since 1980 we travelled all over 50,000 km, visiting numberless collection sites in all nine Amazonian states and at different ecosystems. We collected over 2,500 aromatic part plants including leaves, stems, barks, trunk woods, rhizomes, fruits, flowers, resins, etc. For economic purpose the last 6 years of this work was addressed to collect fast growing plants (herbs and shrubs) occurring in areas of natural fields and secondary forests like savanas and cerrados. Their essential oils were hydrodistilled and analysed by GC-MS using WCOT SE-54 and DB-5 fused silica capillary columns. Individual components were identified by comparison of both mass spectrum and GC retention data with those of authentic compounds previously analysed and stored in the data system or cited in the literature (Maia *et al.* 2002; Adams 1995). Today, the Data Bank contains about 1250 registered specimens providing the following informations: families, scientific and common names and synonyms; collection sites; geographical distribution; habitat; popular uses; oil yields; part of plant furnishing the essential oils; botany, agronomic and ecological sights; GC and GC-MS ion-chromatograms; bibliography and photos. The Data Bank provide also information about the essential oil application according previous bioassays and fragrant and olfative perfurter analyses. At present, from Data Bank were published over 100 scientific papers reporting the Amazon essential oils. It is available for oral presentation in Windows Access ambient.

Acknowledgements: supported by MCT-PPG7/European Community.

Adams RP (1995) Identification of Essential Oil Component by Gas Chromatography/Mass Spectroscopy, Allured, Illinois.

Maia JGS, Silva MHL da, Andrade EHA, Zoghbi MGB, Carreira LMM (2002) *Flavour and Fragrance Journal* 17: 72-74.

Influence of Ylang-ylang oil on mental, emotional and human physiological parameters

Tapanee Hongratanaworakit^a, Gerhard Buchbauer^b

htapanee@hotmail.com, gerhard.buchbauer@univie.ac.at

^aDepartment of Pharmaceutical Chemistry, Faculty of Pharmaceutical Sciences, Srinakharinwirot University, Nakorn-nayok, Thailand

^bInstitute of Pharmaceutical Chemistry, Center of Pharmacy, University of Vienna, Vienna, Austria.

Ylang-ylang oil (*Cananga odorata*) is widely used as a fragrance in perfumery and cosmetic industries. In medicine the interest in the usage of ylang-ylang oil as therapeutically active agent has grown considerably. Especially in aromatherapy, ylang-ylang oil has been used as an antidepressant in the case of depression and nervousness as well as used for reducing the blood pressure in the case of hypertension. In spite of its widespread use, scientific evaluation of the effects of this fragrance on human is rather scarce. The aim of this study was to investigate the influence of ylang-ylang oil on mental, emotional, and human physiological parameters. The fragrance was administered by inhalation. Physiological parameters recorded were blood pressure, breathing rate, skin temperature, and heart rate. Self-evaluation was assessed in terms of alertness, attentiveness, calmness, mood, relaxation, and vigour. Additionally, the fragrance was rated in terms of pleasantness, intensity, and effect.

The present study revealed that ylang-ylang oil affected both physiological parameters and subjective mental and emotional conditions. Ylang-ylang oil caused a significant decrease of blood pressure, which are likely to represent a sympatholytic effect, i.e. dilation of blood vessels and lowering blood pressure. Furthermore, ylang-ylang oil caused a significant increase of subjective attentiveness, which seem to show a harmonizing effect of the oil. Correlational analyses revealed interactions between physiological parameters and self-evaluation in all groups.

These findings give an evidence for the usage of the ylang-ylang oil in aromatherapy such as furnishing a reduction of blood pressure or a relief of depression and stress in humans.

Adverse effects of essential oils and components: Fact and Fiction

Maria Lis-Balchin

lisbalmt@sbu.ac.uk

School of Applied Science, South Bank University, Borough Road, London, SE1 OAA, UK

Many essential oils and extracts used in the food and cosmetics industries are already banned or severely restricted due to their toxic properties (wormseed, wintergreen); carcinogenicity (calamus and saffron); sensitisation (benzoin, Peru balsam); photosensitization (bergamot, expressed). Components like cinnamaldehyde are also restricted due to sensitisation. Extensive research by RIFM has been conducted on the allergic effects of fragrances on skin, many of which have been shown to cause dermal allergic reactions, but they have rarely tested the effects of inhaling fragrance chemicals. An estimated 5.72 million in the US have skin allergy to fragrance. Around 25% of the general population report some sensitivity to chemicals and of these more than 80% report that exposure to fragrances has a major impact on their quality of life. Symptoms include: headaches, dizziness, nausea, fatigue, shortness of breath, difficulty with concentration, and allergy-like symptoms. Fragrances trigger or may actually cause asthma.

Some sceptics (mainly in the cosmetics business) suggest that reactions to fragrance could be psychological i.e. a conditioned response, which could include elements of paranoia, stress-induced psychosomatic symptoms, hypochondriasis, hysteria, and suggestibility.

Due to secrecy laws, adverse reactions to fragrances may be difficult or impossible to link to particular fragrance chemicals. This is complicated by synthetic organic chemicals, which constitute more than 80-90% of the raw materials used in flavour and fragrance formulations. Chemicals that appeared in more than half of cosmetic products included limonene (showed carcinogenicity in male rats, but not in mice or female rats), linalool, β -phenethyl alcohol, and β -myrcene, these other chemicals have not been tested for carcinogenicity. The frequently-occurring α -pinene is however a mutagen. With time, more components are being tested and also found to be carcinogenic or allergenic. The worrying problems are that: when a person is exposed to an odour, the sensation appears to diminish over time as the person adapts to the odour, while sensory irritation occurs in an opposite manner; the higher the number of chemicals being combined, the lower their individual concentrations need to be to cause an adverse reaction and children may be more susceptible than adults.

Is the proposed new EU legislation justified in requiring the labelling of products containing known sensitizers above a certain level like: benzyl alcohol, benzyl salicylate, cinnamal, citral, coumarin, eugenol, geraniol, isoeugenol, anisyl alcohol, benzyl benzoate, benzyl cinnamate, citronellol, farnesol, limonene and linalool; the synthetics: amyl cinnamal, hydroxycitronellal, butylphenyl methylpropional, hexyl cinnamal, methyl 2-octynoate and alpha methyl ionone as well as the naturals, oakmoss and treemoss? Is it the occurrence of these components or their oxidised forms which are sensitizers? Is it the same whether the components are from natural essential oils or the synthetic equivalents, which may have different enantiomeric proportions and have different biological activities? Does it just make it more difficult for perfumers, flavour chemists and for aromatherapists, who may be severely restricted in what they can use? Or, does it benefit the consumer?

The essential oil of Australian Sandalwood (*Santalum spicatum*) – Effects of different samples on human physiology and subjective evaluation

Eva Heuberger^a, Valerie Gearon^b, Steve Birbeck^b, Gerhard Buchbauer^a
Eva.Heuberger@univie.ac.at

^a Institute of Pharmaceutical Chemistry, University of Vienna, Althanstraße 14, A-1090 Vienna, Austria

^b Mt Romance Australia Pty Ltd., Lot 2 Down Road, Mirambeen Park, Albany Western Australia 6330, Australia

The biological properties of the essential oil of Australian Sandalwood (*Santalum spicatum*), e.g. bacteriostatic (Beylier, 1979; Viollon *et al.*, 1996) and anti-inflammatory activity (Mt Romance Pty Ltd., 2002), are very similar to those of the essential oil of East Indian Sandalwood (*Santalum album*). Also, the use of Australian Sandalwood oil in traditional medicine very much resembles that of East Indian Sandalwood oil. Recently, our group was able to show that topical administration of East Indian sandalwood oil has relaxing and harmonizing effects on humans (Hongratanaworakit *et al.*, 2000). However, no such investigations have been carried out with Australian Sandalwood oil. Therefore, the aim of the present study was to elucidate the effects of Australian Sandalwood oil on human physiological parameters and self-evaluation after percutaneous absorption.

In our study three samples of Australian sandalwood oil with varying compositions of the same constituents (dissolved in peanut oil) and a placebo substance (peanut oil) were administered topically for 30 minutes to healthy human subjects while physiological data (skin temperature, skin conductance, breathing rate, blink rate, heart rate, muscle activity and blood pressure) were recorded. Inhalation of the substances was prevented by breathing masks. At the beginning and at the end of the application subjects were asked to rate their mental and emotional condition in terms of relaxation, vigour, calmness, attentiveness, mood, and alertness.

Statistical evaluation showed subtle changes of both physiological and subjective parameters after the administration of Australian Sandalwood oil in comparison to the placebo substance. Moreover, differences in effectiveness were revealed between the three samples of Australian Sandalwood oil.

Beylier M. F. (1979) Bacteriostatic *Perfum. Flavor.* 4: 23-25.

Hongratanaworakit T., Heuberger E., and Buchbauer G. (2000) *poster presentation at the 31st ISEO*, Hamburg, Germany.

Mt. Romance Pty Ltd. (2002) <http://www.mtromance.com.au/applic.html>, 2002-05-06.

Viollon C., Mandin D., and Chaumont J. P. (1996) *Fitoterapia*, 67: 279-281.

Exploring the biological activity and phytochemistry of South African medicinal aromatic plants used in traditional healing rites

A. M. Viljoen^a, S. F. van Vuuren^a, H. Albert^b, S. Hatch^b, B. Demirci^c, K. H. C. Başer^c, R. L. van Zyl^a, K. L. Lindsey^d, J. van Staden^d

viljoenam@therapy.wits.ac.za

^a Department of Pharmacy and Pharmacology, Faculty of Health Sciences, University of the Witwatersrand, 7 York road, Parktown 2193, South Africa

^b TB Laboratory, National Health Laboratory Service, Johannesburg, South Africa

^c Medicinal and Aromatic Plant and Drug Research Center (TBAM), Anadolu University, 26470-Eskişehir, Turkey

^d Research Centre for Plant Growth and Development, University of Natal Pietermaritzburg, P/Bag X01, Scottsville 3209

Despite the technological advancement in modern medicine, many people in South Africa rely on traditional healing practices. It is estimated that there are approximately 200000 traditional healers ('sangomas' and 'inyangas') in South Africa. South Africa also harbours one of the six floral Kingdoms in the world and this biodiversity has provided the traditional healer with an impressive 'natural pharmacy' from which plants are selected as ingredients to prepare herbal preparations. Aromatic plants are often used in these preparations in the form of infusions or aromatic plant material may be burnt and the smoke inhaled. Ten of the most widely used indigenous medicinal aromatic plants have been selected to investigate their chemical composition and biological activity in an attempt to establish a scientific basis for their frequent use. The decongestant and anti-microbial properties of essential oils explain the use of aromatic plants in treating respiratory ailments. Using time kill methods the minimum inhibitory percentage (MIP) has been determined for various indigenous species on pathogens responsible for respiratory ailments (e.g. *Klebsiella pneumoniae*, *Cryptococcus neoformans*, *Bacillus cereus*).

Although the action of essential oils on bacteria and fungi are well documented, almost no information is available on the antimycobacterial properties of mono- and sesquiterpenoids. A preliminary screening of anti-mycobacteria was performed using *Mycobacterium smegmatis*, a rapid-growing, non-pathogenic species, in an agar plate-based format. All oils showed inhibitory activity to varying degrees. Susceptibility of *Mycobacterium tuberculosis* H37rv to the oils was tested by the BACTEC radiometric method. This method measures metabolism of a radioactive substrate into ¹⁴CO₂ as a rapid indicator of growth of the organism. Inhibitory activity of the oils was reported at concentrations between 0.125% and 1.0%.

Many traditional uses of aromatic plants hint at possible anti-inflammatory activity, hence vernacular names such as 'fever tea' or 'fever-berry'. This information prompted us to investigate the *in vitro* inhibition of prostaglandin-synthesis. For the anti-inflammatory assay the essential oils were resuspended to a concentration of 1.0 % in DMSO and anti-inflammatory activity determined using the cyclooxygenase-2 radiometric assay.

Aromatic plants are documented for use in patients with "fever" or "flu-like" symptoms; which are the clinical symptoms used to describe a malaria infection. Thus, preliminary results of the *in vitro* antimalarial activity of the essential oils against a chloroquine-resistant *Plasmodium falciparum* strain will be discussed.

Composition and biological activities of essential oils of *Fagara zanthoxyloides* (dried fruits) from Cameroon as well as of *Syzygium cuminii* and *Syzygium travancoricum* (fresh leaves) from India

Leopold Jirovetz^a, Gerhard Buchbauer^a, Martin Benoit Ngassoum^b, Muhammed Shafi^c

leopold.jirovetz@univie.ac.at

^a University of Vienna, Institute of Pharmaceutical Chemistry, Austria

^b University of Ngaoundere, Department of Applied Chemistry, Cameroon

^c Calicut University, Department of Chemistry, Kerala, India

The composition of the essential leaf oil of *Fagara zanthoxyloides* (syn. *Zanthoxylum zanthoxyloides*, *Z. xanthoxyloides*, Rutaceae) from Cameroon and the essential fruit oils of *Syzygium cuminii* and *Syzygium travancoricum* (Myrtaceae) from India were analyzed by GC, GC-MS and olfactometry.

As main compounds of the essential *F. zanthoxyloides* oil α -pinene, *trans*- β -ocimene, citronellol, sabinene, myrcene, limonene, citronellyl acetate, α -terpinolene, α -phellandrene, geraniol, terpinen-4-ol, *p*-cymene, methyl citronellate and β -pinene were identified.

Pinocarveol, α -terpineol, myrtenol, eucarvone, muurolol, myrtenal, geranyl acetone, α -cadinol and pinocarvone as well as *trans*- β -ocimene, *trans*- β -caryophyllene, α -humulene and α -farnesene were found to be the dominating constituents of the *S. cuminii* and the *S. travancoricum* oil respectively.

The above mentioned essential oils were tested for their antimicrobial activities against *Bacillus cereus*, *Bacillus sphaericus*, *Bacillus subtilis*, *Corynebacterium glutamicum*, *Enterococcus faecalis*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Salmonella typhimurium* and *Staphylococcus aureus*.

The results of these antimicrobial activity testings and additional correlations of such effects of some single compounds (Jirovetz *et al.* 2000, 2001 and 2002; Lis-Balchin *et al.* 1998), identified in the essential oils of *F. zanthoxyloides*, *S. cuminii* and *S. travancoricum*, are discussed in this presentation.

Jirovetz L, G Buchbauer, A Stoyanova, S Metodiev (2000) *Sci Pharm* 68: 323-328.


Jirovetz L, G Buchbauer, MB Ngassoum, JJ Essia-Ngang, NL Tatsadjeu, O Adjoudji (2001) Proceedings of *ISEO* 32: 116.

Jirovetz L, G Buchbauer, MB Ngassoum, JJ. Essia-Ngang, NL Tatsadjeu, O Adjoudji (2002) *Sci Pharm* 70:93-99.

Lis-Balchin M, SG Deans, E Eaglesham (1998) *Flav Fragr J* 13: 98-104.

Posters Abstracts





P. 1. Volatiles from fruit and leaf oils of *Spondias* species growing in the Amazon

Eloisa Helena A. Andrade^a, Maria das Graças B. Zoghbi^a, José Guilherme S. Maia^b

eloisa@museu-goeldi.br

^aMuseu Emilio Goeldi, Departamento de Botânica, CP 399, 66040-170 Belém, Brazil

^bUniversidade Federal do Pará, Departamento de Química, 66075-900 Belém, Brazil

The species *Spondias mombin* L is originated from Amazon basin, though it is also known in the Northeast Brazil, Central America and West Indies. Its fruits are known as "taperebá" and "cajá-pequeno". Its wood barks, leaves and flowers are used in the local folk medicine. The fruits of *Spondias dulcis* Robinson are known as "cajarana" and the species was originated from the South Pacific. Now, it is widespread in many tropical countries, including Brazil. The fruits of *Spondias purpurea* L. are known as "umbu" and the species occurrence is the Northeast Brazil. All three *Spondias* fruits are used for preparation of ice creams, juices and aromatization of pies and liqueurs.

The volatiles of fruits and leaves of *S. dulcis*, *S. mombin* and *S. purpurea* were obtained by hydrodistillation and analysed by GC-MS. Individual components were identified by comparison of both mass spectrum and GC retention data with those existing in the data system libraries and cited in the literature (Maia *et al.* 2000, Adams 1995). The major constituents identified in the oils of *S. dulcis* were α -thujene (fruit: 44.8%), α -pinene (leaf: 62.9%) β -pinene (fruit: 19.7%; leaf: 13.8%) and α -terpineol (fruit: 10.2%). The oils of *S. mombin* were dominated by α -copaene (leaf: 27.1%), β -caryophyllene (fruit: 19.3%), δ -cadinene (leaf: 11.9%), (E, E)- α -farnesene (fruit: 11.1%), ethyl hexanoate (fruit: 10.1%), α -selinene (leaf: 9.4%), (Z)- β -ocimene, (fruit: 8.9%), caryophyllene oxide (leaf: 8.8%) and ethyl butyrate (fruit: 8.2%). The main components found in the fruit oil of *S. purpurea* were limonene (60.8%), Δ^3 -carene (7.7%) and terpinolene (4.1%).

Acknowledgements: supported by MCT-PPG7/European Community.

Adams RP (1995) Identification of Essential Oil components by Gas Chromatography/Mass spectroscopy, Allured, 495p, Illinois.

Maia JGS, Andrade EHA, Zoghbi MGB (2000) *Journal of food Composition and Analysis*: 13: 227-232.

P. 2.**Volatiles from *Lippia grandis* Schau.**

José Guilherme S. Maia ^a, Francisca Socorro N. Taveira ^b, Eloisa Helena A. Andrade ^c, Milton Helio L. da Silva^c,
Maria das Graças B. Zoghbi ^c

Gmaia@museu-goeldi.br

^a Departamento de Química, Universidade Federal do Pará, 66075-900 Belém, PA, Brazil

^b Departamento de Química, Universidade Federal do Maranhão, 65080-040 São Luis, MA, Brazil

^c Coordenação de Botânica, Museu Emílio Goeldi, 66040-170 Belém, PA, Brazil

The genus *Lippia*, belonging to the Verbenaceae, comprises about 200 species of herbs, shrubs and small trees growing in South and Central America and Tropical Africa. Of these, the chemical composition of the essential oils of about forty species has already been reported (Terblanché and Kornelius, 1996). The main compounds previously identified were limonene, *p*-cymene, α -pinene, camphor, β -caryophyllene, linalool and thymol. The species *Lippia grandis* Schau. is a shrub reaching 2.5 m, known as “erva-do-marajó” and occurring at natural fields of the Amazon Basin. Their leaf tea is used for liver ailments and menses disorders. Previously, we report a preliminary analysis of the essential oil of *L. grandis* (Silva *et al.* 1973). Now, we are reporting the volatile composition of five chemotypes of *L. grandis* collected at different locations and soil types at Southeast of Pará state (chemotypes A and B), Northeast of Roraima state (chemotypes C and D) and West of Maranhão State (chemotype E). Voucher specimens from chemotypes of *L. grandis* have been deposited in the herbarium of Museu Emílio Goeldi, in the city of Belém, Pará state, Brazil. The chemotype A was dominated by 1,8-cineole (22.2%), (E)-nerolidol (21.1%), α -pinene (17.4%) and β -caryophyllene (8.2%); The chemotype B by thymol (52.6%), methyl thymol (14.7%), *p*-cymene (21.1%) and γ -terpinene (8.9%); the chemotype C by thymol (67.4%); the chemotype D by carvacrol (21.2%), thymol (19.3%), *p*-cymene (14.0%) and β -caryophyllene (12.1%); and the chemotype E by 1,8-cineole (56.0%) and α -terpineol (13.7%).

Acknowledgements: Supported by MCT-PPG7/European Community.

Terblanché FC, Kornelius G (1996) *Journal of Essential Oil Research*: 8: 471-485.

Silva ML, Maia JGS, Mourão JC, Pedreira G, Marx MC, Gottlieb OR, Magalhães MT (1973) *Acta Amazonica*: 3: 41-42.

P. 3. Composition of the essential oil of *Hypericum glandulosum* Aiton grown on Madeira and its glandular structures

A. Cristina Figueiredo^a, José G. Barroso^a, Luis G. Pedro^a, Teresa Antunes^b, Isabel Sevinate-Pinto^b, Carlos Lobo^b

teresa.antunes@fc.ul.pt

^a Centro de Biotecnologia Vegetal, Dep. de Biologia Vegetal, FCL, C2, Campo Grande, 1749-016 Lisbon, Portugal

^b Centro de Biologia Ambiental, Dep. de Biologia Vegetal, FCL, C2, Campo Grande, 1749-016 Lisbon, Portugal

Nine *Hypericum* (Hypericaceae or Gutiferae) species grow on the Madeira archipelago, three of which are endemic (Tebbs 1994, Vieira 1992). *Hypericum glandulosum* Aiton (= *H. joerstadii* Lid) is a rare endemic species of Madeira, Porto Santo and Canaries Islands, currently known in Madeira as “malfurada” or “hipericão”. The plant is a small evergreen shrub up to 1m heights growing in rocky areas, particularly in the ravines in eastern Madeira. The infusion of the whole plant is used as a diuretic, to treat urinary bladder, kidney and liver diseases and against lithemia (Rivera and Obón 1995).

Scanning electron microscopic studies on vegetative and reproductive organs of *Hypericum glandulosum* revealed three different types of glandular structures: translucent glands or cavities containing essential oil, that occur exclusively at the leaves and two types of black nodules (internal and external) containing hypericin that are found at leaves calix and corola.

The aerial parts of *H. glandulosum* were collected during the flowering phase at Serra de Água, Madeira. The essential oils were isolated by distillation-extraction for 3h and analysed by GC and GC-MS.

The sesquiterpene hydrocarbons (46%) and non-terpenoid compounds (30%) were the main fractions of *H. glandulosum* oil, whereas the monoterpene fraction amounted to 15%. An unidentified sesquiterpene hydrocarbon (25%), β -caryophyllene (19%), *n*-undecane (13%) and *trans*-2-hexenal (11%) were the dominant components of this oil.

Rivera D, C Obón (1995) *Journal of Ethnopharmacology* 46: 73-93.

Tebbs BR (1994) *Hypericaceae*. In *Flora of Madeira*, eds Press JR & MJ Short, HMSO, London, p. 221.

Vieira R (1992) *Flora da Madeira*, Serviço Nacional de Parques, Reservas e Conservação da Natureza, Lisboa, p.65.

P. 4. Quality of supercritical fluid extract from *Levisticum officinale*

Á. Kéry^a, Cs. András^b, É. Lemberkovics^a, Á. Dzurillay^b, P. Apáti^a, M. Pálfi^a, B. Simándi^b

lembi@drog.sote.hu

^a Semmelweis University, Faculty of Pharmacy, Department of Pharmacognosy, Budapest, Hungary, Üllői út 26., H-1085

^b Budapest University of Technology and Economics, Department of Chemical Engineering, Budapest, Hungary, Műegyetem rkp. 3. K ép., H-1521

Lovage (*Levisticum officinale* Koch, Apiaceae) drugs and extracts are well known spices for food- and liqueur industry and are used medicinally for their stomachic, carminative and antirheumatic properties. As various chemical substances are involved both in the culinary value and the bioactivity to obtain high quality extracts of lovage has a great value. Supercritical fluids have been shown to exhibit several advantages in the extraction of natural products from plant matrices. The major effort on supercritical fluid extraction has to date been to evaluate whether or not it can replace traditional organic solvent extraction methods.

The effect of two factors (pressure and temperature of the extractor vessel) were investigated in the ranges of 150 - 450 bar and 40 - 60°C by designed experiments using CO₂. The 3² full factorial design was performed once, with three repetitions in the centre of the design. The extraction yield was used as dependent variable. It was concluded that increasing the pressure and/or temperature the extraction yield rises. The extracts contained the characteristic bioactive components of *Levisticum officinale*. Sixteen volatile constituents were separated by GC from which five were identified: butylphthalide, 2-butylidenephthalide, Z-ligustilide, 3-butyl-4,5-dihydrophthalide, E-ligustilide. The major component in each SFE extract was Z-ligustilide, which was always present in more than 70 %. The characteristic phthalide derivatives were also proved by HPLC and UV spectra. The effect of the extraction pressure for the yield of fatty acids and β-sitosterol was highly significant. Their satisfactory recovery was achieved if the pressure and temperature were higher than 300 bar and 50°C respectively. Fatty acid composition of SFE extracts was typical for Apiaceae species. HPLC analysis proved the presence of some minor compounds as coumarin derivatives beside the main Z-ligustilide. By comparison of retention times and on line UV spectra with authentic samples the linear furanocoumarin imperatorin, isoimperatorin, begapten and the simple coumarin umbelliferon were identified. Efficiency and selectivity of SF CO₂ and traditional extractions support the view that qualitatively new products can be gained in function of various phytochemicals, which may be beneficial in the industrial and therapeutical use.

Acknowledgements: The work was supported by FKFP-0169, ETK-T-08-158/99 (OTKA T 0300344)

Bruneton J (1995) *Pharmacognosy Phytochemistry*, Medicinal Plants, Intercept Ltd, Andover.

Kéry Á, E Rónyai, B Simándi, É Lemberkovics, T Keve, A Deák, S Kemény (1999) *Chromatographia* 49: 503.

Modey WK, DA Mullholland, MM Raynor (1996) *Phytochem. Analysis* 7: 1.

P. 5. An unique *n*-propyl sesquiterpene from *Eryngium creticum* L. (Apiaceae)


Nahla A. Ayoub^a, Mahmoud A. M. Nawwar^b

ayoub.n@link.com.eg

^aDepartment of Pharmacognosy, Faculty of Pharmacy, Ain-Shams University, Cairo, Egypt

^bDepartment of Phytochemistry, National Research Centre, Cairo, Egypt

The sesquiterpenes cadinanes and muurolanes are known to bear isopropyl substituent connected to their perhydronaphthalene cores. Recently, the muurolane derivative, muurol-9-en-15-al, (**1**) has been isolated from *Eryngium maritimum* (Kubeczka, 1998). In a continuation of our previous studies (Kubeczka, 1988, 1999 and Ayoub, 2001) on the essential oils of *Eryngium* we describe herein the isolation, from *E. creticum*, of a new perhydronaphthalene derivative with a *n*-propyl substituent at its position no. 1. It was identified as 1,2,4a,5,6,7,8,8a-octahydro-4-methyl-1-propyl-naphthalene-7-carbaldehyde, (**2**). The compound possesses an unique sesquiterpene carbon skeleton for which we suggest the name eryngane (**3**). The isolated terpenoid (**2**) itself is therefore, renamed, erng-9-en-15-al. We were able, in addition, to isolate the new natural methyl ketone eicos- 8,11-dien-18-ol-2-one, (**4**). Structures of compounds (**2** & **4**) were established by conventional methods of analysis and confirmed by DEPT, COSY, HMQC and HMBC as well.



Ayoub N, Kubeczka KH 2001 [Abstract] 32th ISEO, Wroclaw, Poland.

Kubeczka K.H, Ayoub N, Grande M, Torres P 1998 [Abstract] 29th ISEO, Frankfurt.

Kubeczka KH, Ayoub N, Saleh M, Nawwar M 1999 [Abstract] 30th ISEO, Leipzig.

P. 6. Chemical constituents of the essential oil of *Achillea pachycephala* Rech.f., a species endemic to Iran

Shahrzad Bamasian^a, Abdolhossein Rustaiyan^a, Shiva Masoudi^b

abam@hamgam.com

^aDepartment of Chemistry, Islamic Azad University, North Tehran Branch, Tehran, Iran

^bDepartment of Chemistry, Islamic Azad University, Central Tehran Branch, Tehran, Iran

Achillea is a large genus belonging to the family *Asteraceae*. It is native to Europe and Asia, and can also be found in North America. Nineteen species of the genus *Achillea* are found in Iran, among which seven are endemic (Mozaffarian, 1996). The genus *Achillea* is well known for medicinal properties, in particular, *A. millefolium* is used as a popular antipyretic, because of its analgesic and anti-inflammatory properties (Chandler *et al.*, 1982).

Chemical studies on several *Achillea* species have resulted in the isolation of sesquiterpene lactones, phenolic and acetylenic compounds. Several authors have studied the composition of the oils of different *Achillea* species (Rustaiyan *et al.*, 1998, 1999).

Water distilled essential oil from the herbal parts of *A. pachycephala* Rech. f., which is endemic to Iran, was analysed by means of GC/MS in combination with retention indices. Thirty-three components were characterized in the oil representing 96.9% of the total compounds detected with 1,8-cineole (27.6%) and camphor (27.4%) as major constituents.

The oil was characterised by a large amount of oxygenated monoterpenes (78.7%), and a lesser amount of monoterpene hydrocarbons and sesquiterpenes (15.7% and 2.5%, respectively).

Chandler RF, SN Hooper, MJ Harvey (1982) *Econ. Bot.* 36: 203-223.

Mozaffarian V (1996) *A Dictionary of Iranian Plant Names*, p. 12 Farhang Moaser, Tehran, Iran

Rustaiyan A, H Komeilizadeh, S Masoudi (1998) *J. Essent. Oil Res.* 10: 207-209.

Rustaiyan A, S Masoudi, M Yari (1999) *J. Essent. Oil Res.* 11: 19-20.

P. 7.

Scents from Amazon aromatic plants

Lauro E. S. Barata, Karen F. Discola

lbarata@iqm.unicamp.br

Institute of Chemistry UNICAMP (State University of Campinas, Campinas, São Paulo Brazil).

Over the last 20 years, human incursion into the Brazilian Amazon resulted in the degradation of more than 600000 km² of the natural forest, an area bigger than France. Rosewood, threatened by a half-century of predatory extraction, contributes with ca. 5000 fallen trees per year. Rosewood oil of international commerce currently originates from a destructive process, from an endangered species in which the wood is chopped and extracted by steam distillation (Barata, 2001a).

In consequence of environmental regulations and economic factors, the Brazilian Rosewood industry are progressively pushing away towards establishing plantations of this specie. To face this problem, the author established collaboration with commercial rosewood distillers to establish plantations system trials. In consequence, since the year 2000, ca. 20000 Rosewood trees have been planted in consortium with thousands of others aromatic native plants (Table 1). The goal is to develop new essential oils derived from leaves by coppicing the trees.

Research & Development of the essential oil of the Aromatic Plants is in progress, under a project that shall impact Environmental, Economical, Social and Technological areas in Amazon. A pessimistic estimation considers that 30ha of rosewood plantation shall produce ca. 1000 kg of essential oil under the coppice of 3-5 year-old trees, representing a reasonable cash flow to local producers.

Experiences with rosewood plantations in Amazon have been published (Barata, 2001b).

This presentation reviews the results achieved of recent research on the potential for a new essential oils from Amazon Aromatic Plants. Steam distillation has been made using laboratory extractors and chemical composition was done by GC-MS chromatography.

Table 1. Rosewood and other Aromatic Plants species cultivated by consortium in Amazon

Common Name	Scientific Name	Family	Class	# Trees	Years ¹
Rosewood	<i>Aniba rosaeodora</i>	Lauraceae	Tree	20000	5
Macacaporanga	<i>Aniba fragrans</i>	Lauraceae	Tree	1000	5
Preciosa	<i>Aniba canelilla</i>	Lauraceae	Tree	2100	5
Puxuri	<i>Licaria puchuri-minor</i>	Lauraceae	Tree	NR	5
Andiroba ²	<i>Carapa guianensis</i>	Meliaceae	Tree	5900	4
Copaiba	<i>Copaifera</i> spp.	Caesalpiniaceae	Tree	720	50
Cumaru	<i>Dipteryx odorata</i>	Fabaceae	Tree	980	10

1. Number of years to adult stage. 2. Not an aromatic plant. Produces insecticide substances.

Barata LES (2001a) *Proceedings of IFEAT International Conference*, Buenos Aires.

Barata LES and May, P. (2001b) *Economic Botany* (submitted).

P. 8. Activity of the essential oils of citronella (*Cymbopogum winterianus*) and basil (*Ocimum basilicum*) on the dry wood termite *Cryptotermis brevis*

A. C. Sbeghen, V. Dalfovo, L. A. Serafini, N. M. Barros

n.barros@terra.com.br

Instituto de Biotecnologia e Departamento de Ciências Biológicas - Universidade de Caxias do Sul, Caxias do Sul- RS- Brazil

The termite *Cryptotermis brevis*, commonly known as “dry wood termite”, is usually found in home environments, where it attacks furniture and wooden built structures. The conventional control method of termites is based on the use of chemicals, which are highly toxic for humans and other living beings. For this reason, our work aimed at verifying the effect of citronella and basil essential oils on *C. brevis*, analysing the consumption of substrate (filter paper) impregnated with the oils, and termite mortality. The oils were assayed on concentrations ranging from 94 to 471 $\mu\text{g}/\text{cm}^2$. In all tests, 1 mL of essential oil solutions was applied to the dry filter paper disks, using blank and acetone-impregnated filter paper disks as controls. 50 worker termites were placed in each Petri dish. Daily observations of the use of substrate and insect mortality were carried out during a 25-day period. After that period, the determination of the effective dose that produces a 50% reduction in the feeding (ED_{50}) and the median lethal concentration (LC_{50}) were carried out by the PROBIT method. Higher feeding inhibition effects were observed when basil essential oil was used, resulting in ED_{50} (56 $\mu\text{g}/\text{cm}^2$). Citronella oil also showed anti-feeding activity, but its ED_{50} was 167 $\mu\text{g}/\text{cm}^2$. The toxicity effect of the essential oils was measured by the mortality rate of the termites. The toxic effect of the oils was LC_{50} 262 $\mu\text{g}/\text{cm}^2$ for basil and 290 $\mu\text{g}/\text{cm}^2$ for citronella.

Aknowledgments: Funded by CNPq, FAPERGS and UCS

P. 9. Repellence and toxicity of wax of orange oil in the alternative control of drywood termites

V. Dalfovo^a, E. Fronza ^a, A. C. Sbeghen ^a, C. D. Frizzo ^b, N. M. Barros ^a

n.barros@terra.com.br

^a Instituto de Biotecnologia e Departamento de Ciências Biológicas – Universidade de Caxias do Sul, Caxias do Sul RS- Brazil

^b Aripê Citrus Ltda., RS 124 Km 1.2, 95780-000 Montenegro -RS, Brazil

Cryptotermis brevis is the most common drywood termite. It has been a wide geographic distribution, occurring in all tropical and subtropical areas around the world. In Brazil, it occurs from south to northeast, being the second most important plague among termites in southeast of Brazil. Searching alternatives to prevent and control the infestation by termites, we can emphasise natural products derived from plants. Despite this, the objective of this study was to evaluate the repellence and toxicity of a natural product, which constitutes a residue in the citrus industry, resulting from the cold-pressing process: the wax of orange oil, *Citrus sinensis* (L.) Pers. Another product generated by the cold-pressing process in the citrus industry, wax of lime oil “Tahiti”, *Citrus aurantifolia* (Christ.) Swingle, was used as a parameter for comparison of the tested products. The bioassays were carried out in Petri dishes. Two wooden blocks (*Pinus* sp.) (3.0 x 3.0 x 0.5 cm) and 25 termites were placed in each Petri dish, being one of them treated with wax and other used as control (without treatment). The bioassays were done in triplicate and the results were evaluated for 30 days. The results showed that both waxes tested caused repellence to the insects with an index of mortality of 6.0%.

Aknowledgments: Funded by CNPq, FAPERGS and UCS

P. 10. The essential oil of *Pentapleura subulifera* Hand.-Mazz.

K. H. C. Başer^a, B. Demirci^a, N. Tabanca^a, Z. Aytaç^b, M. Ekici^b

khcbaser@anadolu.edu.tr

^a Medicinal and Aromatic Plant and Drug Research Centre (TBAM), Anadolu University, 26470-Eskişehir, Turkey

^b Faculty of Science and Letters, Gazi University, 06500 Ankara, Turkey

Pentapleura subulifera Hand.-Mazz. (Lamiaceae) is a monotypic genus growing in South Eastern Turkey, Northern Iraq and Western Iran (Davis 1982).

Water-distilled essential oil from the aerial parts of *Pentapleura subulifera* Hand.-Mazz., was analysed by GC and GC/MS. Fifty three components were characterised representing 95.9% of the oil with 1,8-cineole (39.0 %) and T-cadinol (11.1 %) as main constituents.

Davis PH (1982), *Flora of Turkey and the East Aegean Islands*. Vol. 7, p. 313, University Press, Edinburgh.

P. 11. Chemical and microbiological studies of main compounds of *Thymi herba et extracta*

A. Bazylko^a, J. Stefańska^b, H. Strzelecka^a

oklyzab@farm.amwaw.edu.pl

^aDepartment of Pharmacognosy, Faculty of Pharmacy, University of Medicine, ul. Banacha 1, 02-097 Warsaw, Poland

^bDepartment of Pharmaceutical Microbiology, Faculty of Pharmacy, University of Medicine, ul. Oczki 3, 02-007 Warsaw, Poland

Thyme (*Thymus vulgaris* L., *Lamiaceae*) is listed to aromatic plants containing essential oil (1.0 – 2.5%). The essential oil of *Thymus vulgaris* L. would normally be assigned to the aromatic alcohol-rich group. The main phenolic components of the volatile oil of thyme are thymol and carvacrol in varying amounts (thymol 0.2 - 66.5 %, carvacrol 0.2 - 72.0 %). These compounds determine antibacterial (Agnihotri, 1996; Didry, 1993) and fungicidal (Zambonelli 1996) actions of thyme and its volatile oil. On the other hand, in some countries a fundamental part of many galenicals are thyme extracts, and the studies of them were the aim of our investigations.

In the first part we determined the content of essential oil, thymol, luteolin and rosmarinic acid, in the raw material and in a liquid and dry extract. The second part has included an analysis of antiseptic activity of: thyme oil, thymol, luteolin, rosmarinic acid, thyme ethanolic extract and infusion, as well as of its commercialised liquid and dry extract.

The material for research was supplied by PHYTOPHARM, Klęka, Poland. The plant material had been collected in June 1998, at the flowering stage, near Piotrków Trybunalski. It was dried at 35°C. The liquid extract (*extractum fluidum*) was obtained from *Thymi herba* according to DAB X, while the dry extract is *extractum spirituosum siccum*. The essential oil was obtained by steam distillation, using a Deryng apparatus, and was dehydrated with anhydrous sodium sulphate and stored at 4-6°C. The content of thymol was determined by GC method (Shimadzu GC-17A). The contents of luteolin and rosmarinic acid were determined by TLC-photodensitometry (Shimadzu CS-9301 PC). Microbiological actions were investigated against five bacteria and yeast, *Candida albicans*, where a disc diffusion method was used.

Agnihotri S., Vaidya A.D.B. (1996) *Indian J. Exp. Biol.* 34(7): 712-715

Didry N., Dubreuil L., Pinkas M. (1993) *Pharmazie* 48: 301-304

Zambonelli A., Zechini D'Aulerio A., Bianchi A., Albasini A. (1996) *J. Phytopathol.* 144: 491-494

P. 12. Genetic diversity and variability of essential oils from Tunisian *Mentha pulegium* L. populations

Najeh Ben Fadhel, M. Mkaddem, M. Boussaid

najehbenfadhel@lycos.fr

Institut of Applied Sciences and Technologies, Department of Biology, Laboratory of Plant Biotechnology. Centre Urbain Nord, B.P. 676. 1080 Tunis Cedex. Tunisia.

Pennyroyal (*Mentha pulegium*, Lamiaceae) was exploited for a long time for the quality of its essential oils, used in medicine, cosmetics, perfume and fragrance (Simon, Chadwick and Craker 1984). The overexploitation of natural populations has led to great genetic erosion (Voirin *et al.* 1999). Measures of conservation of the species and the elaboration of improvement programs were necessary to ease natural populations.

The analysis of the genetic diversity of populations, by isoenzymatic markers, and the detection of variations in chemical composition of the essential oils could help to the selection of interesting cultivars (Lewis and Crawford 1995; Kang *et al.* 1998).

Fifteen wild *Mentha pulegium* populations, prospected in Tunisia, were surveyed. Eight isozyme systems (PGI, PGM, GOT, ICD, ADH, MDH, LAP and 6-PGD) were analysed by horizontal electrophoresis to access the genetic diversity.

The variation of essential oils composition, within and among populations, was analysed by gas chromatography (GC) and (GC/ MS). Eight major components (pulegone, menthone, carvone, isomenthone, 1-pinene, 1-limonene, menthol...) were considered.

The species maintained high levels of polymorphism within populations. Particular alleles: Adh1-a, Adh1-b, Adh1-d, Icd1-a, Mdh1-a, Mdh3-a and Pgi2-a emerged as a genetic markers for some populations, relating of their geographical localisation. The essential oils composition varied between individuals within the same site and between sites.

There is a positive relationship between the distribution of the isozymic markers (unique alleles) and the variability of the chemical components. Isozyme markers may play an important role in rapid identification of appropriate chemotypes, to promote their exploitation in industrial fields. They could be also used to facilitate conservation and improvement actions in *Mentha pulegium*.

Kang SS, J Noguchi, KB Park, M Gi. Chung (1998) *Nordic Journal of Botany* 18: 581-587.

Lewis PO, DJ Crawford (1995) *American Journal of Botany* 82: 141-149.

Simon, JE. Chadwick, LE. Craker (1984) *Herbs: An indexed bibliography* 1971-1981. The scientific Literature on selected herbs, and aromatic and medicinal plants of temperate zone. Archon Books, Hamden, CT.

Voirin B, C Bayet, O Faure, F Jullien (1999) *Phytochemistry* 50: 1189 - 1193.

P. 13. Genetic diversity, mating system and essential oils in *Rosmarinus officinalis* L., *Myrtus communis* L. and *Mentha pulegium* L.

M. Boussaïd, Y. Zaouali, C. Messaoud, A. Ben Salah, N. Ben Fadhel

mhdboussaïd@rnu.insat.tn

Institute of Applied Sciences and Technologies, Department of Biology, Laboratory of Plant Biotechnology, Centre Urbain Nord, B.P. 676. 1080 Tunis Cedex. Tunisia.

The majority of medicinal and aromatic plants is used to extract essential oils which are exploited in various fields (traditional medicine, perfumery, pharmacy, insecticides, fungicides). However, the bulk of the treated material is still harvested wildly. The over exploitation of spontaneous species has led to the rarefaction and the extinction of many of them (Vinay, 1996; Khiari and Boussaïd, 2002).

Systematic cultivation of species can therefore be introduced in order to conserve biodiversity, protect threatened taxa and promote the culture of interesting varieties.

The quality of essential oils has to depend on the whole process beginning with the selection of material and ending with the final product proposed to the consumer.

Plant selection programs require first the assessment of the genetic diversity and that of the variability of the essential oils in natural populations taking into account the species mating system. These studies may help in the detection and improvement of performant genotypes and contribute to elaborate conservation strategies (Soltis and Soltis, 1991).

Three medicinal and aromatic species: *Rosmarinus officinalis*, *Myrtus communis* and *Mentha pulegium* growing spontaneously in Tunisia were surveyed in this study.

12 to 15 populations collected in different geographical sites represented each species. We analyzed the polymorphism of eight isozyme markers by starch gel electrophoresis. The variation of fourteen essential oil components has been studied by gas chromatography. The mating system in each species was also analyzed in order to interpret the genetic variability.

High levels of genetic variation were observed within populations and unique alleles were detected in *Mentha pulegium* and *Myrtus communis* in relation with geographical sites (Ben Fadhel, 2002; Messaoud, 2002). However, a substantial differentiation between populations was observed, which is a result of low gene flow due to habitat destruction.

Despite their high morphological similarities, individuals within populations exhibited high levels of variation in the composition of their essential oil. This may indicate an important genotype richness, which could be used to promote appropriate varieties.

Isozyme and chemical variations observed in populations may be due to the mating system. In fact, species analyzed were allogamous. So, hybridizations between individuals occurred. It is important to take this parameter into account to ensure efficient selection and conservation.

The presence of particular alleles, revealed by electrophoresis, was in relation with that of some essential oil components. So, isozyme markers could be considered as valid predictors of the chemical composition of plants.

Ben Fadhel N (2002) Proceedings 13th congress of ATSB (Tunisie)

Khiari D, M Boussaïd (2002) Tropicultura n°99F139 AMI/154 mjd.

Messaoud CH (2000) Master FST Bizerte, Tunisie 80 p.

Soltis PS, DE Soltis (1991) Aliso 13: 215-223

Vinay T (1996) Newsletter of the IUCN, 2.

P. 14. Biological activities of basil (*Ocimum basilicum* L.) essential oilBiljana Božin^a, Neda Mimica-Dukić^b, Milan Matavulj^a, Nataša Simin^b, Ružica Pavkov^c

dukaned@eunet.yu

^a Institute of Biology and Ecology, ^bInstitute of Chemistry, Faculty of Natural Sciences, University of Novi Sad, Trg D. Obradovica 2-4, YU-21000, Novi Sad, FR Yugoslavia^c "HEMOFARM" Vrsac., Pharamceutical Company, Vrsac, FR Yugoslavia

Oxygen-derived free radicals and lipid peroxidation (LP) are widely accepted as major contributors to the ethiology of atherosclerosis and its chronic disorders, including coronary heart disease, stroke and ischemic dementia (Knight, 1995). Synthetic antioxidants such as butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA) have been widely used for many years to retard lipid oxidation. However, concern about the safety of synthetic antioxidants together with consumer's preference for natural products has resulted in increased research on natural antioxidants. Since ancient times, aromatic plants have been added to different types of food to improve the flavour. Furthermore, it was recognized that the deterioration and rancidity of food was also prevented, what is primarily due to the antioxidant and antimicrobial ability of plant constituents. The essential oils of many aromatic plants are widely used in cosmetic, pharmaceutical and foodstuff industries, due to their strong antimicrobial activity. Recently, for some terpenes (geraniol, tocotrienol, perillyl alcohol, β -ionone, *d*-limonene) have been found to be effective, non-toxic dietary antitumor agents and hold promise as a novel class of antitumor drugs for human cancer (Karlson et al, 1996).

In this study, the results of the free radical scavenging capacity (RSC) and the antibacterial activity of basil (*Ocimum basilicum*) essential oil are reported.

The essential oil was isolated from dried aerial parts of basil (cultivated in northern part of Vojvodina) by hydrodistillation. The composition of essential oil was evaluated by GC-MS. RSC was assessed using DPPH (2,2-diphenyl, 1-picrylhydrazyl)-method (Kirby and Schmidt, 1997) and measuring OH[•]-radical induced degradation of 2-deoxyribose. Antibacterial activity of essential oil (20 and 50% solution in *n*-hexane) was tested against 13 bacterial strains, by standard antibiogram test.

The examined essential oil strongly reduced free radical formation of DPPH (IC₅₀=0.285 μ g/ml). The inhibition of 2-deoxyribose degradation was 47.1% for pure oil, 50.0% for 50% solution and 44.1% for 20% solution of essential oil. The RSC measured by DPPH-method was dose-dependent. The most effective antibacterial activity of 20% solution of essential oil was on *Shigella sonnei* (multiresistant strain from Public Health Institute, Faculty of Medicine, Novi Sad) and of 50% solution on *Bacillus subtilis* ATCC 10707. Both examined solutions also exhibited inhibitory activities on multiresistant strains of *E. coli*, *Salmonella typhi*, and *Shigella sonnei*.

The obtained results suggest that basil (*Ocimum basilicum*) could serve not only as flavour agent, but also as a safe food supplement in preventing deterioration of foodstuff products. Furthermore, consumption of food prevented with natural basil essential oil is expecting to prevent the risk of many free radicals dependent diseases.

Karlson J, AK Borg, R Unelius, MC Shoshan, N Wilking, U Ringborg and S Linder (1996) *Anticancer Drugs* 7(4): 422-429.

Kirby AJ and RJ Schmidt (1997) *J. Ethnopharmacol.* 56: 103-108.

Knight JA (1995) *Ann. Clin. Lab. Sci.* 25: 111-121.

P. 15. Antimicrobial activity of the *Vanillosmopsis erythropappa* (DC.) Sch.Bip. essential oil

G. B. Oliveira ^a, A. M. A. Nascimento ^a, E. Chartone-Souza ^a, I. C. P. Fortes ^b, L. F. Moreira ^c, M. G. L. Brandão ^c

branlins@dedalus.lcc.ufmg.br

^a Instituto de Ciências Biológicas, ^b Instituto de Ciências Exatas, ^c Faculdade de Farmácia
Universidade Federal de Minas Gerais, Av. Olegário Maciel, 2360. 30180-112 Belo Horizonte. Brazil

Vanillosmopsis erythropappa Schultz-Bip. (*Eremanthus erythropappus* (DC.) Mac Leish/ Asteraceae) is a Brazilian tree used for construction purposes or firewood. The essential oil from its trunk wood has been widely explored by chemical and pharmaceutical industry as source of α -(-)-bisabolol. This compound, typically associated with camomile, is used in cosmetic products to provide soothing effect due to its anti-inflammatory activity. In this work we have investigated the antimicrobial activity of the essential oil of this plant against drug-resistant microorganisms. The trunk wood of *V. erythropappa* was collected from mature trees growing in Barão de Cocais (Minas Gerais) and extracted with hexane. The extract was submitted to antimicrobial assay and showed activity against *Staphylococcus aureus* at concentration of 2%. Quantitative composition of the active oil was calculated from GC/ MS peaks, which showed the presence of several compounds mainly α -(-)-bisabolol (43,6%), β -bisabolene (27,5%), α -humulene (18,0%) and α -bisabolol oxide B (17,0%). Column chromatography of the active oil on silica gel with toluene - ethyl acetate (93:7) yielded 10 different fractions, which were assayed for antimicrobial activity. The results revealed that a fraction constituted by β -bisabolene and α -humulene are highly effective against *Staphylococcus aureus*.

P. 16. Breath-by-breath nosespace analysis while eating banana by Proton-Transfer-Reaction Mass-Spectrometry

Dagmar Mayr^a, Tilmann Märk^a, Werner Lindinger^a, Hugues Brevard^b, Chahan Yeretzyan^b

hugues.brevard@rdls.nestle.com

^a Institut für Ionenphysik, Leopold-Franzens-Universität, Technikerstr. 25, 6020 Innsbruck, Austria

^b Nestlé Research Center, P.O. Box 44, CH-1000 Lausanne 26, Switzerland

The aroma (odour) of food products is related to volatile organic compounds (VOCs) that are released from foods and reach the olfactory epithelium in the upper part of the nose. VOCs can reach the olfactory epithelium from two distinct directions. Either they are sniffed through the nose, via the orthonasal pathway and enter the nostrils from the front. Alternatively, they reach the olfactory receptors through the oral cavity and the pharynx, via the retronasal pathway. When food is eaten, the retronasal path leads to olfactory perception.

Here we report on the in-mouth aroma while eating banana. By analysing banana we have chosen to explore a solid food whose aromatic composition is known. Headspace and nosespace air are sampled using Proton-Transfer-Reaction Mass-Spectrometry (PTR-MS) technique. Nosespace air is collected via two glass tubes fitted into the nostrils. Then, the air is introduced into the drift-tube of the PTR-MS device. Headspace differences between unripe and ripe banana are shown. Furthermore, the time-intensity evolution of selected aroma compounds, the phenomenon of swallowing and the after-odour, will be discussed.




Figure: In-vivo breath-by-breath analysis of nosespace with PTR-MS while eating banana. Three selected masses are shown. The top trace shows mass 89. Two compounds contribute to this ion mass intensity: Ethyl acetate and isopentanol. The middle trace represents Isobutyl acetate, which has a typical banana note and is the single most important key flavour compound of banana. The bottom trace represents isopentyl acetate, which also has a banana odour note. On the left, the traces are shown while eating an unripe banana, while the right traces correspond to volatiles released in the mouth with eating a ripe banana.

P. 17. Quantitative structure activity relationship for synthetic sandalwood fragrance compounds

Assia Kovatcheva ^a, Gerhard Buchbauer ^a, Peter Wolschann ^b

gerhard.buchbauer@univie.ac.at

^a Institute of Pharmaceutical Chemistry, University of Vienna, Althanstrasse 14, A-1090, Vienna, Austria

^b Institute of Theoretical Chemistry and Molecular Structural Biology, University of Vienna, Währingerstrasse 17, A-1090, Vienna, Austria

QSAR is a modern throughout preferred approach to drug design. The main concept of QSAR techniques is to quantify the relationship between chemical structure and biological activity of potential drug candidates. Choices as to the set of compounds, the molecular descriptors, the mathematical form of the relationship between the biological activity and the descriptors as well the statistical method used to analyse the data are most important considerations in QSAR practice.

The lack of information of 3D-structures of the human odorant receptor and the better understanding of physicochemical mechanism of olfaction cause the need of development of new theoretical models for odorant molecules.

East Indian sandalwood oil is an appreciated constituent of perfume compositions because of its excellent odor properties. Its limited availability coupled with the high price is the reason of the incessant search for synthetic products with similar scent. Among all of these substitutes, the group of α -campholenyl derivatives achieved an odor most similar to the natural sandalwood. The unflagged interest in invention of new sandalwood odor substituents and particularly the synthesis of optically pure products are the foundations of providing this research.

A QSAR study for a set of α -campholenyl derivatives with known configuration of their chiral centers was performed. The relationship between the biological activity and structural, electronic and lipophilic properties of the molecules is described by the best model, which was found after applying Multiple Linear Regression. The model is evaluated by cross-validation and successfully applied on a set of test compounds for prediction of their odor thresholds.

The results help to expand the understanding of the interaction between sandalwood molecules and the still unknown odorant receptor site.

P. 18. SPME applied to the study of volatiles from Rosemary submitted to stress and ageing

Fabiana Gomes^{a*}, Tânia M. Pizzolato^a, Eduardo Cassel^b, Maria do Carmo R. Peralba^a, Luciana A. Serafini^c

*fabiana_rs@yahoo.com.br

^a Instituto de Química - UFRGS, Bento Gonçalves 9500, Porto Alegre – RS, 91501-970, C.P. 15003, Brazil

^b Faculdade de Química - PUCRS, Ipiranga 6681, Porto Alegre – RS, 90619-900, Brazil

^c Instituto de Biotecnologia – UCS, Francisco Getúlio Vargas 1130, 95070-560, Caxias do Sul – RS, Brazil

This work reports on the profile of volatiles from Rosemary leaves submitted to stress and ageing. Rosemary was grown in Campestre da Serra, South of Brazil. Solid Phase Micro Extraction (SPME) with 100 μm polydimethylsiloxane fiber and GC/MSD/IT were used to obtain the chromatographic profile of Rosemary volatiles.

Chromatographic analysis showed different results when in *in vivo* Rosemary leaves were compared with those, which had already been cut. The composition of volatiles was monitored through 28 days. The leaves were drying at room temperature (25 °C) and the plant was submitting it to stress conditions as without water. The compounds limonene, α -terpinene, γ -terpinene, terpinolene, are not present in chromatograms of *in vivo* Rosemary, suggesting that these compounds are encapsulated in reservoir bags where essential oils are accumulated. Other compounds like α -bisabolene, δ -2-care-4-ol have been detected only after plant cutting.

This monitoring allows the ideal time for Rosemary storage after harvesting to be determined. Geraniol, ocimene, verbenol, myrtenol, geranial, δ -3-caren-2-ol, myrtenyl acetate, bornilene and other less volatile compounds disappear within 120 hours after harvesting. The terpenes α -pinene, camphene, β -pinene, β -terpinene, terpinolene, 1,8-cineole, γ -terpinene and caryophyllene remain for 28 days after harvesting.

Results obtained so far, allow us to conclude that some compound appear only in stress conditions, such as lack of water or environment change. Echinenone and α -bisabolene were detected shortly after a transplant to a ceramic pot and did not appear afterwards, indicating that their presence is associated with plant non-equilibrium. The increase of quantity of the Rosemary essential oil and the appearance of β -pironene, *p*-mentone and *p*-cimene was observed when the plant was left without water for 3 days.

P. 19. Influence of seasonality and leaf position on leaf biometry and essential oil yield of *Lippia alba* (Mill) N. E. Br. ex Britt & Wilson

Dulce M. Castro^a, Lin C. Ming^a, Márcia O. M. Marques^b, Silvia R. Machado^c

dulce@saloonet.com.br

^a Department of Vegetable Production - FCA/Unesp / Botucatu - SP

^b Center of Genetics, Molecular Biology and Fitoquímica of the Agronomic Institute of Campinas (CGBMFq - IAC)

^c Departamento de Botany-I.B. /Unesp/Botucatu

This work relates variations in leaf biometry and in yield of essential oil of *Lippia alba* as a function of the season of the year and position of the leaves on the branch. Thus the parameters of seasonal variation (spring, summer, autumn, and winter) were considered concerning location of leaves in three parts of the branch (apical, median, and basal). Extractions of essential oil were accomplished through hydrodistillation. For biometric analysis of the leaves, samples were collected of the third median of the blade. The assay was carried out at São Manuel Experimental Farm (Unesp/Botucatu/Brazil), Department of Plant Production. The cuttings were maintained in a greenhouse, and after 60 days transplanted into a definitive location with a 1.0 X 0.60m space utilized for each. The plots are composed by four seasons (Summer'98/99, Autumn'99, Winter'99 and Spring'99), a completely randomized block was employed for the experimental design using 6 replications and different leaf positions (apical, median, basal). Samples were taken from 12 plants. On each plant, 10 leaf pairs were collected from basal (nodes 1-9; old leaves from 7.5-9.0 cm² in area), middle (nodes 10-18; mature leaves from 9.0-9.5 cm² in area) and apical regions (nodes 19-28; young leaves from 6.0-6.5 cm² in area). According to plot, all plant samples were collected during different seasons: summer'98/99 (December/January); autumn'99 (March), winter'99 (June); and spring'99 (September). Tukey (5% ratio) was used to evaluate essential oil yields with respect to season and leaf position. Histological sections were prepared from samples obtained from medial region of the leaf blade taken from the three plant positions. The samples were fixed in Karnovsky solution (12), dehydrated in an alcohol series, and embedded in glycolmethacrylate resin. Sections (8- μ m in thickness) were stained with 0.05% toluidine blue in acetate buffer, pH 4.7 (15), and permanently mounted in Entelan synthetic medium. Quantitative data are based on 24 individual counts for each treatment (station X leaf position). A completely randomized block was employed for the experimental design using 12 treatments and 6 replications. The statistical design was performed at 5% significance. Tukey and Scheffé tests were used to evaluate the means of each station and leaf position. Essential oil extraction was performed by hydrodistillation of fresh leaves (100 g) in a Clevenger type apparatus for 1h and the extract analyzed by GC/MS. An analysis of variance (Tukey 5% ratio) was calculated for the percentage values of the oil yield at each station and leaf position. Data shown in the tables are mean values of four replications. The highest yield of oil was obtained in summer, in leaves located in the apical part of the branch; in these leaves during the same season, the total thickness of the blade was greater, resultant of greater development of the palisade parenchyma and epidermal cells of the adaxial face.

P. 20. Seasonal variation, leaf position on the stem and drying effect on biomass production and essential oil yield and composition of *Lippia alba* (Mill) N. E. Br. ex Britt & Wilson (Verbenaceae)

Dulce M. Castro^a, Lin C. Ming^a, Márcia O. M. Marques^b, Sílvia R. Machado^c

dulce@saloonet.com.br

^a Department of Vegetable Production - FCA/Unesp / Botucatu - SP

^b Center of Genetics, Molecular Biology and Fitoquímica of the Agronomic Institute of Campinas (CGBMFq - IAC)

^c Departamento of Botany-I.B. /Unesp/Botucatu

According to the ethnobotanical data, *Lippia alba* (Mill.) N. E. Br. ex Britt. & Wilson, Verbenaceae, commonly known as “erva cidreira brasileira” or “falsa Melissa”, is one of the most important plants widely used by the population, mainly through the leaf tea, due to its antispasmodic, sedative and stomachic activities among others (Ming, 1992). This work aimed to study the best harvest time for biomass production and essential oil yield and composition of the different plant parts (apical, medial and basal) related to the seasonal variation (spring, summer, autumn and winter). The oils were analysed by GS/MS (Shimadzu, QP-5000) and the components identified by comparison of their mass spectra with those of a MS computer library and/or published in the literature as well as by co-injection with authentic standards. Applying phytochemical tests over fresh and dry mass, the chemical component percentages of essential oils were calculated and identified as follows: citral (neral and geranial), β -myrcene, β -caryophyllene and β -elemene. These results have also been found by Gomes (1993), Correa (1992), in the same species collected at different sites of Brazil. The secretory structures and chlorophyllaceous parenchyma holding essential oils were identified and quantified using microchemical tests through light (LM) and scanning electron microscopy (SEM). Essential oils were extracted by hydrodistillation with a Clevenger apparatus for both fresh and dry mass obtained in field and lab conditions respectively. The seasonal experiment in the field was carried out making possible the compoundable analysis employment for different harvest times in relation to different parts of plant (apical, medial and basal) as well as possible interactions between harvest time and parts of plant. This kind of multivariate analysis is possible, as there is a correlation among the three-plant parts and essential oil yield. The statistical design and methodology were performed at 5% significance in agreement with Morrison (1976). A completely randomised block design was employed for the drying experiment with five treatments and four replications. Tukey and Scheffé tests were used to evaluate the means of each temperature and analyse the differences between the two environments (natural and artificial) respectively, both at 5% significance.

Correa, C.B.V. (1992) Contribuição ao estudo de *Lippia alba* (Mill) N.E.Br. ex Britt & Wilson - erva cidreira. Revista Brasileira de Farmácia, 73: 57-64.

Gomes, E.C., L.C. Ming, E.A. Moreira, O.G. Miguel (1993) Constituintes de óleo essencial de *Lippia alba* (Mill.) N.E. Br. (Verbenaceae). Revista Brasileira de Farmácia 74: 29- 32.

Ming,L.C. Influência de diferentes níveis de adubação orgânica na produção de biomassa e teor de óleos essenciais de *Lippia alba* (Mill)N.E.Br. Verbenaceae. (1992). Curitiba, 206p. Dissertação(Mestrado) em Ciências Biológicas, Universidade Federal do Paraná.

Morrison,D.F. (1976) Multivariate statical methods. 2^{ed}. Tokyo, Kogakusha. 220p.

Silva, J.B. Novo método de extração de óleo essencial de *Lippia citriodora* (K.) do Rio Grande do Sul. (1979) Tribuna Farmacêutica, Curitiba, 4: 96- 8.

P. 21. Composition of the essential oil of *Teucrium lusitanicum*C. Cavaleiro^a, L. Salgueiro^a, G. Miguel^b, A. Proença da Cunha^a

cavaleir@ff.uc.pt

^a Laboratório de Farmacognosia / CEF, Faculdade de Farmácia, Universidade de Coimbra, Rua do Norte 3000 Coimbra, Portugal^b Faculdade de Engenharia de Recursos Naturais, Universidade do Algarve, Campus de Gambelas, 8000 Faro, Portugal

Continuing our research on the composition of the essential oils of *Teucrium* species, we now report on the composition of the essential oil of *Teucrium lusitanicum* Schreber [*T. polium* subsp. *vicentinum* (Rouy) D. Wood] from Portugal. This species grows spontaneously in W and S Iberian Peninsula and NW Africa (Valdés *et al.* 1987). In Portugal occurs only in W Algarve.

The aerial parts of plants from four populations were collected during the flowering period in Sagres and Cabo de São Vicente (Algarve - Portugal) and submitted to water distillation for essential oils isolation.

The oils were analyzed by GC and GC-MS, using two fused silica capillary columns with different stationary phases (polyethylenoglycol and polymethylsiloxane). Components were identified by GC retention indexes and mass spectra.

Sixty-five compounds were identified, representing more than 90% of the total oils. Although qualitative compositions are similar, important quantitative differences were found. (Table 1).

Table 1. Major compounds (>5% in at least one sample) and grouped components of the essential oil of *T. lusitanicum*

Compound	% in samples				Average(Min-Max) %
	A	B	C	D	
α -Pinene	8.2	2.5	0.8	8.5	5.0 (0.8-8.5)
Sabinene	9.6	3.2	2.1	7.7	5.7 (2.1-9.6)
β -Pinene	10.5	4.7	2.5	11.9	7.4 (2.5-11.9)
β -Myrcene	7.3	2.8	2.5	4.1	4.2 (2.5-7.3)
Limonene	11.5	1.2	3.0	2.3	4.5 (1.2-11.5)
Terpineol-4	1.9	2.7	5.5	2.8	3.2 (1.9-5.5)
Germacrene D	3.1	5.3	6.0	1.0	3.9 (1.0-6.0)
δ -Cadinene	2.0	4.1	5.3	7.3	4.7 (2.0-7.3)
Elemol	6.1	11.2	12.0	2.6	8.0 (2.6-12.0)
T-Cadinol	5.4	5.2	5.5	6.2	5.6 (5.2-6.2)
α -Cadinol	4.2	4.9	4.7	9.1	5.7 (4.2-9.1)
Grouped components					
Monoterpene hydrocarbons	49.8	15.7	12.4	36.2	28.5 (12.4-49.8)
Oxygen containing monoterpenes	6.8	11.0	15.6	10.9	11.1 (6.8-15.6)
Sesquiterpene hydrocarbons	14.3	30.1	32.2	20.2	24.2 (14.3-32.2)
Oxygen containing sesquiterpenes	20.5	34.2	32.9	25.8	28.4 (20.5-34.2)
Other compounds	1.2	1.6	1.9	0.6	1.3 (0.6-1.9)
Total identified	92.6	92.6	95	93.7	93.5 (92.6-95.0)

Valdés B, Talavera S, Fernández-Galiano E (eds.). Flora Vasculare de Andalucía Occidental, Ketres Editora, S.A. Barcelona, 1987

P. 22. Essential oil from leaves of *Cryptocarya moschata* Nees (Lauraceae): constitution and population variability

Marcelo Telascrea^a, Carla C. de Araújo^a, Márcia O. M. Marques^b, Roselaine Falcalani^b, Alberto J. Cavalheiro^a

albjcava@iq.unesp.br

^a NuBBE – Núcleo de Bioensaios, Biossíntese e Ecofisiologia de Produtos Naturais, Instituto de Química – UNESP, Departamento de Química Orgânica, CEP 14800-900, Araraquara-SP, Brazil

^b Centro de Genética, Biologia Molecular e Fitoquímica - Instituto Agrônomo de Campinas, CEP 13020-902, Campinas - SP, Brazil

Cryptocarya moschata, the most common and numerous *Cryptocarya* species from Brazil, is a tree with 20-30 meters high and a very important alimentary source to mastofauna, mainly primates, as *Brachyteles arachnoides* (“mono-carvoeiro” or “muriqui”), and cracid birds, as *Pipile jacutinga* (“jacutinga”), that eat their leaves and fruits. Previous phytochemical studies revealed the occurrence of several styrylpyrones in barks (Cavalheiro and Yoshida, 2000). Flavonoid glycosides and styrylpyrones are found in leaves, and intraspecific variability on contents of styrylpyrones in these organs was reported (Nehme *et al.* 2002). In previous studies about essential oils from *Cryptocarya* species, the occurrence of linalool was described in leaves of *C. aschersoniana* and *C. moschata*, and β -myrcene, 1,8-cineol and stereoisomeric linalool oxides were found only in the former (Naves *et al.* 1963).

To complement the phytochemical investigation on *C. moschata* for further ecophysiological studies, a detailed description of the components of essential oils from leaves was made, using GC-MS, retention index, chromatographic fractionation and ¹³C NMR spectroscopy. Seventy compounds were identified, with predominance of isomeric sesquiterpenes (MW 204). The more intense peaks observed on the chromatogram obtained by GC-MS were attributed to germacrene-D, β -caryophyllene, bicyclogermacrene, α -copaene, δ -cadinene, spathulenol, α -humulene, linalool, caryophyllene oxide and β -cubebene. In addition, forty trees of three populations of *C. moschata* from Atlantic Rain Forest of São Paulo State – Brazil, were analyzed by GC-FID, in order to assess intraspecific variability on essential oil components. The cluster analysis led to identification of two groups with different essential oil profile, characterized by quantitative differences of (on the) main components. Intra and interpopulational variability were verified. Essential oil group with predominance of β -caryophyllene (19-34%), germacrene-D (14-25%) and bicyclogermacrene (10-25%), named CGB group, is more frequent at South populations, and essential oil group with predominance of germacrene-D (26-41%), β -caryophyllene (9-20%), and bicyclogermacrene (9-18%), named GCB, is more frequent at North population, indicating a clinal variation in quantitative profile of essential oil components from leaves of *C. moschata*. These results are concordant with intraspecific chemical variability found to styrylpyrones from leaves of the same individuals of *C. moschata* (Nehme 2001; Nehme *et al.* 2002).

Acknowledgements: Financial support: FAPESP and CNPq

Cavalheiro AJ; Yoshida, M. (2000) *Phytochemistry* 53 (07): 811-819.

Naves, Yves-René, et al. (1963). *Helvetica Chimica Acta*, 46: 1056-9. (Please give the complete reference authors)

Nehme CJ *PhD Tesis*. Instituto de Química – UNESP, Araraquara-SP, Brazil, 2001.

Nehme CJ, Moraes PLR, Cavalheiro AJ (2002) *Bioch. System. and Ecol.* 30, (6): 613-616.

P. 23. Contribution to the composition of the essential oils from herb and fruits of Greek coriander

P. S. Chatzopoulou^a, S. T. Katsiotis^b

xatzlin@yahoo.gr

^a National Agricultural Research Foundation, Agricultural Research Centre of Macedonia & Thrace, Department of Aromatic and Medicinal Plants, P.O. Box 60458, Thessaloniki 57001, Thessaloniki, Greece

^b Department of Pharmaceutics, School of Pharmacy, Aristotle University of Thessaloniki, 54124, P.O.Box 1589, Thessaloniki, Greece

Seeds of *Coriander sativum* L (fam. Umbelliferae), well known as an important ingredient of many ethnic foods, produce a commercial valuable essential oil utilized widely in beverages, condiment and in the perfumery industry, as well as in pharmaceuticals.

The essential oils of Greek coriander herb and seeds were studied, considering the influence of the maturity stages, the comminution of the seeds and the distillation time applied, on their yield and quality.

The main constituents of the oils, analysed by GC/MS, were 2-decenal for the herb, and linalool for the seeds. Significant variability in the essential oil content and composition were noticed in the seeds samples from different ontogenetic stages, resulting in an increase of the linalool content from 44.5 to 82.5%, throughout the whole maturity period.

The treatment of the seeds affected also significantly the obtained essential oils. The intact seeds yielded higher linalool amounts (max 82.5%) than the comminuted ones (max 71.7%). The latter were richer in oxygenated monoterpenes when shorter distillation times were applied.

P. 24. *Foeniculum vulgare* Mill. from different origin: variation of essential oil content and main components

H. G. Chung^a, S. M. Kim^b, N. S. Seong^a, E. Nemeth^c

^a haegon@rda.go.kr

^aDivision of Industrial Crop of National Crop Experiment Station Rural Development Administration, 209 Seodun-dong Kwansunku Suweon, Republic of Korea

^bCollege of Industrial Science, Konju National University, Yesan Chung-Nam Republic of Korea

^cDepartment of Medicinal and Aromatic plants Faculty of Horticultural Sciences, Szeint Istvan University, Villany ut 29-35 Budapest, Hungary

The aim of our investigation of Korean and Hungarian origin fennel was to examine the variation of essential oil content and its main components by hydrodistillation and capillary GC method with standards during the ontogenesis and plant growing seasons from 1999 to 1999. The essential oil contents of the shoots (2nd year old) and that of the roots (2nd and 3rd year old) exhibited a higher accumulation level in the Hungarian population at the leaf-rosette-stage. The essential oil content varied according to vegetation years and phenological phases and the maximum essential oil values were measured (10.0-12.8% in the Hungarian and 7.1-11.8% in the Korean taxon) in the two-year old plants of seeds. We observed the maximal essential oil content of seeds at a certain phase, when the dry matter content was between 22 and 26%. From this stage, the increase of dry matter content and the decrease of essential oil content have a significant linear correlation ($r > 0,5$) with each other in both taxa. Anethole (60.9-82.2%) proved to be the main component of the over-ground organs, while dillapiol (53.9-92.6%) was the major compound of the roots in each year and in both populations. Intra-specific differences concerning the reproductive organs are reflected first of all in levels of anethole (maximum values by 12-50% higher in the Hungarian) and of α - and β -pinenes (more in Hun. By 0.9-3.2%). Korean population can be characterised as a high anethole - low methyl chavicol chemoform of the anethole chemovariety (fenchone <15%; anethole > 68%; methyl chavicol <3,2%) respectively.

P. 25. Extraction of *Satureja montana* L. essential oil with supercritical carbon dioxide

A. P. Pereira^a, J. Burillo^b, J. S. Urieta^b, A. C. Figueiredo^c, J. G. Barroso^c, R. L. Mendes^a, J. A. Coelho^a,
A. M. F. Palavra^a

jcoelho@deq.isel.ipl.pt

^a Centro de Química Estrutural, IST, Lisbon, Portugal

^b Dep. Química-Física, Universidad - Servicio de Investigación Agroalimentario, Zaragoza, Spain

^c Centro de Biotecnología Vegetal, Dep. de Biología Vegetal, FCL, C2, Campo Grande, 1749-017 Lisbon, Portugal

Aromatic plants are important raw materials for the perfumery, pharmaceutical and food industries. Essential oils from these plants are traditionally obtained by hydrodistillation and solvent extraction, although these techniques have several disadvantages, such as the heat instability of essential oils and the difficulty to remove the solvent from the extract.

With supercritical fluid extraction (SFE) it is possible to obtain solvent-free extracts and to avoid the degradation of thermally labile components (Palavra *et al.*, 1993, 1995, 1999). Therefore, the natural odour and flavour of the initial material are maintained.

Satureja montana L. is an aromatic plant that grows in arid, sunny and rocky habitats. The corresponding essential oil, obtained by hydrodistillation, can reach about 1.8% (w/dw). The principal use of this plant is in the treatment of stomach ailments due to its carminative and desiccative properties.

Flowers and leaves of *Satureja montana* L., grown at Servicio de Investigación Agroalimentario, Zaragoza, Spain, were used as matrices for supercritical CO₂ extractions, followed by two stage fractional separations. The oil was obtained with a flow apparatus, with an extraction vessel (1L), and two separators (0.27L) in the pressure and temperature ranges of 90 to 100 bar and 40 to 50 °C, at the selected separations conditions, and analysed by GC and GC-MS. The essential oil was dominated by the monoterpene fraction, carvacrol (56-71%) being its main component. The amount of the sesquiterpene components was always lower than 2%.

The most convenient conditions of extraction and separation, to obtain the oil from this plant, are discussed.

Bruno T, CAN Castro, JF Hammel, AMF Palavra (1993) Supercritical Fluid Extraction of Biological Products. In Recovery Processes for Biological Materials, eds. JF Kennedy, JMS Cabral, J. Wiley & Sons, Chichester.

Mendes RL, JP Coelho, HL Fernandes, IJ Marrucho, JMS Cabral, JM Novais, AF Palavra (1995) "Applications of Supercritical CO₂ Extraction to Microalgae and Plants", J. Chem. Tech. Biotechnol., 62: 53-59.

Reis-Vasco EMC, JP Coelho, AF Palavra (1999) "Comparision of Pennyroyal Oils Obtained by Supercritical CO₂ Extraction and Hydrodistillation", Flav. Frag. J., 14: 156-160.

P. 26.**Madeira Laurel essential oils**

M. C. Costa^a, P. Castilho^b, F. Venâncio^a, C. Camacho^b, A. Rodrigues^a

ceu.costa@mail.ineti.pt

^a INETI-DTIQ, Instituto Nacional de Engenharia e Tecnologia Industrial, Estrada do Paço do Lumiar, P-1649-038 Lisboa, Portugal

^b Universidade da Madeira, Dept. de Química, Campus Universitário da Penteada. P-9000-390 Funchal, Portugal

Laurus azorica is a *Lauraceae* endemic of the Macaronesia (Press *et al.*, 1994) confined to the archipelagos of Azores, Canaries and Madeira. In Madeira, *Laurus azorica* is the most abundant tree in the thermo-Mediterranean (altitude 14-700m in the South coast, 0-300m in the North coast) and meso-Mediterranean (700-1500m in the South coast and 300-600m in the North coast) bioclimatic steps.

Essential oils may be obtained from different morphological parts of the plant; the better studied are the oils from leaves and berries. A fixed (lipidic) oil is expressed from the ripe berries and used in traditional medicine as anti-infective, anti-rheumatic, blood depurative, haemostatic and apoplexy preventive (Rivera *et al.*, 1995) This oil is much richer in volatiles (11.7%) than any other known vegetable oils (<5%).

Comparative studies on leaves and berries, complemented with results on the chemical composition of essential oils from stems and the fixed oil are presented. Three different locations in Madeira Island were used for plant material collection: Ponta do Pargo, Ribeiro Frio and Chão de Louros.

Volatiles were obtained by hydrodistillation with a Clevenger type apparatus, during 4 hours. Yields ranged from 0.19% in stems to 9.1% in the fixed oil. Over this lipidic product the hydrodistillation was left to proceed for two other (uninterrupted) periods of 4 hours until no more volatiles were obtained; two new portions (1.7 and 0.8% yield, respectively) were thus isolated. These were found to be very poor in monoterpenes but richer on sesquiterpenes and oxygenated compounds. Sample collection at regular intervals over a long time distillation proved to be a useful method for the crude separation of compounds for further NMR analysis.

Studies over the composition of foliar essential oils are in good agreement with literature results, showing a dominance of the monoterpene fraction, with α -pinene (13-15 %) as the major compound, followed by 1,8-cineole (9-10%) and β -pinene (7-8%).

The ratio hydrocarbons/oxygenated compounds presented a significant variation with collection location; correlation of this ratio with botanical diversity is still to be proved.

The phenylpropanoid elemicin was found in amounts varying from 2 to 9%, with the most expressive values on the stems of trees from Chão de Louros (Camacho *et al.*, 2001). This compound was previously detected (1.7%) in *Laurus azorica* leaves from Madeira Island although it was absent from samples collected in Azores archipelagos (Pedro *et al.*, 2001).

Chemical composition of the first fraction obtained by distillation from the expressed oil correlates well with the volatiles obtained from the ripe berries, suggesting that even though the oil production involves the boiling of the berries, the process does not affect the aroma of the product.

Camacho C, A Rodrigues, F Venâncio, P Castilho, MC Costa 2001 XV *Encontro Galego Português de Química*, La Coruña, Spain.

Pedro LG, PAG Santos, JA da Silva, A C Figueiredo, JG Barroso, S G Deans, A Looman and J J Scheffer 2001 *Phytochemistry* 57 (2): 245-250

Press JR, MJ Short 1994 (ed) *Flora of Madeira*, The Natural History Museum, London, , p.102.

Rivera D, C Obón 1995 *J. Ethnopharm* 46: 73-93.

P. 27. Compositon of the essential oil of *Micromeria juliana* (L.) Bentham ex Reichenb. (*Lamiaceae*)

M. Couladis^a, O. Tzakou^a, V. Slavkovska^b, B. Lakusic^b, R. Jancic^b

kouladi@pharm.uoa.gr

^a Department of Pharmacognosy and Chemistry of Natural Products, School of Pharmacy, University of Athens, Panepistimiopolis Zografou, 15771 Athens, Greece

^b Institute of Botany, Faculty of Pharmacy, University of Belgrade, Vojvode Stepe 450, 11000 Belgrade, Yugoslavia

Micromeria juliana (L.) Bentham ex Reichenb. belongs to the section *Eumicromeria* (Boissier, 1879), tribe *Mentheae*, subfamily *Nepetoideae* (Cantino *et al.*, 1992).

This species grows in the Mediterranean region, spreading westwards to S.E. France, and in C. Portugal (Chater and Guinea, 1972). In Yugoslavia, *M. juliana* is distributed in Kosovo and Montenegro (Silic, 1979).

The object of our investigation was the essential oil composition of *M. juliana* collected from three different localities in Western Montenegro: Moraca canyon, Cijevna canyon and Mt Orjen (c. 500 m).

The samples were gathered in the flowering period. The essential oils were extracted from dry aboveground parts of the plants by hydrodistillation. The yield of essential oils was lower than 0.1%.

The analyses of the oils were carried out using GC/MS. The identification of the compounds was based on comparison of their Kovats indices (KI), their retention times (RT) and mass spectra with those obtained from authentic samples and/or the MS library (Adams, 1995).

The main component in all oils was caryophyllene oxide. However, the contribution of certain other major components varied.

The population from Moraca canyon had a high content of caryophyllene oxide (18.1%), carvacrol (18.1%), *o*-cymene (10.8%) and *p*-cymene (5.3%). Traces of isomenthone and pulegone were found.

In the essential oil from the Cijevna canyon population there was a considerable presence of caryophyllene oxide (20.4%), (E)-caryophyllene (7.1%), α -cadinol (6.6%) and borneol (6.3%). Isomenthone, pulegone, thymol and carvacrol were found in traces, while no *o*-cymene was detected in the essential oil of this population.

The Mt. Orjen population contained essential oil rich in caryophyllene oxide (15.9%), isomenthone (10.1 %), pulegone (8.1%), thymol (7.3%), (E)-caryophyllene (5.6%), *allo*-aromadendrene (5.3%) and germacrene D (5.2%). There were traces of borneol and carvacrol, while there was no presence of *o*-cymene.

The species *M. juliana* is similar to other species of *Eumicromeria* section, according to essential oil characteristics.

Supported by Serbian Ministry of Science, Technology and Development.

Adams RP (1995) Identification of Essential Oil Components by Gas Chromatography/Mass Spectroscopy. Allured Publishing Co, Illinois, USA.

Boissier E (1879) Flora Orientalis, Basileae, Genève & Lugduni, 568-575.

Cantino PD, RM Harley, SJ Wagstaff (1992) Genera of Labiatae: Status and classification. In: Harley RM, Reynolds T. (eds.), Advances in Labiatae Science, Royal Botanic Gardens, Kew, 511-522.

Chater AO, E Guinea (1972) *Micromeria* Bentham. In: Tutin TG, VH Heywood, NA Burges, DM Moore, DH Valentine, SM Walters, DA Webb (eds.), Flora Europaea, Cambridge University Press, Cambridge, 3: 167-170.

Silic C. (1979) Monographie der Gattungen *Satureja* L., *Calamintha* Miller, *Micromeria* Bentham, *Acinos* Miller und *Clinopodium* L. in der Flora Jugoslawiens, Zemaljski muzej BiH, Sarajevo, 209-219.

P. 28. Chemical Composition of the essential oil of *Teucrium polium* ssp. *capitatum* L. (Labiatae) from Corsica (France)

Stéphanie Cozzani^a, Jean-Marie Desjobert^a, Alain Muselli^a, Antoine François Bernardini^a, Félix Tomi^b,
Joseph Casanova^b

tomi@vignola.univ-corse.fr

^a Université de Corse - Equipe Chimie des Produits Naturels UMR CNRS 6134 - Quartier Grossetti - 20250 Corte, France

^b Université de Corse - Equipe Chimie et Biomasse, UMR CNRS 6134, Route des Sanguinaires, 20000 Ajaccio, France

The genus *Teucrium* exhibit more than 300 species, which were distributed around the world (1). Among them, *T. polium* (L.) has been used in folk medicine (2). Therefore, several studies reports the identification of components extracted from *T. polium* (2-3). Concerning the chemical composition of the essential oil of *T. polium*, three compositions were reported in the literature (4-5). In Corsica, the essential oil obtained by steam distillation of the aerial parts from *Teucrium polium* ssp. *Capitatum*, grown wild in the stony ground of the island, takes an interest in perfume industries. To our knowledge, the chemical composition of the essential oil of *T. polium* subsp. *capitatum* was never reported in the literature.

In the present work, the essential oil of *Teucrium polium* ssp. *capitatum* L. from Corsica, was investigated using combination of capillary GC, GC/MS and Carbon-13 NMR spectroscopy after fractionation by column chromatography. More than eighty compounds, accounting for 81% of the total amount were identified. The main components were α -pinene (19.9%), p-cymene (7.9%) and β -pinene (5.0%).

α -Agarofurane, β -dihydroagarofurane, 4 α -hydroxydihydroagarofurane, 3 β ,4 β -oxydo-agarofurane and 6,7-bisepoxycalamenene-seco, were identified as minor components and to our knowledge, this is a first report on these compounds in the essential oil of *Teucrium polium*.

1. Coste H., Flore descriptive et illustrée de la France, de la Corse et des contrées limitrophes, 3 volumes, Librairie Blanchard, Paris, 139-140, 1909.
2. Bello R., Calatayud S., Moreno L., Beltrán B., Primo-Yúfera E., Esplugues J., Effects on arterial blood of the methanol extracts from different *Teucrium* species, *Phytotherapy research*, 1997, **11**, 330-331.
3. Hassan M., Al-Hazimi G., Ghulam A. Miana, The diterpenoids of *Teucrium* species (Part II), *Journ. Chem. Soc. Pak.*, 1993, **15** (3), 215-226.
4. Hassan M. M. A., Muhtadi F. J., Al-Badr A. A., GLC-Mass Spectrometry of *Teucrium polium* oil, *J. of Pharmaceutical Sciences*, 1979, **68** (6), 800-801.
5. Çakir A., Emini Duru M., Harmandar M., Ciriminna R., Passannanti S., Volatile constituents of *Teucrium polium* L. from Turkey, *J. Essent. Oil Res.*, 1998, **10**, 113-115.

P. 29. Effects of *Thymus vulgaris* L. (Thyme) essential oil as an *in vivo* dietary supplement on chicken intestinal microflora

D. E. Cross, K. P. Svoboda, K. Hillman, R. McDevitt, T. Acamovic
D.Cross@au.sac.ac.uk

Scottish Agricultural College, Auchincruive Estate, Ayr, Scotland, KA6 5HW, UK

Thyme oil possesses antimicrobial properties *in vitro*, due to the presence of phenolic terpenes in *Thymus vulgaris* L. (Dorman & Deans, 2000; Helander *et al.*, 1998). Terpenoids may also act as nutraceuticals against certain antibiotic-resistant bacteria (Nascimento *et al.*, 2000), by reducing the bacterial load. This paper describes the effects on the gut microflora when thyme oil was added to chicken diets in the presence or absence of supplementary dietary enzymes.

Using a factorial design, 240 female broiler chickens (Ross 308) were reared on litter from 0-42 days, arranged as 10 birds per pen and 6 pen replicates of 4 dietary treatments. All birds were reared on a basal wheat/soya bean meal ration (ME=12.88MJ kg⁻¹; CP=190.6g kg⁻¹), either with or without a thyme oil supplement at 5g kg⁻¹. Both diets were then fed with and without an enzyme at 0.5g kg⁻¹ (xylanase=2776 IU & β-glucanase=117 IU). Caecal contents were sampled at 19 & 42 days, serially diluted and plated within 12 hrs onto MRS, MacConkey No.3, Wilkins Chalgren, Perfringens OPSP, charcoal cefaperazone deoxycholate (CCDA) and Slanetz-Bartley agar media (Oxoid Ltd., UK). After transformation to log₁₀, all counts were analysed by GLM in Genstat Release 5.

Table Effects of *T. vulgaris* supplementation in the diet on chicken intestinal microflora at 19 days

Treatment	Inclusion	Lactic Acid Bacteria	Coliforms	Total Anaerobes	Cl. perfringens
Oil	+ve	10.03	8.88 ^a	10.63	2.54
	-ve	9.47	9.94 ^b	10.73	2.62
	s.e.d.	0.352	0.451	0.320	1.062
	Sig	NS	0.028	NS	NS
Enzyme	+ve	9.58	9.56 ^a	10.99	1.53 ^a
	-ve	10.03	8.76 ^b	10.95	4.06 ^b
	s.e.d.	0.264	0.338	0.240	0.797
	Sig.	NS	P<0.05	NS	P=0.006
N		3	3	3	3

All counts are expressed as log₁₀ colony forming units per gram sample. Significance comparisons are separate for each treatment factor and are based within each column

The trial birds were compromised by a coliform infection, unrelated to treatment from study days 1-10, with 7.4% mortality. At 19 days, thyme oil reduced the numbers of coliforms, whereas the enzyme reduced *Cl. perfringens*, indicating separate modes of action. No significant effects were observed at 42 days due to either treatment (not shown), and no treatment interactions were present.

These results indicate the protective potential of dietary thyme oil and enzymes in birds with an immature bacterial flora. Bacterial count reduction will aid recovery, representing reduced potential either for re-infection or the transfer of infection to other birds. The bacterial count variations masked any effects due to treatment at 42 days. Inclusion of more thyme oil in diets may compensate for volatilisation (only 60% was recovered), but may prove unpalatable, thus terpene mixtures may be more effective. In conclusion, terpenes show potential for dietary inclusion, although more detailed studies are required to elucidate the length of the protective period.

The authors thank SEERAD for funding, Danisco Animal Nutrition for enzymes & Ian Nevison for his statistical advice

Dorman, H.J.D., S.G. Deans (2000) *Journal of Applied Microbiology* 88: 308-316

Helander, I.M., H-L. Alakomi, K. Latva-Kala, T. Mattila-Sandholm, I.Pol, E.J. Smid, L.G.M. Gorris, A. von Wright (1998) *Journal of Agriculture & Food Chemistry* 46: 3590-3595

Nascimento, G.G.F, J. Locatelli, P.C. Freitas, G.L. Silva (2000) *Brazilian Journal of Microbiology* 31: 247-256

P. 30. Seasonal variation of the leaf essential oil from two populations of *Pittosporum undulatum* Vent. grown in the Portuguese mainland

Nicolau J. Ferreira, Inês G. Meireles de Sousa, Tiago Cunha Luís, António José M. Currais, A. Cristina Figueiredo, José G. Barroso, Luis G. Pedro

tiago.c.luis@netcabo.pt

Centro de Biotecnologia Vegetal, Dep. de Biologia Vegetal, FCL, C2, Campo Grande, 1749-016 Lisbon, Portugal

The genus *Pittosporum* comprises about 200 species distributed throughout the world. Three species of this genus are described for Portugal: *P. coriaceum* Dryander ex Aiton [Mocano, Pitosporo], *P. undulatum* Vent. [Árvore do incenso] and *P. tobira* (Thunb.) W. T. Aiton.

P. undulatum is an evergreen 4-8m tree with white-creamy flowers in loose umbellate cymes, orange capsule when ripe and wavy edged glabrous leaves. Nowadays considered an alien weed, it was introduced in most countries because of its ornamental value, with its perfumed white-creamy flowers, orange capsules and fast-growing leaf canopy (Palhinha 1946, Goodland and Healey 1996). *P. undulatum* is abundant in several hill forest areas in the Portuguese mainland and it is one of the most frequent species in wet hill forest near Lisbon.

The aerial parts of *P. undulatum* were collected fortnightly during six months at the Parque de Saúde de Lisboa. Two populations were considered: pruned (pP) and non-pruned trees (npP). The essential oils were isolated from deep-frozen aerial parts by hydrodistillation to estimate the oil yields, and by distillation-extraction to determine their percentage composition, and analysed by GC and GC-MS.

The essential oils isolated from the leaves of pruned and non-pruned trees of *P. undulatum* were obtained in yields ranging from 0.05% to 0.15% (v/w) and from 0.03% to 0.20% (v/w), respectively.

Monoterpenes were dominant in all oils (>85%), the monoterpene hydrocarbons constituting the main fraction of both populations (pP >65%, npP >67%). Table 1 gives the percentage ranges for the main components. Sesquiterpenes occurred in a relative amount <8% in pP and <10% in npP, bicyclogermacrene being the dominant component of this fraction, Table 1.

Table 1: Percentage range of the main components from the two populations of *Pittosporum undulatum* studied.

Percentage range	pP	npP
α -Pinene	4 - 8	5 - 11
Sabinene	13 - 36	13 - 25
Myrcene	4 - 6	3 - 5
α -Terpinene	3 - 6	3 - 5
Limonene	12 - 36	14 - 37
γ -Terpinene	4 - 10	4 - 7
Terpinen-4-ol	8 - 24	10 - 16
Bicyclogermacrene	2 - 3	2 - 4

Acknowledgements: The authors gratefully acknowledge the permission of the Parque de Saúde de Lisboa for collection of plant material.

Palhinha R. T. (1946) Plantas aromáticas de Portugal, *Brotéria* 15: 97-113.

Goodland T., J. R. Healey (1996) The invasion of Jamaican mountain rainforests by the Australian tree *Pittosporum undulatum*, School of Agricultural and Forest Science, University of Wales, Bangor, UK.

P. 31. Chemical composition and antimicrobial activity of the essential oil of *Eucalyptus globulus* Labill., Myrtaceae, from Montenegro

Biljana Damjanovic^a, Jovanka Damjanovic^b, Vladimir Zivkovic^c

bibana@cg.yu

^a Faculty of Metallurgy and Technology, Univ. of Montenegro, Cetinjski put bb., 81000 Podgorica, Montenegro, Yugoslavia

^b Institute for Public Health - Montenegro, Ljubljanska bb., Podgorica, Montenegro, Yugoslavia

^c Center for Ecotoxicological Researches of Montenegro, Put Radomira Ivanovica 2, Podgorica, Montenegro, Yugoslavia

Essential oils, due to their bioactive components, are indeed promising to be used as effective antibacterial, antifungal and antioxidant agents. With the growing interest in the use of essential oils in both food and pharmaceutical industries, a systematic examination of plant extracts has become increasingly important (Baratta *et al.*, 1998).

The *Eucalyptus* genus comprises over 500 species of aromatic trees and shrubs. Previous research suggests that some species counteract influenza viruses; others are antimalarial or highly active against bacteria (Ody, 1994). Of all Australian eucalypts, *Eucalyptus globulus* Labill., Myrtaceae (Tasmanian Blue Gum) is the species most widely introduced overseas and has been established especially throughout the Mediterranean region. It contains great amount of essential oil (*Oleum eucalypti*), up to 3.5% (w/w), which is used in medicine, aromatherapy and perfumes (Bremness, 2000).

In this study, fresh leaves of *E. globulus*, collected in March 2002 from the southern part of Montenegro (Boka Kotorska Bay), were extracted by hydrodistillation in a Clevenger-type apparatus for 2.5h. The obtained oil was dried over anhydrous sodium sulphate and stored at 6°C until analysed.

The yield of obtained oil was evaluated gravimetrically and chemical composition was determined by gas chromatography-mass spectrometry (GC/MS).

Diffusion technique on Mueller-Hinton agar was applied to test sensitivity of certain human pathogenic bacteria and fungi to *E. globulus* essential oil. The essential oil (5, 10, 15, 20 and 30µl) was poured on the Mueller-Hinton base prepared as required.

In this paper the following microbial strains, isolated from patients treated at the hospital or under out-patient regime, were examined: *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Enterococcus* sp., *Streptococcus pyogenes*, *Morganella morganii*, *Providentia stuarti*, *Enterobacter cloacae*, *Acinetobacter baumannii*, *Citrobacter freundii*, *Salmonella infantis* and *Candida albicans*.

In addition, reference antibiotic disks such as ofloxacin, ampicillin, erythromycin, tetracycline, amixacin, ceftriaxon, imipenem etc. were used for comparison (provided by the Institute for Serums, Vaccines and Diagnostic Preparations - Torlak, Yugoslavia).

Eucalyptus essential oil yield was 1.8% (w/w) on a fresh weight basis, whereas GC/MS analysis resulted in the identification of a total of 11 constituents, being eucalyptol (85.8%), α -pinene (7.8%) and β -myrcene (1.9%) the main components. Other identified compounds in gained oil were β -pinene, α -phellandrene, γ -terpinene, linalool, pinocarveol, terpinen-4-ol and α -terpineol.

The results of testing antimicrobial activity are shown in diameters of inhibition zones. The gained results revealed that the essential oil of *E. globulus* has very strong antimicrobial activity, especially against *Streptococcus pyogenes*, *Escherichia coli*, *Candida albicans*, *Staphylococcus aureus*, *Acinetobacter baumannii* and *Klebsiella pneumonas*.

Baratta MT, HJ Dorman, SG Deans, AC Figueiredo, JG Barroso, G Ruperto (1998) *Flavour Fragr. J.*, 13: 235-244.

Bremness L (2000) *Herbs*, Dorling Kindersley, 54.

Ody P (1994) *The complete guide medicinal herbal*, Dorling Kindersley, 61.

P. 32. Comparative headspace-SPME and hydrodistillation of two *Artemisia* sp. from Turkey

B. Demirci, F. Demirci, K. H. C. Başer

bdemirca@anadolu.edu.tr

Medicinal and Aromatic Plant and Drug Research Centre (TBAM), Anadolu University, 26470-Eskişehir, Turkey

Fresh plants of *Artemisia spicigera* C. Koch and *A. scoparia* Waldst. et Kit. (Compositae) were subjected to Headspace-SPME (HS-SPME) and the same plants were air-dried and hydrodistilled to obtain their essential oils. Headspace volatiles were analyzed by GC/MS. The oils were analysed by GC and GC/MS.

Camphor (37.5% and 43.9%) was found as main constituent in both SPME and oil samples of *A. spicigera*. However, in the headspace-SPME samples of *Artemisia scoparia*, capillene (a diacetylene) (53.1%) was found as the main constituent while β -pinene (20.8%), β -caryophyllene (16.4%), (Z)- β -ocimene (16.4%), myrcene (12.8%) and limonene (11.0%) were main components in the oil.

P. 33. The chemical composition and antimicrobial properties of South African *Plectranthus* (Lamiaceae) species

K. Maistry^a, A. M. Viljoen^a, S. F. van Vuuren^a, Betül Demirci^b, K. Hüsni Can Başer^b

viljoenam@therapy.wits.ac.za

^aDepartment of Pharmacy and Pharmacology, Faculty of Health Sciences, University of the Witwatersrand, 7 York road, Parktown 2193, South Africa

^bMedicinal and Aromatic Plant and Drug Research Center (TBAM), Anadolu University. 26470-Eskişehir, Turkey

The genus *Plectranthus*, a member of the mint family, has a cosmopolitan distribution of 300 species worldwide of which 50 are indigenous to South Africa. Many members of this highly aromatic group of plants are used extensively by the Zulu to treat various respiratory disorders. Guided by both traditional use and chemotaxonomic criteria, 12 indigenous species were selected, the leaves hydrodistilled and the essential oils analysed by gas chromatography (GC) and GC coupled to mass spectrometry. Antimicrobial assays incorporating 4 bacteria, 2 yeasts and 2 fungi were performed on the essential oils.

Disc diffusion assays on *Bacillus cereus* showed *P. grandidentatus* being the most active essential oil exhibiting inhibition (at 50 mg/ml) comparable to that of the Neomycin control. Three species showed promising activity against *Bacillus* species; *P. ciliatus* accumulates bicyclogermacrene (17%) and spathulenol (16%), *P. hadiensis* produces T-cadinol (27%), α -fenchone (18%) and cubenol (15%) as major compounds while *P. porphyranthus* produced cubenol (36%) as the major metabolite. A TLC bioautographic assay indicated that several essential oil compounds contribute to the observed antimicrobial activity.

P. 34. Detoxification of terpinolene by *Botrytis cinerea*

Afgan Farooq^{a, c}, Fatih Demirci^b, M. Iqbal Choudhary^c, Atta-ur-Rahman^c, Satoshi Tahara^a, K. Hüsnü Can Başer^b

fdemirci@anadolu.edu.tr

^a Division of Applied Biosciences, Graduate School of Agriculture, Hokkaido University, 060-8589 Sapporo, Japan

^b Medicinal and Aromatic Plant and Drug Research Centre (TBAM), Anadolu University, 26470-Eskişehir, Turkey

^c International Centre for Chemical Sciences, H. E. J. Research Institute of Chemistry, University of Karachi, 75270-Karachi, Pakistan

In the course of our microbial transformation studies of monoterpenes, detoxification of terpinolene by the plant pathogenic fungus *Botrytis cinerea* resulted in the formation of two hydroxylated metabolites: 2,3-dihydro-3 β ,6 β -dihydroxy-terpinolene (39 %) and 2,3-dihydro-1 α ,3 α -dihydroxy-terpinolene (20 %), respectively. Terpinolene showed good levels of antifungal activity while both the metabolites were inactive against the other tested plant pathogenic fungus *Cladosporium herbarum*. The antifungal activity of the substrate and the metabolites were tested at a concentration of 10 microgram per ml using a bioautographic assay.

P. 35.**HS-SPME of *Osyris alba* L. flowers**Fatih Demirci and K. Hüsnu Can Başer

fdemirci@anadolu.edu.tr

Medicinal and Aromatic Plant and Drug Research Centre (TBAM), Anadolu University, 26470-Eskişehir, Turkey

Osyris alba L. (Santalaceae) was subjected to HS-SPME (Head Space- Solid Phase Micro Extraction) and subsequent GC/MS analysis due to the strong fragrance of volatiles emitted by the fresh cut flowering branches. SPME with polydimethylsiloxane/divinylbenzene (PDMS/DVB) fibre resulted in the identification of thirteen hexyl and hexenyl derivatives with a fruity, pear type, leafy green odour.

P. 36. Effect of nitrogen on *Pimpinella anisum* hairy root growth and essential oil production

Carla S. Duarte^a, A. Cristina Figueiredo^b, M. Graça Miguel^a, José G. Barroso^b, Luis G. Pedro^b, J. J. C. Scheffer^c

cduarte@ureach.com / cduarte22@hotmail.com

^aFaculdade de Engenharia de Recursos Naturais, Universidade do Algarve, Campus de Gambelas, 8000-117 Faro, Portugal

^bCentro de Biotecnologia Vegetal, Dep. de Biologia Vegetal, FCL, C2, Campo Grande, 1749-016 Lisbon, Portugal

^cDivision of Pharmacognosy, Leiden University, Gorlaeus Labs, PO Box 9502, 2300 RA Leiden, The Netherlands

Pimpinella anisum hairy root cultures have been maintained, for over six years, in SH liquid medium (Schenk and Hildebrandt, 1972), in darkness, on orbital shakers (80 rpm) at 24°C (Santos *et al.*, 1998), and subcultured every three weeks. Different nitrogen sources, NO₃⁻ / NH₄⁺ ratios and nitrogen amounts were assayed (Table 1) in order to study their effect on the anise hairy root growth and the essential oil production.

Table 1. NO₃⁻ / NH₄⁺ ratio and nitrogen source of the different media assayed.

Media	NO ₃ ⁻ /NH ₄ ⁺	Nitrogen source
SH	25/3	KNO ₃ / (NH ₄) ₂ HPO ₄
SHa	25/1	KNO ₃ / (NH ₄) ₂ SO ₄
SHb	39/21	KNO ₃ / NH ₄ NO ₃
SH ⁺	25/3	Ca(NO ₃) ₂ ·4H ₂ O / (NH ₄) ₂ HPO ₄
SH ⁺⁺	0/3	(NH ₄) ₂ HPO ₄

The hairy root cultures growth was determined by the dissimilation method (Schripsema *et al.*, 1990) and by dry-weight and fresh-weight determination. Morphological characterisation was performed by scanning electron microscopy. The essential oils from each of the culture types were isolated by distillation-extraction and analysed by GC and GC-MS. Both for the growth and essential oil composition studies, hairy root samples were collected periodically along the growth cycle.

In comparison to SH-grown hairy root cultures, all other culture conditions assayed led to a slower growth, a higher tendency to browning and to development of *callus* tissue.

The dominant components of the constitutive essential oil from the SH anise hairy root cultures were *trans*-epoxypseudoisoeugenyl 2-methylbutyrate (>69%) and β-bisabolene (>9%). The former component was found to be dominant also in the SHa, SHb and SH⁺ grown hairy root cultures (>27%). The dominant component of the oil from the hairy root cultures maintained in SH⁺⁺ was octyl isobutyrate, in a percentage higher than 12%. This compound occurred in a similar percentage in the essential oils isolated from the cultures maintained in SHa (13%), SHb (13%) and SH⁺ (12%), but in a much lower relative amount (1%) in the oil from the SH anise hairy root cultures.

Santos PM, AC Figueiredo, MM Oliveira, JG Barroso, LG Pedro, SG Deans, AKM Younus, JJC Scheffer (1998) *Phytochemistry*, **48**: 455-460.

Schenk UR, AC Hildebrandt (1972) *Can. J. Bot.*, **50**: 199-204.

Schripsema J, A H Meijer, F Van Iren, H J G Ten Hoopen, R Verpoorte (1990) *Plant Cell Tiss. Org. Cult.*, **22**: 55-64.

P. 37. Composition of the volatiles from *Gennaria diphylla* (Link) Parl. (Orchidaceae) grown on the Madeira Island

Francisco M. Fernandes^{a,d}, A. Cristina Figueiredo^b, José G. Barroso^b, Luis G. Pedro^b, Christopher C. Wilcock^c, M. A. Pinheiro Carvalho^d

^a goodyera@dragoeiro.uma.pt

^a Jardim Botânico da Madeira, Caminho do Meio, Bom Sucesso, 9050 Funchal, Madeira, Portugal

^b Centro de Biotecnologia Vegetal, Dep. de Biologia Vegetal, FCL, C2, Campo Grande, 1749-016 Lisbon, Portugal

^c University Aberdeen, Dpt. of Plant and Soil Science, Aberdeen AB9 2UD, UK

^d BIOTecMOL – CCBG – UMA, Edifício Penteada, 9000-300 Funchal, Portugal

Gennaria diphylla (Link) Parl. [= *Peristylus cordatus* (Willd.) Lindl.] is the only species of the genus. It is a very unusual terrestrial orchid, which characteristically possesses only two cordate leaves at the base. The plant is 8 to 45cm height with a glabrous inflorescence bearing 8 to 46 flowers. In the Madeiran archipelago, this orchid can be found both on Madeira, in the higher parts of Porto Santo and also on Deserta Grande. On Madeira, *G. diphylla* grows in the Laurisilva, but can be seen also at lower altitudes. This species can also be found in Southern Spain, Sardinia, Corsica and Tunisia, Canary Islands and in the Portuguese mainland where it grows in small populations sheltered by *Corema album* along the coastal woods (Pena and Cabral 1997, Neiland 2000).

The aerial parts of *G. diphylla* were collected during the flowering phase at Montado do Sabugal, Madeira. The essential oils were isolated from the inflorescences, by distillation-extraction for 3h as well as by headspace sorption on charcoal filters for 48h, and analysed by GC and GC-MS.

The essential oils isolated by distillation-extraction possessed very different qualitative and quantitative compositions compared to those isolated by headspace sorption. Of the thirty-two components identified in the essential oil isolated by distillation-extraction, only eleven were detected in the volatiles isolated by headspace sorption. Conversely, of the eighteen components identified in the volatiles isolated by headspace sorption, seven were not detected in the former oil.

The non-terpenoid compounds were the main components (79%) of the oils isolated by distillation-extraction, whereas the monoterpene fraction amounted only to 7%. *n*-Octacosane (19%) and *n*-heptacosane (13%) were the dominant components of this oil.

Oxygen-containing monoterpenes were the major compounds of the volatiles isolated by headspace sorption, amounting to 67% of the total oil. Among those 5-methyl-5-vinyl-tetrahydrofuran derivatives were the dominant components (61%), *cis*-arbusculone (28%) and lilac alcohol (26%) being the main volatile components of this fraction.

The considerable differences between volatile compounds isolated by distillation-extraction and headspace sorption are related to the isolation procedure used. On the one hand the large amount of hydrocarbons detected in the distillation-extraction oil is probably due to the decomposition of the leaf cuticle waxes. On the other hand, charcoal traps are known to be ineffective in trapping the full range of floral fragrance compounds, moreover, the trapping time influences the relative amount of the fragrance components. In addition, some compounds, namely lilac aldehydes, decompose on charcoal during collection and storage, as it has been found for other orchid species (Patt *et al.* 1988).

Neiland R, (2000) *Genera Orchidacearum, Vol. 2, Orchidoideae*, Ed. Pridgeon, A., Royal Botanic Gardens, Kew.

Patt JM, DF Rhoades, JA Corkill (1988) *Phytochemistry* 27: 91-95.

Pena A, Cabral J (1997) *Roteiros da natureza – Algarve*, Temas e Debates, Lisboa, p.85.

P. 38. Essential oils from cultivated plants of *Hypericum androsaemum* L.Ana P. Guedes ^a, Lúcia R. Amorim ^a, Ana Vicente ^b, M. Fernandes-Ferreira ^a

mfferreira@bio.uminho

^a Department of Biology, University of Minho, Campus Gualtar, 4710-057 Braga, Portugal^b ERCA/DRAEDM, S. Pedro de Merelim, Braga, Portugal

Plants from *Hypericum androsaemum* L. were cultivated by March 1998 in an experimental farm from Direcção Regional de Entre Douro e Minho (DRAEDM), at the northern region of Portugal, by planting slips of about 15 cm long obtained from wild plants growing near Ponte de Lima (Facha).

The cultivated plants were pruned by July 1999 (S1), November 1999 (S2) and June 2000 (S3) and the respective aerial parts were subjected to hydrodistillation in a Clevenger-type apparatus over one hour. The essential oils were analysed by GC and GC-MS. The identification of the compounds was performed with the help of mass spectra libraries and confirmed, whenever possible, by comparison with the relative retention times and mass spectra of authentic standards.

Identification was achieved for 90, 98, and 97% of the essential oils from the samples S1, S2, and S3 respectively, in a basis of the total area of the peaks plotted in the respective GC chromatograms and reports. More than seventy compounds were identified, majority distributed by five groups: monoterpene hydrocarbons (MH), oxygen-containing monoterpenes (MO), sesquiterpene hydrocarbons (SH), oxygen-containing sesquiterpenes (SO) and other hydrocarbons (H). The percentages of each one of these compound groups varied from one sample to another. In any case however, SH was the main compound group accounting for 58, 77, and 43% of the essential oils from the samples S1, S2, and S3 respectively. Following this sequence, the relative amounts of each one of the other four main compound groups were: MH (17, 7, and 29%); SO (6, 3, and 4%); H (4, 4, and 13%); MO (0.5, 0.3, and 0.5%). The ten main compounds and respective percentages (%) in each one of the three samples were the following ones: S1 - β -caryophyllene (12.5), γ -elemene (8.5), β -gurjunene (6.1), limonene (5.9), β -pinene (4.2), terpinolene (3.9), γ -muurolene (3.8), *n*-nonane (2.9), 2-hexenal (2.5), *trans*- β -ocimene (1.4); S2 - γ -elemene (17.9), β -gurjunene (15.5), β -caryophyllene (15.1), 2-hexenal (5.3), γ -muurolene (4.4), β -elemene (2.7), β -pinene (2.2), limonene (2.1), 1-octene (1.6), terpinolene (1.4); S3 - limonene (15.4), β -caryophyllene (9.4), γ -elemene (8.0), 2-hexenal (7.9), β -gurjunene (7.6), 1-octene (7.3), β -pinene (6.4), *n*-nonane (3.6) α -pinene (3.2), γ -muurolene (2.1).

Variations in essential oil composition could be induced by different physiological or environmental factors whose variation over the vegetative cycle may influence the compounds turnover. The attack of the cultivated plants by some kind of organisms constitute another type of factors that contingently could influence the essential oil composition. On the *H. androsaemum* cultivated plants, here reported, we identified two contaminant organisms; a rust fungus - *Uromyces* sp. - and an aphid - *Aphis gossypii* Glover. The populations of these contaminants appeared by April of 1998 and 1999, rising over the following four months. It remains speculative however, any interpretation inherent to the respective effects on the essential oil composition, because no type of control was made.

Sponsored by Programme AGRO / 8 / key action 8.1 / Project 338.

P. 39. The effect of phytohormones on the essential oils produced by in vitro shoots of sage (*Salvia officinalis* L.)

Paula C. Santos-Gomes, M. Fernandes-Ferreira

mfferreira@bio.uminho

Department of Biology, University of Minho, Campus Gualtar, 4710-057 Braga, Portugal

In vitro shoot cultures of sage (*Salvia officinalis* L.) were established as previously described (Santos-Gomes and Fernandes-Ferreira, 2002). 2,4-Dichlorophenoxyacetic acid (2,4-D) at 0.1 or 0.05 mg/l, combined with one cytokinin: benzyladenine (BA) or zeatine (ZEA) or kinetin (KIN), at 1.5 mg/l, and 2.0 or 4.0 mg/l, in the case of KIN, were tested in the evaluation of the effect of the phytohormones in the production of essential oils by these type of cultures. From the four types of primary explants tested (nodal shoot segments, inter-nodal shoot segments, leaves, and root segments, all of them excised from aseptic seedlings), only the nodal shoot segments were efficient in promoting direct shoot regeneration through organogenesis. Concentration of 0.05 mg/l 2,4-D has promoted higher number of shoots *per* explant and higher linear growth than 0.1 mg/l 2,4-D concentration, except when combined with KIN at 1.5 mg/l. In this case, differences were not weighty. However the percentage of explants that originated shoots did not change by the variation of 2,4-D concentration from 0.1 to 0.05 mg/l but was influenced by the variation of the KIN concentration. The increase in KIN concentration from 1.5 to 2.0 and 4.0 promoted multiple shoot formation.

After the fifth subculture to the same medium and same supplementation conditions, the shoots grown over 33-35 days were subjected to hydrodistillation in a Clevenger type apparatus over one hour. The essential oils were analysed by GC and GC-MS. The identification of the compounds was performed with the help of mass spectra libraries and confirmed, whenever possible, by comparison with the relative retention times and mass spectra of authentic standards. The quantification was performed according the methodology previously described (Santos-Gomes and Fernandes-Ferreira, 2001).

The highest content of essential oil (~24 mg/g of biomass dry weight) was obtained from shoots grown on Murashige and Skoog (1962) medium supplemented with 0.05mg/l of 2,4-D and 2.0mg/l of KIN. From the three cytokinin tested, KIN was the most effective in the essential oil production, specially when combined with 0.05 mg/l 2,4-D. The phytohormone supplementation influenced the relative amounts of the main group of compounds. However, at the individual level, the percentages of the major compounds were not greatly influenced. Camphor was the major compound (20-24%) followed by α -thujone (13-16%), β -pinene (9-15%), β -thujone (~7%), camphene (6-7%), manool (4-8%), α -humulene (5-7%), 1,8-cineole (4-6%), α -pinene (4-5%), and limonene (~2%). These composition parameters are majority within the ranges of the profile defined by standard ISO 9909 for essential oils of sage (Bruneton, 1999) with exception for the levels of α -thujone (somewhat low) and perhaps, β -pinene and manool, which are not referred by this author as taking part of the same standard profile.

Sponsored by FCT through the Project POCTI / AGR / 43482 / 2001.

Bruneton J (Ed.) (1999). *Pharmacognosy, Phytochemistry, Medicinal Plants*, Intercept Ltd, London
Murashige T, Skoog F (1962) *Physiol. Plantarum* 15: 473-497.
Santos-Gomes PC, M Fernandes-Ferreira (2001) *J. Agric. Food. Chem.* 49: 2908-2916.
Santos-Gomes PC, M Fernandes-Ferreira (2002) *Plant Science* (in press).

P. 40. Glandular trichomes and volatile oils from two *Plectranthus* species

Lia Ascensão ^a, Marília de M. Castro ^b, Cláudia Ferreira ^a, Mafalda Selas ^a, Luis G. Pedro ^a, A. Cristina Figueiredo ^a, José G. Barroso^a

c.ferreira@iol.pt

^a Centro de Biotecnologia Vegetal, Dept. Biologia Vegetal, FCL, C2, Campo Grande, 1749-016 Lisbon, Portugal

^b Departamento de Botânica, Instituto de Biologia, Universidade Estadual de Campinas, Brazil

The genus *Plectranthus* L'Herit belongs to the Lamiaceae, known for their aromatic and medicinal properties. It is a large genus of about 350 species distributed through Africa, Asia and Australia, with over 40 species of indigenous to Southern Africa (Codd, 1985). *P. ecklonii* and *P. verticilatus*, ornamental plants in Europe, occur in South Africa along forest margins of woodlands in semi-shady places where frost is not too severe. This study compares the morphology and distribution of the trichomes of these two species, as well as the histochemistry of the secretory products and the composition of the essential oils produced.

Glandular and non-glandular trichomes are spread over the vegetative and reproductive organs. The glandular trichomes observed are of the same types as described previously for the Lamiaceae – peltate and capitate. However, on *P. ecklonii* and *P. verticilatus*, they have *in vivo* an uncommon and characteristic orange-to-reddish colour and, unlike the other peltate trichomes present in the Lamiaceae, never develop a large subcuticular space. They exhibit a lobated shape, somewhat flattened-to-depressed in the centre, even during the secretory phase. Capitate trichomes have an ovoid unicellular head and a short cell or an elongated two-to-three-celled stalk slightly enlarged at the base. In the short-stalked trichomes the secretion, mainly polysaccharidic, is probably exuded by micropores, whereas in the long-stalked a heterogeneous secretion stores temporarily in a subcuticular space, being released by cuticular rupture.

Glandular trichomes occur on both leaf sides. Peltate trichomes are more numerous on the abaxial surface, where they are particularly concentrated at basal intervein areas. Short capitate trichomes predominate along the veins of the adaxial surface. The reproductive organs, namely the calyx and corolla, exhibit scarce long-stalked capitate trichomes in addition to those already reported. Peltate trichomes are also abundant on the stamens, between the two anther lobes and on the basal portion of the style, between the four lobes of the ovary.

The aerial parts of *P. ecklonii* and *P. verticilatus* were collected, during the flowering and the vegetative phases at the Botanical Garden of Lisbon. The essential oils were isolated separately from flowers and from leaves by hydrodistillation, for yield determination, and by distillation-extraction for analysis by GC and GC/MS. All the oils were obtained in a yield <0.05% (v/w). In *P. ecklonii* monoterpenes represented the main fraction of the oil from the flowers, amounting to 74%, α -pinene (35%), sabinene (16%) and β -pinene (10%) being the main compounds. The oils from the leaves of this species, both from the flowering and the vegetative phases, were dominated by octen-3-ol, representing 49% and 62%, respectively. Monoterpenes represented only 7% and 2% of the total oils, respectively. In *P. verticilatus* all the oils were dominated by octen-3-ol that attained 41%, 84% and 85% in the oil from the flowers and in those from the leaves collected during the flowering and the vegetative phases, respectively. The monoterpene fraction amounted to 35% in the flower oil, limonene (15%) and sabinene (10%) being the most representative compounds, but accounted only 10% and 3% of the oils from the leaves collected during the flowering and the vegetative phases, respectively.

Codd LE (1985) *Lamiaceae*. In: Flora of Southern Africa, vol. 28, part 4, pp: 137-172, Leistner OA (ed.). Botanical Res. Inst. Dept. of Agriculture and Water Supply, Pretoria.

P. 41. *Plectranthus laxiflorus*: morphology and histochemistry of glandular trichomes and essential oil composition

Cláudia Ferreira, Lia Ascensão, Luis G. Pedro, A. Cristina Figueiredo, José G. Barroso
c.ferreira@iol.pt

Centro de Biotecnologia Vegetal, Dept. Biologia Vegetal, FCL, C2, Campo Grande, 1749-016 Lisbon, Portugal

Plectranthus laxiflorus, an aromatic perennial herb that emanates a characteristic citronella-like scent, occurs in Southern Africa along forest margins and on shady stream banks (Codd, 1985). In local medicine it is reputed to be antiseptic and analgesic. Infusions from leaves are commonly used for coughs and colds. Powdered leaves are also employed by Zulus as an enema for feverishness, abdominal upsets and influenza (Watt and Breyer-Brandwigk, 1962). Pursuing our studies on the genus *Plectranthus*, this work reports on the morphology, distribution and histochemistry of the glandular trichomes of *P. laxiflorus* and the composition of its essential oil.

Four distinct types of glandular trichomes were identified on the vegetative and reproductive organs. Unlike the majority of the peltate trichomes reported for *Plectranthus* species, which *in vivo* have a characteristic orange-to-reddish colour and never develop a large subcuticular space, the peltate trichomes of *P. laxiflorus* are colourless and become nearly spherical during the secretory phase. Capitate trichomes are divided in three subtypes according to the dimension of the stalk and the secretion process. In addition to the short and long-stalked trichomes, similar to the others previously described in the Lamiaceae, an unusual type of capitate trichomes (ellipsoidal trichomes) are reported for the first time in this family. They consist of an elongated four-to-twelve-celled stalk, which bears an ellipsoidal head cell that stores the secretion in the vacuoles.

Peltate trichomes are abundant at the intervein areas of the abaxial leaf surface, whereas ellipsoidal ones predominate on the leaf base and along the veins and margins. Short capitate trichomes are numerous on the adaxial surface above the veins. The flower shows all the trichome types here reported. Ellipsoidal trichomes are densely arranged on the abaxial surface of the upper corolla lip and on the calyx veins and teeth. However, glandular trichomes are absent on the stamens and carpels. The histochemical characterization of the secretion reveals the presence of terpenoids, flavonoids and polysaccharides as the main constituents.

The aerial parts of *P. laxiflorus* were collected, during the flowering and the vegetative phases at the Botanical Garden of Lisbon. The essential oils were isolated separately from flowers and from leaves by hydrodistillation and by distillation-extraction, and analysed by GC and GC/MS. All the oils, isolated from the flowers (F), leaves from the flowering phase (LF) and leaves from the vegetative phase (LV), were obtained in a yield <0.05% (v/w). Monoterpenes represented the main fraction, amounting to 55% (F), 70% (LF) and 51% (LV). The identified sesquiterpenes attained 27% (F), 26% (LF) and 39% (LV). The main components were α -phellandrene (10%), *trans*- β -ocimene (7%), citronellal (7%) and citronellol (6%) in the oil from the flowers, α -phellandrene (24%), *trans*- β -ocimene (15%), germacrene-D (10%) and β -phellandrene (10%) in that from the leaves of the flowering phase, and α -phellandrene (21%), germacrene-D (19%), *trans*- β -ocimene (9%) and β -phellandrene (9%) in that from the leaves of the vegetative phase. These results are in close agreement with those previously reported for the oils isolated from flowers and leaves collected during the flowering phase of the plant (Figueiredo *et al.*, 2000).

Codd LE (1985) *Lamiaceae*. In: Flora of Southern Africa, vol. 28, part 4, pp: 137-172, Leistner OA (ed.). Botanical Res. Inst. Dept. of Agriculture and Water Supply, Pretoria.

Figueiredo AC, FMS Silva, L Ascensão, JG Barroso, LG Pedro, JJC Scheffer (2000) 31st Intern. Symp. on Essential Oils, Hamburg (Abstract A-30).

Watt JM, MG Breyer-Brandwigk (1962) The Medical and Poisonous Plants of Southern Africa. 2nd edn., E & S Livingstone Ltd, London.

P. 42. *Pimpinella anisum* hairy root cultures: growth and essential oil production in a two-phase system

Francisco F. Silva^a, A. Cristina Figueiredo^b, M. Graça Miguel^a, José G. Barroso^b, Luis G. Pedro^b, Johannes J. C. Scheffer^c

^b acsf@fc.ul.pt

^a Faculdade de Engenharia de Recursos Naturais, Universidade do Algarve, Campus de Gambelas, 8000-117 Faro, Portugal

^b Centro de Biotecnologia Vegetal, Dep. de Biologia Vegetal, FCL, C2, Campo Grande, 1749-016 Lisbon, Portugal

^c Division of Pharmacognosy, Leiden University, Gorlaeus Labs, PO Box 9502, 2300 RA Leiden, The Netherlands

In an attempt to avoid the toxicity of some essential oil components, and of making it easier to collect the volatiles secreted by *in vitro* cultures, the growth and essential oil composition of *Pimpinella anisum* hairy root cultures has been determined in a two-phase system. This system is based on the addition to the culture medium of an adsorbent that temporarily accumulates the products secreted by the cells. In this experiment Miglyol 812N (a non-toxic triglyceride immiscible with water) was added to the culture medium (5% v/v) before autoclaving.

The hairy root cultures growth in bubble-column bioreactors containing 1 litre of SH (Schenk and Hildebrandt 1972) medium, both with and without Miglyol (control cultures), was determined by dry-weight and fresh-weight determination, over a period of five weeks. The essential oils from each of the culture types were isolated by distillation-extraction and analysed by GC and GC-MS. Both for the growth and essential oil composition studies, the hairy roots were weekly collected along the growth cycle, and the essential oils isolated separately from the hairy roots and culture medium.

The control cultures and those grown with Miglyol showed comparable growth patterns, attaining a specific growth rate of 0.24/day and 0.22/day, respectively.

The dominant component of the oils isolated from the control cultures was the phenylpropanoid *trans*-epoxypseudoisoeugenyl 2-methylbutyrate (*trans*-EPIE2MB), attaining a maximum of 1.12 µl/g d.w. β-Bisabolene (0.47 µl/g d.w.), zingiberene (0.23 µl/g d.w.), dodecanal (0.13 µl/g d.w.) and *trans*-pseudoisoeugenyl 2-methylbutyrate (*trans*-PIE2MB; 0.11 µl/g d.w.), were the other four main components of the oils. The same components were found to be the dominant ones in the Miglyol-grown cultures, but sometimes in lower concentrations; *trans*-EPIE2MB attained a maximum of 1.07 µl/g d.w., β-bisabolene 0.19 µl/g d.w., zingiberene 0.10 µl/g d.w., dodecanal 0.05 µl/g d.w. and *trans*-PIE2MB 0.09 µl/g d.w.

Some of the volatiles isolated from the Miglyol culture medium such as dodecanal (0.019 µl/L), zingiberene (0.032 µl/L) and β-bisabolene (0.49 µl/L) attained, particularly at the end of the growth cycle, higher concentrations than those obtained with the control culture medium (0.005 µl/L, 0.012 µl/L and 0.022 µl/L, respectively). Interestingly, while compounds such as *trans*-anethole, that is usually found in trace amounts in the oils from the control hairy roots and not detected in the control medium, attained a maximum of 0.031 µl/L in Miglyol culture medium, *trans*-EPIE2MB was only detected in amounts below 0.008 µl/L throughout the culture period.

In conclusion, although *P. anisum* Miglyol hairy root cultures revealed comparable growth to control cultures and were even less prone to oxidation on harvest, there was no *de novo* synthesis and a smaller amount of volatiles was recovered from these cultures. The larger amounts of some volatiles recovered from medium of the Miglyol hairy root cultures was probably due to their higher adsorption to Miglyol than to the hairy roots, in this way not reaching toxic levels to the cells.

Acknowledgements: The authors are grateful to CONDEA Chemie GmbH for a generous gift of Miglyol.

Schenk UR, AC Hildebrandt (1972) *Can. J. Bot.* **50**: 199-204.

P. 43. Study and comparison of two mandarin essential oils from southern Brazil

Caren D. Frizzo ^a, Eduardo Dellacassa ^b

aripe@terra.com.br

^a Aripê Citrus Ltda., RS 124 Km 1.2, 95780-000, Montenegro -RS, Brazil

^b Cátedra de Farmacognosia y Productos Naturales, Facultad de Química, General Flores 2124. 11800 Montevideo, Uruguay

Only a detailed knowledge on the composition of the oils obtained by different cultivars of mandarin can provide the choice of the cultivars which will produce the best oils (Calvarano *et al.*, 1989; Dellacassa *et al.*, 1992; Cotroneo *et al.*, 1994; Dugo *et al.*, 1999; Verzera *et al.*, 2000). To exemplify this topic, this study reports the composition of the essential oils of two varieties of mandarins (*Citrus reticulata*, Blanco), commonly called "Caí" and "Montenegrina," growing in Southern Brazil. Mandarin fruits were collected in a small scale experimental area located at Aripê Citrus, Montenegro, from 4-5 years old trees and processed during the 2002 Season (February-May). The volatile fractions were obtained by hydrodistillation of fresh mandarin peels using a Clevenger-type apparatus. The oil composition was analyzed by HRGC and HRGC-MS. Repeatability of the measuring system showed variation coefficients under 5% for all the components reported. The essential oil components were identified by comparison of their linear retention indices on methyl silicone. Comparison of fragmentation patterns in the mass spectra with those stored on databases (Jennings and Shibamoto, 1980; Adams, 1995) was also performed.

Sixty-seven components were identified in the oils from both mandarin varieties, which were similar in composition. Nevertheless, some characteristics such as the content of the main components with commercial interest, particularly dimethyl anthranilate and α -sinensal, make the oils from both varieties different from each other.

Adams RP (1995) Identification of essential oil components by gas chromatography/mass spectroscopy. Allured Publ. Corp., Illinois, USA.

Calvarano M, I Calvarano, E Dellacassa, P. Menendez, A Di Giacomo (1989) *Essenze Deriv. Agrumari* 59: 397-406.

Cotroneo A, L Mondello, I Stagno D'Alcontres (1994) *Essenze Deriv. Agrumari* 44: 275-285

Dellacassa E, C Rossini, P Menéndez, P Moyna, A Verzera, A Trozzi, G Dugo (1992) *J. Essent. Oil Res.* 3: 265-272.

Dugo G, KD Bartle, I Bonaccorsi, M Catalfamo, A Cotroneo, P Dugo, G Lamonica, H McNair, L Mondello, P Previtte, I Stagno D'Alcontres I, A Trozzi, A Verzera (1999) *Essenze Deriv. Agrumari* 69: 79-111.

Jennings W, T Shibamoto (1980) Qualitative analysis of flavor and fragrance volatiles by glass capillary gas chromatography. Acad. Press, New York.

Verzera A, A Trozzi, A Cotroneo, D Lorenzo, E Dellacassa (2000) *J. Agr. Food Chem.* 48: 2903-2909.

P. 44. Characterization of fragrance notes used in perfumery by GC-MS

Vera Mata, Paula B. Gomes, A. E. Rodrigues
pgomes@fe.up.pt

LSRE, Faculty of Engineering of University of Porto, Portugal

Gas chromatography (GC) associated with mass spectrometry (MS) is currently the best analysis method of complex mixtures, namely essential oils and perfumes.

A perfume has a well-defined structure that is usually designed according to the fragrance pyramid (Figure 1). The perfume composition is therefore based on three kinds of odour notes:

- ❖ **Top notes**, which are responsible for the first impression of a perfume when it is smelled for the first time, such as citrus notes;
- ❖ **Middle notes** that become evident some time after the application, mostly floral and spicy notes, constituting the heart of the perfume and defining its unique character;
- ❖ **Base notes** make up the long-lasting scent that in the end smoothly fade away, working as well as fixative materials, which include vanilla, musk, amber and sandalwood




Figure 1 – The Fragrance Pyramid




Figure 2 – GC-MS chromatograms of lemon (a), geranium (b) and vanilla (c).




Figure 3 – GC-MS chromatogram of a perfume with triangular structure.

The retention time of the fragrance chemicals using a GC capillary polar column is related to the evaporating profile of a perfume when applied on the skin or on the smelling strips that are used for olfactory tests.

In this work, the fragrance notes of lemon, rose geranium and vanilla are characterized by GC-MS using a polar column (CP-Wax 52 CB), representing a top, middle and a base note, respectively (Figure 2). In general, it can be observed that top notes are the first to evaporate and therefore present the lowest retention times. On the other hand, the base notes have the highest retention times since they are the ones that last longer until they evaporate completely. When considering a perfume composition, it can be also identified the three areas, corresponding to the three kinds of odour notes, as can be seen in Figure 3 : I – top; II – middle and III- base note.

The characterization by GC-MS analysis of fragrance notes can provide a significant insight into the perfume composition, enabling to break down its structure into top, middle and base notes, as well as to evaluate the balance of the fragrant materials in the final composition. This way it is possible to establish patterns in perfume composing and GC-MS is a fundamental tool to help the perfumer in this work.

Tudor, E., (1997) *J. Chromatography A*, 779: 287-297

Mariotti, P. et al. (2001) *J. Chromatography A*, 936: 1-22

P. 45. Composition and antibacterial activity of the essential oils from *Zanthoxylum rhoifolium*

Wellington de A. Gonzaga, Ademir F. Morel, Andréia D. Weber, Sandro R. Giacomelli, Euclésio Simionatto, Ionara I. Dalcol, Emilia C. Dessoy

afmorel@baseufsm.br

Departamento de Química, Universidade Federal de Santa Maria, 97105-900 Santa Maria RS, Brazil

Zanthoxylum rhoifolium (Rutaceae), locally called "mamica-de-porca", "mamica-de-cadela", "juvê" is a plant that grows in South America (Brazil, Uruguay, Paraguay and Argentina). It has been used in Brazilian folk medicine against a variety of diseases. As a continuation of our chemical studies on Rutaceae plants (Morel *et al.*, 1997), we now report on the volatile constituents of leaves, fruits and flowers of *Z. rhoifolium*. The oils were analysed by GC and GC/MS. The identification of the chemical constituents was based on comparison of their relative retention times and mass spectra with those obtained from authentic samples and/or the Wiley and NBS/NIST libraries and those published by Adams (1995).

Fourteen constituents were identified representing *ca* of 96% of the total oils. β -Myrcene (32%) and limonene (38%) were the most abundant components in the oil of the fruits, while β -myrcene (65%) was the main component in that from the flowers. Germacrene-D (34%) and bicyclogermacrene (23%) were the most abundant components in the oil from the leaves. As revealed by Bioautography (Homans 1970), the antibacterial studies showed that the oils of leaves and fruits were active against *S. aureus* (Gram⁺) and *K. pneumoniae* and *S. setubal* (Gram⁻) bacteria, while the oil of the flowers was inactive (see Table 1). The microorganisms used in the antibacterial assays have been maintained at the Departamento of Chemistry of the Universidade Federal of Santa Maria, RS, Brasil. For the antimicrobial assays, 50.0, 25.0, 12.5, 6.25, 3.12, 1.06 μ g of the oils were applied to pre-coated TLC plates. The inoculum was prepared by culturing each organism in tryptone soya agar (TSA, Oxoid) at 37°C to a turbidity equivalent to McFarland 0.5 standard (1.5×10^8 CFU/mL). One microliter of each diluted inoculum (10^4 - 10^6 CFU) was applied onto Mueller Agar (MHA-DIFCO). Amoxicillin was used in order to control the sensitivity of the test organisms.

Table 1 Antibacterial activity: minimum amount required for inhibition on bacteria growth on TLC plates (μ g)

Material	<i>S. aureus</i>	<i>S. efidermidis</i>	<i>K. pneumoniae</i>	<i>S. setubal</i>	<i>E. coli</i>	<i>M. luteus</i>
Leaves	3.12	NA	1.06	12.50	NA	NA
Fruits	12.50	NA	50.00	50.00	NA	NA
Flowers	NA	NA	NA	NA	NA	NA

NA: No active.

Acknowledgements: FAPERGS, CNPq

Adams RP (1995) Identification of Essential Oil Components by Gas Chromatography Mass Spectroscopy, Allured Publishing Co

Homans AL and A Fuchs, (1970) Direct bioautography on thin-layer chromatograms as a method for detecting fungitoxic substances. *Journal of Chromatography* 51: 327.

Morel AF, NF Moura, HB Ribeiro, ECSM Machado, EM Ethur, N Zanatta, (1997) Benzophenanthridine Alkaloids from *Zanthoxylum rhoifolium*. *Phytochemistry* 46: 1443-6.

P. 46. Changing of terpene composition in single peltate trichomes of *Salvia officinalis* leaves by direct liquid, non-equilibrium SPME application

Paolo Grassi, Johannes Novak, Heidi Steinlesberger, Chlodwig Franz

Paolo.Grassi@vu-wien.ac.at

Institute for Applied Botany, University of Veterinary Medicine, Veterinarplatz 1, 1210 Vienna, Austria

A new non-equilibrium SPME method was developed for rapid extraction of single oil glands from sage and compared with solvent extracts and the essential oil. Differences between ageing of single peltate trichomes on three insertions of *Salvia officinalis*, an immature apical young leaf, an expanding, and a fully developed leaf were studied. Chemical variation between the leaves showed high sesquiterpene and bornyl acetate contents in the young leaf and high camphene and camphor contents in the old leaf, according to Croteau *et al.* (1981). The intermediary, still expanding leaf was used to carry out an age assignment of single oil glands. The basal region of the intermediary leaf contains oil gland compounds in high conformity with the young leaf trichomes, the remaining oil glands showed inconsistent accumulation patterns.

The oil glands of sage appear as opaque or as transparent heads. Chemical differences in their essential oil composition were found in the content of sesquiterpenes only, whereas Tirillini (1999) showed strong qualitative differences between them. Opaque glands showed a significant higher sesquiterpene rate than the transparent glands.

Croteau R, Felton M, Karp F, Kjonaas R (1981) *Plant Physiology* 67: 820-824.

Tirillini B., Ricci A (1999) *J Essent. Oil Res.* 11: 565-569.

P. 47. Essential oil of *Nepeta haussknechtii* Bornm. from Iran

Zohreh Habibi, Sajjad Sedaghat

zohre1340@hotmail.com

Dept. of Chemistry, Shahid Beheshti University, Tehran, Iran

The genus *Nepeta* comprises about 250 species distributed in the world, of which 67 occur in Iran.

N. haussknechtii plants were collected, during the flowering period, at Chalous, North of Iran, in June 2001. Air dried aerial parts of the plants were grounded and submitted to hydrodistillation, for 4 hours, using a Clevenger-type apparatus. The oil was pale greenish and had a pleasant odour, with a yield of 0.15% (w/w). The oil was analysed by GC/MS.

Twenty-two components, representing 90% of the total oil, were identified of which 4a β , 7 α , 7a β -nepetalactone (35.3%), diisooctyl 1,2-benzene dicarboxylate (16.3%) and 1,8-cineol (15.0%) were the main constituents.

P. 48. Composition of the essential oil of *Satureja isophylla* Rech. f. from Iran

Zohreh Habibi, Sajjad Sedaghat

zohre1340@hotmail.com

Dept. of Chemistry, Shahid Beheshti University, Tehran, Iran

The genus *Satureja* comprises 12 species occurring in Iran. The aerial parts of *S. isophylla* were collected, during the flowering period, at Niknam village, Chalous, North of Iran, in June 2001.

The dried aerial parts of the plants were hydrodistilled, for 4 hours, using a Clevenger-type apparatus to yield 0.1% (w/w) of oil. The oil was yellowish and had a pleasant odour. The chemical composition of the essential oil of *S. isophylla* was analysed by GC/MS. Twenty-six components, representing 95% of the total oil, were identified, with α -eudesmol (24.2%), β -caryophyllene (12.1%) and camphor (9.5%) as main constituents.

P. 49. Utilisation of essential oils and fatty acid esters with the formulation of the “solvent/active” for application in phytosanitary industry

Leon Hernandez Ochoa, G. Vilarem, Z. Mouloungui

lhernandez@ensct.fr, gvilarem@ensct.fr

Laboratoire de Chimie Agro-Industrielle, UMR 1010 INRA/INP-ENSIACET, Ecole Nationale Supérieure des Ingénieurs en Arts Chimiques et Technologiques, 118 route de Narbonne, F-31077 Toulouse Cedex, France

Solvent extraction, co-hydrodistillation and hydrodistillation were compared as methods to extract essential oils from *Origanum dictamnus*, *Origanum vulgare*, *Carum carvi*, *Acore calamus*.

Associating “hydrodistillation and solvent extraction” as traditional techniques of essential oils extraction, co-hydrodistillation process was obtained. Traditionally water is used as solvent in hydrodistillation process, nevertheless in the latter ethyl ester of fatty acid was found to be a good extraction solvent.

Gas chromatography-mass spectrometry analysis (GC-MS) showed that the composition of the isolated essential oils varies depending on the period of harvest the plants and conditions of co-hydrodistillation (solvent/raw matter report).

The products obtained from solvent extraction and co-hydrodistillation with solvents of vegetal origin named “solvent/active “ will be tested for application on phytosanitary industry.

P. 50. GC/MS analysis of the oil from the berries of *Juniperus oxycedrus* L. subsp. *oxycedrus* grown in Spain

Ana Iñigo ^a, Arturo Velasco-Negueruela ^a, M^a José Pérez Alonso ^a, Ginés López ^b

ainigo@terra.es

^a Departamento de Biología Vegetal I (Botánica), Facultad de Biología Universidad Complutense, 28040-Madrid, Spain

^b Real Jardín Botánico de Madrid CSIC Plaza de Murillo 2, 28014-Madrid, Spain

The composition of the essential oil obtained from the berries of *Juniperus oxycedrus* L. subsp. *oxycedrus* was analysed by GC/MS. Sixty-three compounds have been identified from which α -pinene (68.0%) and myrcene (12.1%) were the main constituents. We have also detected germacrene D (4.4%), abietadiene (1.0%), limonene (0.9%) and manoyl oxide (0.9%)

From 63 identified components in the oil of the berries of this Juniper, 32 were monoterpenes which represented 88% of the oil, 22 were sesquiterpenes summing 8% and 9 were diterpenes amounting 3.2% of the oil.

Other studies on the essential oil of *J. oxycedrus* have been done by different authors. Motl *et al.* (1960) cited ylangene and viridiflorol and De Pascual Teresa *et al.* (1977,1978) detected for the first time β - and γ -bulgarenes, but we have not found these compounds. Guerrero Hernández *et al.* (1987) cited monoterpenes camphene, sabinene, α -terpineol, 1,8-cineole, γ -terpinene, terpinolene, terpinen-4-ol, camphor, linalool, and the sesquiterpene α -copaene. We have detected these monoterpenes and also several sesquiterpenes like α -cubebene (0.4%), β -caryophyllene (0.4%), δ -cadinene (0.5%) and α -cadinol (0.3%) and diterpenes like manoyl oxide (0.9%), abietadiene (1.0%) and pimarinal (0.4%). It is worth mentioning the presence of germacrene D (4.4%) as the major sesquiterpene found in our oil. This compound had not been mentioned before in the literature.

De Pascual Teresa J, AF Barrero, A San Feliciano, MC Caballero (1977) *An. Quim* 73: 1527.

De Pascual Teresa J, AF Barrero, MC Caballero, A San Feliciano (1978) *An. Quim* 74: 966.

Guerrero Hernández E, MC López Martínez, R García Villanova (1987) *J. Chromatography* 396: 417-420.

Motl O, V Herout, F Sorm (1960) *Collect. Czech. Chem. Commun* 25: 1656.

P. 51. The bioactive essential oil of *Heracleum sphondylium* subsp. *ternatum* (Velen.) Brummit

Gökalp İşcan^a, Fatih Demirci^a, Mine Kürkçüoğlu^a, Merih Kıvanç^b, K. Hüsnü Can Başer^a

giscan@anadolu.edu.tr

^a Medicinal and Aromatic Plant and Drug Research Centre (TBAM), Anadolu University, 26470-Eskişehir, Turkey

^b Faculty of Sciences, Department of Biology, Anadolu University, 26470-Eskişehir, Turkey.

The essential oil of *Heracleum sphondylium* subsp. *ternatum* (Velen.) Brummit (Umbelliferae) was isolated from crushed seeds by means of hydrodistillation and analyzed by GC and GC/MS. Major components were identified as 1-octanol (50.3%), octyl butyrate (24.6%) and octyl acetate (7.3%). Furthermore, antimicrobial activity of the oil was evaluated using micro-dilution broth and agar diffusion methods. The bioactive constituent of the essential oil was determined as 1-octanol by using the bioautography assay.

P. 52. The ultrastructural features of plastids relative to synthesis of naphthoquinones in leaf glands cells in *Droseraceae* plants

L. E. Muravnik, Alexandra N. Ivanova
alyx@KD1537.spb.edu

Komarov Botanical Institute RAS, Prof Popov str., 2, 197376, St.-Petersburg, Russia

Different naphthoquinones are taxonspecific for genus of *Droseraceae* family. Plumbagin was found in *Drosera* (Durand, 1971), *Drosophyllum* (Durand, 1974), *Dionaea* and *Aldrovanda* (Culham, 1994) but 7-methyljuglone is peculiar to *Drosophyllum* and *Drosera* (Culham, 1994).

The most colored glands of unstimulated leaves were fixed with 3% glutaraldehyde or 2.5% glutaraldehyde with 2% formaldehyde in 0.1 M phosphate buffer pH 7.2 at room temperature and postfixed with 2% osmium tetraoxyde. After dehydration in alcohols material was prepared for electron microscopy by standard procedure.

Plastids in secretory cells of above-listed genera are leucoplasts. Their volume densities are similar within the error of calculations and come to 2.2%. In secretory cells of slime glands of *Drosophyllum lusitanicum* there are 9.0 plastids per cell section; one plastid section area is as small as 0.45 μm^2 . Stroma is condensed; it contains light thylacoids with vacuole-like dilatations. Peripheral reticulum consists of one or two elements parallel to plastid membrane. Tubular complex is frequent. Plastoglobuli are small and rare. Sometimes short elements of reticular envelope occur near plastids.

In secretory cells of tentacles of *Drosera rotundifolia* there are 2.2 plastids per cell section; one plastid section area is 0.53 μm^2 . Stroma is condensed, contain light thylakoids with local dilatations. In some plastids tubular complex forms dense accumulations. Periferal reticulum consists of single elements. Plastoglobuli are large and numerous. Plastids are enclosed in reticulum.

In secretory cells of digestive glands of *Dionaea muscipula* there are 3.7 plastids per cell section; one plastid section area is 0.15 μm^2 . Stroma is light; it contains tubular complex, single thylakoids, large plastoglobuli of different electron density, and small starch grains. Reticular envelopes near plastids were not found.

In secretory cells of digestive glands of *Aldrovanda vesiculosa* there are 1.2 plastids per cell section; one plastid section area is 0.87 μm^2 . Stroma is light. Thylakoids have small dilatations; they assembled in loose network. Tubular complex was not found. Plastoglobuli are small and rare. There is no reticular envelope around plastids but sometimes lipid bodies were found nearby plastid membrane.

On the strength of these data some considerations were made. Reticular envelopes around plastids in *D. lusitanicum* and *D. rotundifolia* cells indicate synthesis of secondary methabolites. Presence of reticular envelope distinguish both species containing juglone from that ones without it. Therefore it may be leucoplast reticular envelope which is related to juglone synthesis. Condensed stroma of *D. lusitanicum* and *D. rotundifolia* plastids may be due to high concentration of carbohydrates in cytoplasm. *D. lusitanicum* has maximal plastid number per cell section at the minimal area of plastid section among tested species therefore it has maximal total area of plastid surface. It may provide for greater intensity of synthesis and transmembrane transport of methabolites. In fact *D. lusitanicum* synthesizes plumbagin in greater amount then some other representatives of *Droseraceae* (Durand, 1974).

This work was supported by RSCI (grant No 01-04-48806).

Durand R, MH Zenk (1971) *Tet. Lett.* 32: 3009-3012.

Durand R, MH Zenk (1974) *Phytochemistry* 13: 1483-1492.

Culham A, RJ Gornall (1994) *Biochemical. Systematics & Ecology* 22: 507-515.

P. 53. Diurnal variation in the concentrations of volatiles in leaves and florets of *Thymus vulgaris* and *Salvia officinalis*

H. B. Jakobsen^a, L. P. Christensen^b

henrik.byrial@get2net.dk

^a Vibevadgaard, Blaesenborgvej 9, 4320 Lejre, Denmark

^b Department of Horticulture, Danish Institute of Agricultural Sciences, Research Center Aarslev, Kirstinebjergvej 10, DK-5792 Aarslev, Denmark

The major volatiles of *Thymus vulgaris* and *Salvia officinalis* leaves and florets were identified and quantified in order to detect diurnal oscillations in the concentration of volatiles in the tissues. This trial was conducted in an attempt to determine the optimal time of the day for harvest of the essential oil under Scandinavian conditions. The samples were collected at the initiation of flowering at 4 h intervals throughout the 24 h period. Leaves, open flowers and floral buds were extracted separately prior to GC-MS analysis.

In order to determine the optimal developmental stage for harvest of the essential oil, the entire plants were harvested from 1 m² plots at 14 d intervals from pre-flowering till maturation of the seeds. The theoretical yield of essential oil pr. Ha was evaluated after steam distillation.

Quality aspects of the oils together with the yield levels are discussed in order to evaluate the possibility for essential oil production in Scandinavia.

P. 54. Chemical composition of the essential oils of *Anthemis altissima* L.

K. Javidnia^a, R. Miri^a, M. Kamalinejad^b, H. Sarkarzadeh^a, Az. Jamalian^a

kjavidnia@albarta.com

^aFaculty of Pharmacy, The Medical Sciences University of Shiraz, Shiraz, Iran

^bDepartment of Pharmacognosy, Faculty of Pharmacy, The Medical Sciences University of Shaheed Beheshti, Tehran, Iran

Anthemis, is one of the most important genera of the Compositae family. The genus *Anthemis* is represented by 37 species grown wildly in Iran (Rechinger, 1982). *Anthemis nobilis* has been known from Roman times. It has been used as antispasmodic and sedative in folk treatment of digestive and rheumatic disorders (DerMarderosian, 2001). The essential oil of *A. nobilis* possess anti-inflammatory and sedative properties (Rossi *et al.*, 1988). *A. cotula* showed antimicrobial activity against both gram-negative and gram-positive microorganisms (Quarenghi *et al.*, 2000).

In the present work we report constituents of the essential oil of the flowers, leaves and stems of *Anthemis altissima* L. growing wild in Iran. The essential oils were isolated separately from flowers, leaves and stems of *A. altissima* and analyzed by GC-MS. A total of 123 components were identified in the oils of flowers, leaves and stems. The main components of the oil of both the flowers and leaves were β -caryophyllene (18.9%, 10.7%) and caryophyllene-oxide (9.0%, 17.0%), respectively. Nonterpenic compounds were the main components of the oil of stem. Palmitic acid (30.4%) and linoleic acid (27.9%) were the main compounds of the stem oil.

DerMarderosian A (2001): *The Review of Natural Products*. 1th edn, Facts and Comparisons, Missouri, pp. 518.

Quarenghi MV, ML Tereschuk, MD Baigori, LR Abdala (2000) *Fitoterapia* 71(6): 710-712.

Rechinger KH (1982): *Labiatae*. In: Edit. K.H. Rechinger, Austria, *Flora Iranica*, Vol. 150, 25-44, Akademische Druck-u. Verlagsanstalt, Graz.

Rossi T, M Melegari, A Bianchi, A Albasini, G Vampa(1988) *Pharmacol Res Commun* 5: 71-74.

P. 55. Seasonal variation in yield and composition of essential oil from *Artemisia vulgaris* L. growing wild in Poland

Jean Baptiste Nsanzimana, Danuta Kalemba

dakal@snack.p.lodz.pl

Institute of General Food Chemistry, Technical University of Łódź, Poland
90-924 Łódź, Stefanowskiego 4/10

Artemisia vulgaris L., the common mugwort, occurs wild in Europe and Asia. The herb has been used as a digestive in folk medicine and a flavouring agent in food and beverages. It is regarded as a thujone-containing plant although there are some chemotypes lacking this ketone.

The aim of the presented work is to investigate the yield and chemical composition of the mugwort essential oil during vegetation. Plants were collected in 3-weeks intervals from June to September 2001 in three places in Poland. The essential oils were obtained by hydrodistillation from the air-dried material and separated to fractions by vacuum distillation and flash chromatography. The chemical constituents were identified by GC-RI, GC-MS and ¹H-NMR.

The yield of the oil was low at the beginning of vegetation period (0.03%), increased up to 0.19% at the stage of buds (mid-July) than decreased gradually to 0.10% in full flowering and to 0.04% after flowering. Simultaneously with change of the yield great seasonal variation was observed in the chemical composition of the oil. Up to stage of bud the sesquiterpene hydrocarbons were the main constituents of all essential oils, germacrene-D (20-30%), β-caryophyllene (13-20%) and α-humulene (6%), being the major components. On the bud emergence, the composition of the oils changed suddenly and monoterpenes became the major constituents. 1,8-Cineole (10-15%), camphor, borneol and terpinen-4-ol (each 3-5%) were the main oxygenated monoterpenes and α-pinene and sabinene (each more than 5%) were the main monoterpene hydrocarbons in both stage of buds and full flowering. Other important components of mugwort essential oil were camphene, β-myrcene, artemisia ketone, α-copaene, β-elemene, bicyclogermacrene, spathulenol, β-caryophyllene oxide, T-muurolol, α-cadinol. It is noteworthy that even traces of thujones, which are considered as neurotoxic and limited by food regulations, were not found in mugwort essential oil.

P. 56. Phytopesticidal and repellency effect of four ethnobotanicals from N. E. India against mosquitoes and their GC/MS analysis

S. Phukan, M. C. Kalita

mckalita1@sancharnet.in

Dept. of Biotechnology, Gauhati University, Guwahati 781014, Assam, India

The use of botanicals as an alternative to the chemical compounds is gaining tremendous momentum because of its multifarious advantages. The essential oils of four species (*Acorus calamus*, *Cymbopogon flexuosus*, *C. winterianus* and *Ocimum americanum*) used traditionally by the various tribes of N.E. India have been subjected to bioassay against two mosquitoes species (*Culex quinquefasciatus* and *Aedes aegypti*) by the standard bioassay procedure (WHO/VBC/81.807). The essential oils were extracted using a Clevenger apparatus. Of the four species that were subjected to bioassay, *A. calamus* showed the highest potency in terms of pesticidal activity, with 80% mortality towards *A. aegypti* and 60% mortality in the case of *C. quinquefasciatus* at 25 ppm after 24 hrs. *C. flexuosus* also exhibited high potency, with 60% and 55% mortality against *A. aegypti* and *C. quinquefasciatus* respectively, within 6 hrs at 100. ppm. *C. winterianus* showed moderate efficacy with 70% and 20% mortality against *C. quinquefasciatus* and *A. aegypti* respectively, within 24 hrs at 50 ppm. The oil of *O. americanum*, which was reported to have pesticidal activity against some insects (Tawatsin *et al.* 2001), did not show any toxic effect to the target organisms. However, it showed a growth regulatory activity, with complete inhibition of pupal development even at 10% concentration. Similar effects are also seen for the other species, except for *A. calamus*, where 10% of *C. quinquefasciatus* and 15 % of *A. aegypti* formed pupae showed no subsequent adult formation.

The repellency effect of 55% and 60% was observed for *O. americanum* and *C. flexuosus*, after up to 4hrs, whereas *A. calamus* and *C. winterianus* exhibited 100% repellent activity after 1hr only. *C. winterianus* however showed a 70% repellency effect for *A. aegypti* and 80% for *C. quinquefasciatus*, after 3hrs.

The GC/MS analysis of *C. winterianus* oil showed 59 constituents, with citronellal constituting 26.5%, citronellol 11.4%, geraniol 20.3%, geranyl acetate 5.2%, nerol 0.5% and eugenol 0.5%. In the oil of *A. calamus* 40 constituents were analysed, with β -asarone as the major component, amounting to 82.6%. The analysis of *O. americanum* revealed 53 compounds of which thymol, constituting 52.9% of the total oil, was the main component. The essential oil of *C. flexuosus* showed 10 major compounds, with geranyl acetate (76.0%) being the main constituent.

Tawatsin A, SD Wratten, RR Scott, U Thavara, Y Techadamrongsin (2001) *J. Vector Ecol.* 26(1): 76–82.

P. 57. Essential oil of *Juniperus horizontalis* Moench. cultivated in Spain: seasonal variation of leaf and berry oil

Mohamed Kardali, María José Pérez-Alonso, Arturo Velasco Negueruela

kardali@bio.ucm.es

Dpto de Biología Vegetal I. Facultad de Biología, Universidad Complutense. 28040 Madrid. Spain

The volatile oils from leaves and green unripe-berries of *Juniperus horizontalis* Moench., an ornamental prostrant shrub (Peña Arribas and Peña Arribas, 1998), cultivated under gardening conditions in a central park in Madrid (Spain) were analysed by means of capillary GC and GC-MS in combination with Kovats indices. A seasonal investigation of both leaves and berries oils was also performed. Significant differences in yields and in chemical composition from a quantitative and qualitative point of view were detected when comparing all the oil samples analysed. Among the almost 60 components investigated, monoterpenes were dominant which is in accordance with data reported in a previous publication (Fretz, 1976). The sesquiterpenic fraction, relatively small, is characterized by high percentages of both cadinane-type sesquiterpenes and elemol/eudesmol-type; these results stand in sharp contrast with those previously published by Von Rudloff (1975). In addition, a small amount of diterpenes has also been detected.

The autumn-collected leaf oil was characterized by a high amount of monoterpenes namely: β -myrcene (16.9%), endobornyl acetate (19.4%) and sabinyl acetate (14.3%) followed by limonene (9.4%) and sabinene (9.3%) while, in decreasing order, α -thujene (3.7%), α -pinene (3.2%), β -thujone (2.4%), terpinen-4-ol (2.3%), camphene (2.0%), γ -terpinene (1.3%) and α -terpinolene (1.3%) were found to be in moderate amounts. Elemol (2.3%), δ -cadinene (1.9%) and α -cadinol (1.5%) were the main sesquiterpenes found.

On the other hand, the leaf oil chemical composition of the same plant collected in springtime of the following year has shown a significant increase of sabinene (19.2%) limonene (25.8%), γ -terpinene (4.2%), terpinen-4-ol (6.6%) and cadinol (4.0%) whereas β -thujone and endobornyl acetate have decreased to 2.4% and 4.4%, respectively.

Furthermore, berries oil was found to contain a very big amount of limonene (28.6%), terpinen-4-ol (11.6%) and myrcene (18.9%) when compared to autumn-leaf oil but has shown a slight variation with the springtime-leaf volatiles except with regard to terpinen-4-ol.

Fretz TA (1976) *J. Amer. Soc. Hort. Sci.* 101 (5): 611-613.

Peña Arribas J, J Peña Arribas (1998) *Coníferas ornamentales*. Floraprint, España Ed.

Von Rudloff E (1975) *Phytochemistry* 14: 1319-1329.

P. 58. Effect of essential oils for prevention of steel corrosion

Tatsuo Kawahigashi

L1kawahi@cced.kindai.ac.jp

Institute for Science and Technology, Kinki Univ., Higashi-Osaka, 577 Japan

Corrosion of steel in concrete is one of the damage-causes for the durability of the concrete structures. At present, several techniques are used for the prevention of the steel corrosion. For example, as for the prevention methods of the corrosion, "painting the surface of steel with the resin", "using the steel of the corrosion-resistant material" or "using the material of FRP (fiber reinforced plastic)" and so on are done. However, these are not excellent methods for prevention of the steel corrosion at present. Therefore, it is necessary to propose a new opinion of prevention of the corrosion of steel in concrete.

In this study, the steel corrosion in the saltwater environment was evaluated by the electrochemical technique. In salt water, the steel corrosion without being processed by essential oils was compared with being processed by essential oils on the steel surface.

From these examined results, this study is discussed about the prevention effect of essential oils against the corrosion of steel.

P. 59. Glandular trichomes and essential oil of *Salvia glutinosa* L.

Ayla Kaya, Betül Demirci, K. Hüsni C. Başer

aykaya@anadolu.edu.tr

Medicinal and Aromatic Plant and Drug Research Centre (TBAM), Anadolu University, 26470 Eskisehir, Turkey

The aerial parts of *Salvia glutinosa* L. bear indumentum with two types of trichomes: simple and multicellular nonglandular trichomes, and stalked and sessile dense glandular trichomes. Glandular trichomes are extremely long-stalked and dense on the stem and calyx surfaces. But, sessile glands are rare on the stem, calyx and leaf adaxial surfaces and dense on abaxial surface of the leaf. Secretion accumulates in a subcuticular space and is released to the outside by cuticle rupture. Water-distilled essential oil from dried aerial parts of *S. glutinosa* was analysed by GC and GC/MS. The main constituent was found as 1-octadecanol (11.0 %).

P. 60. Comparative Study of the Essential Oils of *Cymbopogon schoenanthus* ssp. *proximus* from Central Sudan

Hatil Hashim Elkamali ^a, Hassan ElSubki Khalid ^b, Babiker Fadlalla ^c Mohamed Ahmed Hassan ^d, Hizabr Hashim Elkamali ^e
htlkamali@yahoo.com

^aDept. Of Botany, Omdurman Islamic University, P.O. Box 382 Omdurman Sudan

^bDept. of Phytochemistry and Taxonomy, Medicinal & Aromatic Plants Research Institute, P.O. Box 11496 Khartoum Sudan

^cDept. Of Botany, University of Khartoum P.O. Box 321, Khartoum Sudan

^dDubai Police Crime Laboratory

^eAlwasl Hospital, Pharmacy Dept. 9115, Dubai UAE.

There are seven wild species of *Cymbopogon* in Sudan: *C. caesius*, *C. commutatus*, *C. excavatus*, *C. giganteus*, *C. nervatus*, *C. sennarensis* and *C. schoenanthus* ssp. *proximus* (Andrews, 1956).

Cymbopogon schoenanthus (L.) Spreng ssp. *proximus* (Hochst.ex A.Rich.) Maire and Weiller (Poaceae), Camel's Hay, known in Sudan as Maherib and in Arab Countries as Halfa Bur is a well known aromatic and medicinal plant of North and Central Sudan. In Sudan this plant is used as antispasmodic and for treatment of stomach and kidneys pain.

The chemical composition of Maherib oil (*Cymbopogon schoenanthus* ssp. *proximus*) was studied by Modawi (1975), El Gamal and Wolff (1987). In our present study, we analyzed the oils isolated by hydrodistillation from leaves collected at different regions of Central Sudan, i.e., El Damer (North Central), El Obeid (West Central) and El Gedaref (East Central). The oil yield is.

The oils obtained in a yield of 3.5%, 1.2 % and 2.1 % respectively, were analysed by GC/MS. Eleven components were reported for the first time. Pipertione, which identification was accomplished, with CNS excitatory, was the main component, ranging from 60.8% to 89.7 %. The major constituents were piperitone (60.8%), Δ -2-carene (11.4%) and elemol (6.4%) for El Obied sample, and piperitone (89.7%), α -terpineol (1.7 %) and β eudesmol (1.6%) for ElGedaref sample. Eldamer sample was dominated by piperitone (63.0%), elemol (11.1%) and Δ -2-carene (8.6%)

The essential oils from central Sudan are predominately monoterpenoid in character (75.5-93.8%), sesquiterpenoids being the only significant at 18.4 % of El Damer sample.

Upon comparing the composition of essential oils of the Mahereib Sudanese populations with that from plants with Indian origin, considerable intraspecific variations were noted (Shahi and Tava, 1993).

Andrews, FW (1956) The Flowering Plants of the Anglo- Egyptian Sudan Vol. (III) Buncl; Co. Arbroath.

El-Gamal, MH, Wolff, P (1987) Planta Medica. 53 (3): 293-294,

Modawi, BM (1975): Examination of the terpenoids of *Cymbopogon* species Ph.D. Thesis – University of Khartoum, Sudan

Shahi, AK and Tava, A (1993) J. Essent.oil Res. 5 (6): 639 – 643

P. 61. Variation in essential oil content and composition of *Levisticum officinale* Koch. from different origins

S. M. Kim^a, H. G. Chung^b, N. S. Seong^b, E. Nemeth^c

^aCollege of Industrial Science, Konju National University, Yesan Chung-Nam Republic of Korea

^bDivision of Industrial Crop of National Crop Experiment Station Rural Development Administration, 205 Seodun-dong Kwansunku Suweon, Republic of Korea.

^cDepartment of Medicinal and Aromatic plants, Faculty of Horticultural Sciences, Szeint Istvan University, Villany ut 29-35 Budapest, Hungary

The variation in the essential oil content of lovage (*Levisticum officinale* Koch.) from Korean and Hungarian origin was studied from 1997 to 1999. Different plant organs and different times of harvest during the vegetative and reproductive periods were also considered. The essential oils were isolated by hydrodistillation and their composition determined capillary GC method with standards. The essential oil content showed significant differences between the two populations on the vegetative organs. The essential oil level of the leaves and roots was considerably higher in the Korean population at full flowering and waxy ripening stage but essential oil content of the roots was significantly higher in the Hungarian taxon at leaf rosette stage. We observed the essential oil accumulation tendency was mainly dependent on plant organs and intra-specific taxon during the vegetation period.


Butylidene-phthalide was proved to be the main component of the oil in both population roots (50.9-73.3%), while dimethyl-acetate was showed as a major compound on the over-ground parts (56.7-62.0%). The qualitative composition of the essential oil in the reproductive organs concerning the identified compounds was the same as the vegetative parts with the main component α -phellandrene (4.8-28.1%) and butylidene-phthalide (9.7-16.1%). The quantitative composition showed some changes during the ontogenesis phases. Most characteristic ones are the decreasing proportion of dimethyl-acetate (from 7.3% to 1.1%) and the appearance of α -pinene (from 0.5 % to 1.5 %) only after fruit setting in both populations.

P. 62. Microbial transformation of (-)-carvoneFatih Demirci^{a*}, Neşe Kırimer^a, Yoshiaki Noma^b, Yoshinori Asakawa^c, K. Hüsni Can Başer^a

nkirimer@anadolu.edu .

^a Medicinal and Aromatic Plant and Drug Research Centre (TBAM), Anadolu University, 26470-Eskişehir, Turkey^b Faculty of Domestic Sciences, Tokushima Bunri University, Yamashiro-Cho180, Tokushima City, 770-8055, Japan^c Faculty of Pharmaceutical Sciences, Tokushima Bunri University, Yamashiro-Cho180, Tokushima City, 770-8055, Japan

The enantiomeric monoterpene (-)-carvone (**1**) was subjected to microbial transformation studies using the fungi *Abisidia glauca* ATCC 22752. After 4 days of incubation a new diol: 10-hydroxy-(+)-neodihydrocarveol (**4**) was formed. The absolute configuration and structure of this crystalline substance was identified with the aid of X-ray refractometry and supported by spectroscopic techniques such as MS, IR and NMR. Antimicrobial activity of the substrate and metabolite was also investigated against human pathogenic microorganisms.



*Part of the PhD thesis of Demirci F (2000), Eskişehir, Turkey.

P. 63. Quantitative data of plastidome in secretory canals of *Eleuterococcus senticosus*

M. R. Kolalite, A. N. Ivanova

milana_kolalite@mail.ru

Plant Resources laboratory, V.L. Komarov Botanical Institute, Prof. Popov str. 2, 197376, St.-Petersburg, Russia

Eleuterococcus senticosus Maxim. is a wide-spread species in the Russian Far East. Its young leaves, bark and roots are fragrant and are used by local population for non-alcoholic drinks, beer and sweets production and in seasonings, the essential oil of *E. senticosus* is used in the perfumery. Monoterpenes represent up to 0.8% of the secretion isolated from the roots and shoots (Wild useful plants of Russia, 2001). Leucoplasts are the most specific feature of epithelial cells in secretory canals, already in the meristematic zone. They become dominant in epithelial cells at the stage of secretion. As plastidome is proved to be an important compartment that takes part in monoterpene biosynthesis (Cheniclet and Carde, 1985) we have studied their morphological properties. Plastids in the secreting cells are elongated, have rather dense stroma and no ribosomes. One or two large inclusions enclosed in a membrane are present in plastids. We distinguished two membrane systems in plastids: well developed peripheral reticulum and vesicular membrane system located in the central part of plastid. Plastoglobuli are numerous and are associated with the vesicular membrane system. Starch grains are often present and they are electron dense as the starch grains of the plastids in parenchyma cells.

The quantitative parameters were: linear and square dimensions, the number and partial volumes of membrane systems, plastoglobuli, starch grains and inclusions. Five plastid types were found in secreting cells of *E. senticosus* according to the quantitative data. They differed by partial volume of the main plastidome features. Depending on plastid type the stroma occupied from 30.9% to 48.1%. Moderately dense inclusions were present in most plastids but occupied different partial volumes ranging from 2.9% to 30.0%. Electron comparatively dense inclusions bordered by a membrane were present only in 3 plastid types, but occupied, when present, a relatively large volume from 39.0% up to 47.7%. The partial volume of starch grains had negative correlation with the dense inclusion volume. They were absent in plastids with the largest partial volume of dense inclusions and in the plastids that had no such inclusions ranged from 7.4% to 33.3% per plastid volume. We interpret the described plastid types as a developmental succession and propose that the cells at the same spatial canal level have different developmental or functional status.

Plastids appear to participate actively in the secretory processes. They are always sheathed by endoplasmic reticulum. At least one, sometimes two electron dense inclusions, that we interpret as terpenes, and mitochondria are adjacent to plastids at secreting stage.

In the senescent part of the canals plastids are smaller, rounded, and they have neither vesicular membrane system nor starch grains. The inclusions occupy the most part of plastid volume. There are no associations between plastids, electron dense droplets of secretion and mitochondria.

We acknowledge the Russian Academy of Sciences for the financial support of the present study according to the 6th young scientists supporting program, grant N 245.

Wild useful plants of Russia (2001) Budantsev AL and EE Lesyovskaya (Ed.), Saint-Petersburg Pharmaceutical Academy Press, Saint-Petersburg. 663 p.

Cheniclet C, J-P Carde (1985) *Isr.J.Bot.* 34: 219-238.

P. 64. Evidence of cell divisions in the epithelial cells of *Pinus sylvestris* secretory canals

N. K. Koteyeva, M. R. Kolalite

milana_kolalite@mail.ru

Plant Resources laboratory, V.L. Komarov Botanical Institute, Prof. Popov str. 2, 197376, St.-Petersburg, Russia

Secretory canals are traditionally regarded as a character of great importance for the *Pinus* genera. They produce oleogumresin that is valued for its pharmaceutical and technical properties. The canals in *Pinus sylvestris* are present already in the buds and produce the secretion that takes part in buds protection during winter. Within a bud all the needles primordia are formed. In spring the bud starts to grow very rapidly. It elongates from the size 5-10 mm to 50-300 mm during 2-3 weeks. The bud grows both due to cell divisions in meristematic zones and to the cell linear dimensions increase. During the shoot growth the canal size increases greatly both in diameter and length. There may be two mechanisms of canal growth. The first is: the epithelial cell number grows only due to apical meristeme activity and below the apex the canal grows due to cells elongation. The second mechanism assumes that secretory cells may divide below the apex being already differentiated. In the literature we have found no evidence of cell division during the active secretion stage.

Longitudinal sections of *P. sylvestris* shoots, fixed in a mixture of glutaraldehyde and paraformaldehyde in June, when the shoot elongation is most evident, were examined using light and electron microscopes. Already in a bud the epithelial cells of the secretory canals have the features common for active secreting terpenoidogenic cells with hypertrophied leucoplastidome. Endoplasmic reticulum is present mostly in cisternal form and its membranes are associated with plastid envelope.

At a distance 3-4 cm from the apex canal lumen has maximal diameter, the cells surrounding it are moderately vacuolated. A large elongated or lobed nucleus occupies the central part of secretory cells. Plastids are located on both sides of the nucleus, they are very long, flexed or have invaginations. Plastid stroma is dense. Numerous osmiophilic droplets, which we interpret as terpenes, are located within plastid envelope. Different mitotic stages have been registered in the epithelial cells. The dividing cells had all the features of actively secreting cells. The number of such divisions found in different shoot samples allows regarding them as normal phenomenon.

The data obtained provides direct evidence of canal elongation due to both linear dimensions increase and cell divisions of differentiated epithelial cells. The capacity to divide at the stage and in the course of active secretion may be regarded as a unique feature in secretory canal epithelial cells. The main reason of this event may be a high growth rate of *P. sylvestris* shoot.

We acknowledge the Russian Academy of Sciences for the financial support of the present study according to the 6th young scientists supporting program grant N 245.

P. 65. Essential oil of laurel (*Laurus nobilis* L., Lauraceae) from MontenegroNada Kovačević^a, Mirjana Simić^a, Violeta Slavkovska^a, Mihailo Ristić^b

nadak@pharmacy.bg.ac.yu

^a Faculty of Pharmacy, Vojvode Stepe 450, 11000 Belgrade, Yugoslavia^b Institute for Medicinal Plant Research "Dr Josif Pancic", Tadeusa Koscuska 1, 11000 Belgrade, Yugoslavia

Laurel, bay laurel, *Laurus nobilis* L. (Lauraceae), is an ornamental tree, indigenous to the western as well as in the southeast part of Europe. The main products are leaf of laurel and essential oil obtained from the leaf by steam distillation. The leaf is chiefly used as spice in culinary and the essential oil in the food technology. Rarely, leaf of laurel is used as a traditional drug for digestive disorders. The leaf of laurel contains approximately 1-3 % of essential oil, sesquiterpenoid lactones and isoquinoline alkaloids. The main constituents of this oil are 1,8-cineole (30-70 %), linalool (3-17 %) and methyleugenol (2-8 %).

The essential oil of laurel is one of the main export products of Yugoslavia. The main area for collected leaves of wild growing laurel is the southeast part of Adriatic coast in Montenegro. The essential oil is produced by many little, private distilleries without standardised technology of distillation. Mainly, the essential oil is obtained by steam distillation of the young shoots, not pure leaves. Besides this, the request for quality of this essential oil still does not exist as an ISO or AFNOR standards. The Yugoslavian standard (JUS) is from the 1968. All this constitutes a major problem for laurel essential oil producers. This work deals with our effort to define the quality of leaf essential oil as one of the most export drugs from Yugoslavia and Montenegro. Besides, we made the comparison of the composition of essential oils obtained from different parts of laurel tree.

The autumn of 2000 was very weak with high daily temperature (up to the 20 °C) unusual for that part of the year. The laurel trees were in the second flowering phase just in period of October to November, the usual time for collecting leaves for commercial purpose. The samples of young shoots, bark and fruits were collected from the same tree, what was done at two different localities (Budva and Bar) in the Adriatic coast of Montenegro. The air-dried plant material was pulverized and the essential oil was obtained by hydrodistillation using a Clevenger-type apparatus. The content of essential oil was as following: 1.4 % in young shoots, 0.75 % in bark, 1.5 % in the separated leaves and 0.7 % in separated stems (locality Budva).

The chemical composition of these oils was determined by GC (FID) and GC/MS techniques. The main constituents of all investigated oils were, as it is usual for laurel, 1,8-cineol, methyleugenol and α -terpinyl acetate. Besides, α -pinene, β -pinene, sabinene and linalool were also present. It was interesting and important for commercial samples of laurel essential oil that there was no significant difference among the essential oils obtained from young shoots and separately from leaves and stems.

The main constituents of the essential oil obtained from the bark (from the trunk and branches) were 1,8-cineol (26%), methyleugenol (10%) and α -terpinyl acetate (9%). By comparison with the essential oil obtained from the leaves of laurel, the bark oil contained more methyleugenol, and the same level of α -terpinyl acetate but less 1,8-cineole, as well as α -pinene, β -pinene, sabinene and linalool.

The essential oil obtained from dry flowers of laurel contained the same components but in lower concentration: 1,8-cineol (15,7%), α -terpinylacetate (6.5%), methyleugenol (3.9%), sabinene (1.6%), linalool (1.4%), β -pinene (0.4%) and α -pinene (0.3%). Besides, the flower oil contained higher concentration of *trans*-caryophyllene (9.5%) and γ -muurolene (7.1%).

P. 66. Comparative study of the volatile compounds of different types of poppy seeds and -oils by SPME-GC-MS-Detection and by Olfactorimetry

Sabine Krist^a, Heidrun Unterweger^b, Gerhard Buchbauer^a, Franz Bandion^b

martina.hoeferl@univie.ac.at

^a Institute of Pharmaceutical Chemistry, University of Vienna, Althanstr.14, 1090 Vienna, Austria

^b Institute for Foodstuffs, Federal Office and Research Center for Agriculture, Spargelfeldstr.191, 1226 Vienna, Austria

The opium poppy (*Papaver somniferum*) is said to be the oldest recognised medicinal plant in the world and it is still very important, as it furnishes a most important painkiller, morphine. Analysis of the remains from the Neolithic age verified that poppy was known to cavemen living in the territories of Spain, France, Germany, Austria and Hungary 4-5 thousand years BC.

The poppy, more than any other medicinal species, has been used and misused by humankind from the early days of history. Even today, the production and sale of this plant is politically charged. In Europe, especially in Hungary (6000-10 000 ha), Czech Republic (7000-8000 ha), Slovakia (7000-8000 ha), Romania (3000-5000 ha) and the Netherlands (1000-1500 ha) poppy is cultivated for its edible and oil producing seed. The poppy never lost its therapeutic and economic importance.

Although numerous investigations have been carried out with respect to the constituents of poppy capsules and seeds, very little was yet known about the volatile components of poppy seeds and -oils, which contribute to their characteristic odour.

In this work, the distribution of components in the headspace of different types of poppy seeds and -oils from white, blue and grey-varieties and the influence of different storage conditions on their odour are shown. For this work SPME-GC-MS-Detection and Olfactorimetry were used.

P. 67. Variability of methyleugenol and estragole of Basil varieties in field and greenhouse cultivation

H. Krüger

h.krueger@bafz.de

Federal Centre for Breeding Research on Cultivated Plants, Institute for Plant Analysis, Neuer Weg 22/23, D-06484 Quedlinburg, Germany

There are two opinions of the Scientific Committee on Food of the European Commission (Council of Europe, 2001, 2002) which point out, that estragole und methyleugenol have been demonstrated to be genotoxic and carcinogenic. Therefore the existence of thresholds cannot be assumed and the Committee could not establish a safe exposure limit. But consequently, reduction in exposure and restrictions in use levels are indicated. It is known, that both substances can be constituents of basil essential oils. The content of estragole in these oils can be up to 40 %, that of methyleugenol up to 90 %.

Comparisons of varieties from 2001 show that the composition of the basil essential oils are not only determined genetically. They are also affected substantially by the cultivation conditions. The differences between the contents of methyleugenol of all varieties are particularly explicit when field and greenhouse cultivation are compared (Tab. 1). Estragole types were not contained in the 2001 assortment.

Tab.1: Contents of methyleugenol in the basil essential oils of different varieties under field and greenhouse conditions

	Contents of methyleugenol [%] in the essential oils of the varieties							
	Genova	Bavires	Aton	Bageco	Genua Star	Sanremo	Pesto	Genoveser
Field cultivation	1.1	0.6	0.7	0.0	0.5	0.0	0.7	0.3
Greenhouse cultivation	32.4	8.0	30.3	20.1	23.2	3.3	25.5	27.1

The results from 2001 are to be complemented by systematic investigations of the variability and ontogenesis from the 2002 assortment of 12 varieties cultivated in the field and in the greenhouse.

Council of Europe – Scientific Committee on Food (2001) Opinion of the Scientific Committee on Food on Methyleugenol (4-Allyl-1,2-dimethoxybenzene): SCF/CS/FLAVOUR/4 ADD1 FINAL 26 September 2001

Council of Europe – Scientific Committee on Food (2001) Opinion of the Scientific Committee on Food on Estragole (1-Allyl-4-methoxybenzene): SCF/CS/FLAVOUR/4 ADD2 FINAL 26 September 2001

P. 68.**Volatile constituents of *Sanguisorba officinalis***Karl-Heinz Kubeczka^a, Carsten Vollmann^b, Nahla Ayoub^c^a kubeczka@t-online.de^a Untere Steigstr. 12 b, D-97276 Margetshöchheim, Germany^b Dept. of Pharmaceutical Biology, University of Hamburg, Bundesstr. 45, D-20146 Hamburg, Germany^c Dept. of Pharmacognosy, Faculty of Pharmacy, Ain-Shams University, Cairo, Egypt

Sanguisorba officinalis L. belonging to the Rosaceae, subfamily Rosoideae is a semi-rosette shrub up to 100 cm with composite 1 to 2 cm dark red-brown ovate-oblong flower-heads. The rosette leaves are 20 to 40 cm long and consist of 7 to 15 ovate leaflets.

The plant is widespread in the northern, temperate regions of Europe, temperate Asia and North America and has been formerly used in folk medicine for staunching wounds. Phytochemical studies performed on the herb and the roots including the rhizomes led to the identification of numerous hydrolysable and condensed tannins, triterpene glycosides, sterols, and flavonoids including flavonoid sulphates, which seem to be characteristic of the genus *Sanguisorba*.

Since no previous reports concerning the chemical composition of the volatile constituents of *S. officinalis* were found, we have investigated the volatiles from the roots including the rhizomes and the fresh leaves obtained by hydrodistillation by means of GC and GC-MS.

In the yellowish root oil numerous constituents mainly belonging to monoterpenoids (34.7 %), sesquiterpenoids (16 %), and several fatty acids (16.8 %) were identified.

In contrast to the root oil the small amount of volatiles obtained from the leaves consisted mainly of degradation products of chlorophyll (phytol and phytadienes), aliphatic alcohols and aldehydes, alkanes and fatty acids. Terpenes were almost completely lacking.

P. 69. Analysis of a historical oil, Ninde oil (*Aeolanthus gamwelliae* Taylor) revisited

K. H. C. Başer, M. Kürkçüoğlu, B. Demirci

mkurkcuo@anadolu.edu.tr

Medicinal and Aromatic Plant and Drug Research Centre (TBAM), Anadolu University, 26470 Eskisehir, Turkey

Aeolanthus gamwelliae Taylor (Ninde) (Lamiaceae) is an aromatic plant native to northern Zambia. In 1930s, water and steam distilled oil from flowers of this plant was commercially available due to its high geraniol content. The cultivation of Ninde plant was also initiated in the region in the same years. Long ago, the cultivation practices have come to an end and the oil is not, at present, a commodity of commerce.

The oil subjected to this study was obtained from the collection of Natural Resources Institute (NCI), which had been produced by water and steam distillation in 1934/35.

The oil was analysed by GC and GC/MS. Forty one compounds were characterised representing 95% of the oil with geraniol (66%), geranyl acetate (12.5%), geranial (4%) and geranyl formate (3.7%) as main constituents.

P. 70. Leaf essential oil of *Laurus nobilis* L. from different localities of Croatia

I. Jerković, J. Mastelić, D. Kuštrak

igor@ktf-split.hr

Department of Organic Chemistry, Faculty of Chemical Technology, University of Split, N. Tesle 10/V, HR-21 000 Split, Croatia

Bay laurel, sweet laurel, *Laurus nobilis* L., an evergreen shrub or small tree native to the Mediterranean region and Asia Minor, has been admired for its beauty and aromatic leaves since ancient times. Currently, the plant is both cultivated and collected from the wild. Dried leaves are used to flavour meats, fish, vegetables, and soups and also as an ingredient in pickling spice. The essential oil, distilled from the leaves, with bactericidal and fungicidal properties (Schnaubelt, 1998) is used in perfumery and for flavouring food products (Duke, 1987). This is the first report on the Croatian bay leaf essential oil.

The leaves were harvested at three different localities in October 2001 (Table 1.). The dried leaves were crushed to powder and the oil was isolated by hydrodistillation, for 3 hours, in a modified Clevenger-type apparatus. The oil yield varied from 0.87 – 1.53 % (mass/dry weight). The isolated oils were analysed by GC-MS using two columns with different stationary phase polarity (HP-20M and HP-101). The most abundant oil compounds are listed in Table 1.

Table 1. Comparative chemical composition of *Laurus nobilis* L. leaf essential oils - main components (%)

Compound	Localities		
	Opatija	Rijeka	Krk-Kras
1,8-Cineole	32.8	30.9	33.6
Sabinene	8.7	10.2	10.7
α -Terpinyl acetate	10.2	8.6	9.1
Linalool	7.3	8.8	4.7
Methyl eugenol	4.8	5.6	8.8
Eugenol	3.4	3.9	2.8
α -Terpineol	3.5	4.6	3.7
Terpinene-4-ol	3.3	2.7	4.6
α -Pinene	3.3	3.8	4.7
β -Caryophyllene	1.6	2.2	0.7

Our results are comparable with those reported for Portugal (Roque, 1989), Turkey (Lawrence, 1990) and Italy (Caredda *et al.*, 2002) bay oils.

Caredda A, B Marongiu, S. Porcedda, C Soro (2002) *J. Agric. Food Chem.* 50:1492-1496.

Duke JA (1987) *Handbook of Medicinal Herbs*: 271-272.

Lawrence BM (1990) *Parfum. Flavor.* 15: 67-68.

Roque OR (1989) *J. Ess. Oil Res.* 1: 199-200.

Schnaubelt K (1998) *Advanced Aromatherapy*: 35-38.

P. 71. Steam-volatile leaf oils of some Western Australian species of the family MyrtaceaeErich V. Lassak^a, Joseph J. Brophy^b^a Phytochemical Services, 254 Quarter Sessions Road, Westleigh NSW 2120, Australia^b School of Chemistry, University of New south Wales, Sydney NSW 2052, Australia

The volatile leaf oils of nine Western Australian species have been extracted by means of cohobative steam-distillation and analysed by GC/MS.

The leaf oils of *Agonis obtusissima* F. Muell., *Eremaea pauciflora* (Endl.) Druce, *Kunzea pulchella* (Lindl.) A. S. George and *Malleostemon tuberculatus* (E. Pritzel) J. W. Green have not been previously investigated whilst *Thryptomene australis* Endl. and *Thryptomene kochii* E. Pritzel have received only cursory attention. Additional data have been obtained for *Melaleuca uncinata* R. Br., *Eucalyptus erythrocorys* F. Muell. and *Eucalyptus stoatei* C. Gardner.

The oils of *Kunzea pulchella* (both the red and the white flowered forms) contain globulol as the main constituent (83-88%). The oils of *Thryptomene australis* are rich in geranic acid (ca 52%) and α -pinene (22%) whilst *Thryptomene kochii* contains α -pinene as its main constituent (58-60%). *Eremaea pauciflora* oil contains 1,8-cineole (22%) and the three isomeric eudesmols (α -, β -, and γ -, 26%). *Agonis obtusissima* contains substantial amounts of α -pinene (12%), *trans*- β -ocimene (16%) and globulol (39%). *Malleostemon tuberculatus* oil contains α -pinene (33%), 1,8-cineole (21%) as well as E,E-farnesol (6%) as its main constituents.

The oil of *Melaleuca uncinata* obtained from bushes growing near Jaurdie contains about 89% of α -pinene and represents the 4th chemical form of this species postulated by Watson (1943/44). The three other forms are characterised by: 1,8-cineole and the three eudesmol isomers; 1,8-cineole and no eudesmols (Brophy *et al.*, 1990); and terpinen-4-ol (Brophy and Lassak, 1992).

The composition of the leaf volatiles of *Eucalyptus erythrocorys* and *Eucalyptus stoatei* obtained by steam-distillation differ in the former case, but are comparable in the latter, with results published on vacuum extracted oils (Bignell *et al.*, 1994, 1996).

Bignell CM, PJ Dunlop, JJ Brophy, JF Jackson (1994) *Flavour Fragr. J.* 9: 113.

Bignell CM, PJ Dunlop, JJ Brophy, JF Jackson (1996) *Flavour Fragr. J.* 11: 145.

Brophy JJ, EV Lassak (1992) *Flavour Fragr. J.* 7: 27.

Brophy JJ, EV Lassak, DJ Boland (1990) *Flavour Fragr. J.* 5: 43.

Watson EM (1943/44) *J. Roy. Soc. Western Australia* 30: 83.

P. 72. Influence of extraction methods on the composition of essential oilsÉ. Lemberkovics^a, Á. Kéry^a, B. Simándi^b, A. Kakasy^a, É. Szöke^a

lembi@drog.sote.hu

^a Semmelweis University, Faculty of Pharmacy, Department of Pharmacognosy, Budapest, Hungary, Üllői út 26., H-1085^b Budapest University of Technology and Economics, Department of Chemical Engineering, Budapest, Hungary, Műegyetem rkp. 3. K ép., H-1521

The aim of this presentation is to show our results on comparison of composition of the essential oil fractions obtained by traditional steam distillation and supercritical fluid extraction. The latter technique avoids the degradative heat processes, hydrolysis, isomerisation and racemisation. It is a special separation technique to obtain extracts and fractions of natural origin for therapeutical or cosmetic uses (Chouchi *et al.* 1995, Bicchi *et al.* 1999).

The plant material for the various extraction methods was selected from the *Asteraceae* and the *Lamiaceae* families. For the supercritical fluid extraction (SFE) carbon dioxide was used as supercritical solvent. The extracts were collected by stagewise precipitation in two separators. The waxy product and extract rich in essential oil were collected in the 1st and in the 2nd separator respectively. The traditional water steam distillation (SD) was carried out according to the Hungarian Pharmacopoea 7th ed. and the GC analysis performed on capillary silica fused columns coated with DB-1701 and the specific chiral columns coated with Rt- β DEXm or Rt- β DEXsm.

Comparing the composition of steam distilled oils with that of volatile SFE fractions the following general characteristics were established. The SFE fractions were richer in monoterpene-esters and poorer in alcohols than the traditional essential oils (clary sage, lavender, dragonhead). Regarding the distribution of the monoterpene and sesquiterpene compounds, the SFE fractions contained sesquiterpene hydrocarbons in higher percentage than the distilled oils (*Salvia triloba*). Moreover, the proportion of sesquiterpenes increased in SFE fractions collected successively in time (*Salvia officinalis*) similarly to the ratio of oxygenated monoterpenes to monoterpene hydrocarbons (*Rosmarinus officinalis*). In more cases it was verified that a part of mono- and sesquiterpenes were present originally in bounded form (glycosides) in plants. So they appeared only in essential oil fractions after previous acidic treatment (*Thymus*, *Origanum*, *Satureja* species) [Oszagyán *et al.*, 1996, Lemberkovics *et al.*, 1998]. During the supercritical extraction the azulene sesquiterpene lactones did not transform to azulenes in camomile and yarrow, but the non volatile SFE fraction of some *Asteraceae* plants (feverfew, blassed thistle) contained sesquiterpene- γ -lactons of unchanged structure.

Acknowledgements: The work was supported by Hung. Nat.Sci.Found (OTKA T 30034).

Bicchi C, A Binello, P Rubiolo (1999) *Phytochemical Analysis* 10: 17-21.

Chouchi D, D Barth, E Reverchon, DG Porta (1995) *Ind.Eng.Chem.Res.* 34: 4508-4513.

Lemberkovics É, Á Kéry, B Simándi, G Marczal, M Then (1998) *Proceedings of Supercritical Fluid Material and Natural Products* 2: 567-572.

Oszagyán M, B Simándi, J Sawinsky, Á Kéry (1996) *J.Essent. Oil.Res.* 8: 333

P. 73. Indumentum features and essential oil composition of the Hungarian *Salvia* species

M. Then, É. Lemberkovics, G. Marczal

Thenm@drog.sote.hu

Semmelweis University, Faculty of Pharmacy, Department of Pharmacognosy, Budapest, Üllői út 26., H-1085 Hungary

In this work we report on the anatomical features of the indumentum of four *Salvia* species (Lamiaceae), *S. officinalis* L., *S. sclarea* L., *S. pratensis* L., *S. nemorosa* L, namely the structure and position of glandular and covering trichomes, as well as on the composition of their essential oils.

The chemical composition of the oils was investigated in the different plant organs (leaf, calyx and petal) of the various species. The plants used in this study were cultivated in the garden of the Ecological and Botanical Research Institute of the Hungarian Academy of Sciences. The microscopical studies were carried out by Axioscop apparatus. The essential oils were isolated by the methods described in the Hungarian Pharmacopoea and analysed by gas chromatography.

The indumentum of each organ of the *Salvia* species investigated is composed by glandular and covering trichomes. The characteristic Labiatae-type glandular trichomes occurred along with other glandular trichomes, composed of a unicellular head. Numerous glandular and covering trichomes can be seen on the calyx and petal. Their structure is similar to those of *S. officinalis*. A few of glandular trichomes are on the petals, too. A lot of whip-like trichomes are present, but conical unicellular trichomes are rare. The big covering trichomes, having short basal cells and a sharp terminal cell are typical of the *S. sclarea* leaf. Some conical covering trichomes are also found. The leaf of *S. pratensis* is poor in trichomes. There are rarely multicellular covering and glandular trichomes with unicellular head.

Comparing the essential oil composition of the fresh and dried plant organs of the four *Salvia* species, differences exist only in *S. officinalis*. The dominant components in the fresh leaf oil are the sesquiterpenes α -humulene (27,9%) and β -caryophyllene (13,0%). In dried leaf oil the monoterpene thujone isomers were the main ones. The qualitative oil composition of the three organs does not differ from each other, although some qualitative differences are recorded. The thujone isomers are dominant (17-20%) in the leaf and calyx oils but they represent only 5,9% of the petal oil. The main component in petal oil is the pharmacologically active β -pinene (24,4%). The oil from *S. pratensis* is dominated β -caryophyllene (7,1%) and γ -muurolene (24,0%). Their ratio in the calyx and petal oils is similar, but in the leaf oil this is reversed.

The structure of covering trichomes and the percentage composition of the oils provide a good basis for the differentiation of *Salvia* species.

Hegnauer R (1962) *Chemotaxonomie der Pflanzen* IV. 295-296.

P. 74. Comparative pharmacology of essential oils with a floral to pungent odour

Maria Lis-Balchin^a, Stephen Hart^b

lisbalmt@sbu.ac.uk

^aSchool of Applied Science, South Bank University, Borough Road, London, SE1 0AA, UK

^bMessengers and Signalling Research Group, School of Biomedical Sciences, King's College London, Hodgkin Building, Guy's Campus, London, SE1 9RT, UK

Essential oils and absolutes with a floral odour like lavender, geranium and jasmine all have a spasmolytic effect on the guinea-pig ileum *in vitro* which is most likely mediated through the secondary messenger cAMP. The oils of a number of floral-smelling *Pelargonium* species are also similar, but the more pungent, camphoric species, though still exhibiting a spasmolytic effect, show a mechanism which is more likely to be through calcium channel blockade.

Peppermint oil is also similar to the florals, as is one of the so-called pungent Tea tree oils from New Zealand, Manuka (*Leptospermum scoparium*). However the other tea tree oil from New Zealand, Kanuka (*Kunzea ericoides*) and the Australian tea-tree (*Melaleuca alternifolia*) has an initial spasmogenic action followed by a spasmolytic response. Valerian oil, which can only be described as pungent again behaves spasmolytically like the floral oils but appears to act through calcium channels like the pungent *Pelargonium* oils. Oils like citrus, frankincense, nutmeg, dill and fennel show a very strong contraction. Two 'catty-like' Buchu oils, *Agathosma betulina* and *A. crenulata* showed differences in the mode of action although both showed an initial spasmogenic action followed by spasmolysis. The former alone showed an action via cAMP.

The mode of action on smooth muscle appears to be directly related to the chemical composition of the oils. Spasmolysis is effected by EOs with high concentrations of alcohols, aldehydes, esters, ketones and sesquiterpenes, regardless of the actual component. The spasmogenic effect is effected by EOs with high concentrations of monoterpenes. The actual mechanism of the spasmolytic effect is more complicated but seems to involve the floral or pungent odours of the oils.

The pharmacological effect on the uterus *in vitro* is remarkably constant, showing a decrease in spontaneous contractions by all EOs studied. This raises the question of safety when using oils during childbirth as they could invoke cessation of uterine contractions and therefore prove a danger to the baby's life.

The action on skeletal muscle, using the rat isolated phrenic nerve diaphragm preparation or chick biventer muscle, is more complex. Some EOs show a decrease in the twitch response alone, like lavender. Some like dill, frankincense and nutmeg show a contracture with a decrease in the twitch response, others like thyme show a contracture without the effect on the twitch response. Camphor caused an increase in the twitch response and angelica root showed no response.

All these aspects will be discussed and illustrated.

P. 75. New odoriferous terpenoids containing bicycle [3.1.0] hexane skeleton using (+)-3-carene as a starting substrate

Bożena Frąckowiak, Stanisław Lochyński

lochyński@kchf.ch.pwr.wroc.pl


Institute of Organic Chemistry, Biochemistry and Biotechnology, Wrocław University of Technology, W. Wyspiańskiego 27, 50-370 Wrocław, Poland

This paper is a continuation of our investigations on elaboration of simple methods for the synthesis of new chiral odoriferous compounds when using monoterpene hydrocarbon (+)-3-carene **1** (a major constituent of Polish turpentine, the essential oil from *Pinus sylvestris* L.) as a starting material.

In our previous paper we reported syntheses of odorants with preserved carane system (Lochyński, 2002), as well as those with di- or trimethylbicyclo[3.1.0]hexane system substituted at C-2 (Lochyński, 2001) and C-3 positions (Lochyński, 1997).

Here we present further study starting from allylic alcohols **2** and **4**, which were subjected to the Claisen rearrangement (orthoacetate modification) giving γ,δ -unsaturated esters and their derivatives **3** and **5**.

All newly obtained terpenoid compounds possess interesting olfactory properties. Odor characteristics will be presented with special emphasis on structure-activity relationship.



Acknowledgements: Supported by State Committee for Scientific Research, Grant No 6 P06B 031 20

Lochyński S, *J. Soc. Cosmet. Chem.*, 1997, 48, 107-116

Lochyński S, Frąckowiak B, Olejniczak T, *32nd International Symposium on Essential Oils*, Wrocław (Poland), September 9-12, 2001, P 99

Lochyński S, Kowalska K, Wawrzeńczyk C, *Flavour Fragr. J.*, 2002, 17, 181-186

P. 76. Aromatic species with economic potential on the cerrado- Goiás, Brazil

Lucia Lopes¹, Marcus Vinicius de Miranda Martins²

lulopes7@terra.com.br; mvmm67@terra.com.br

SAS Quadra 05 Bloco H Sala 204 Brasília - DF/Brazil CEP 70070-914. Centro de Desenvolvimento Sustentável- CDS/UnB

The Brazilian Cerrado (Savannah) is rich in plants species that have economic potential, most of them unexplored. The increase of agricultural areas plus uncontrolled extractives actions is gradually reducing the natural areas where medicinal species grow wild. O Biome Cerrado (Brazilian Savannah) has 2 million of km², representing 23% of Brazilian territory. It is the second vegetation type of Brazil. Unfortunately, on last the 30 years this Biome has been losing its genetic material unknown scientifically mainly because of devastation. Local groups use the plants and sell products made from it on local markets making perfumes, crafts and liquors extracted from flowers, leaves, roots and bark.

A pilot research project is in course at the Public School Zeca de Farias working with 180 children and their families to increase the popular interest on this matter. Besides the educational purposes, this project focuses on economic interest in collecting seeds and creating greenhouses. The population of this school is originally from rural areas and this pilot project may help them to reconnect with their roots. The methodologies used are interviews and field work, collecting seeds and information about the place where is the mother tree, popular name, its use and how to use it and what part of the plant can be used. The plants, which show the most interest for the population, will be planted in the school ground and/or in the city.

Alto Paraíso is located at the Northeast portion of the State of Goiás, in the microregion denominated Chapada dos Veadeiros, occupying an area of 2603,4 km². The altitudes vary between 1180 and 1200 meters in the São Bartolomeu river basin. It is part of the territory São Jorge's District and the Rural Town of the Moinho. Due to its altitude the climate it is characterised by presenting soft summers and winters, with temperatures varying between 18°C and 20°C and the distribution of rains is seasonal.

A lack of economic alternatives pushed the small landowners to sell their properties to entrepreneurs who buy their land to introduce business forcing them to live in the city. The tourism, an important potential on the region, is make with no planning. The National Park, which has tourism and research potential, has lost its protected territory. In 1961 was approximately 625 thousand hectares, in 1981 was reduced to 65 thousand hectares.

Species	Popular Name	Active Principles	Use
<i>Caryocar brasiliense</i> Camb.	pequi	several substances were on essential oils like 1,8 cineole	liquors, home made food
<i>Achyrocline satureoides</i> D.C.	macela	essential oils and flavonóides	Aromatics pillows and shampoos
<i>Lychnophora ericoides</i> Less.	arnica	essential oils	Perfumery and cosmetics
<i>Spiranthera odoratissima</i> A. St. Hill.	manacá	essential oils	dry flowers, tea and source of jasmine
<i>Brosimum gaudichaudii</i> Trec.	mama-cadela	furonocumarinas	aromatic for pipe, food and medicine

¹ Coordenação da Especialização em Educação e Gestão Ambiental e Doutoranda em Desenvolvimento Sustentável/CDS/UnB

² Agrônomo e Membro do Cerrado *in Vitro*/FINATEC

P. 77. Seasonal essential oil variation of *Aniba canelilla* (H.B.K.) Mez

Francisca Socorro N. Taveira^a, Waterloo Napoleão de Lima^b, Eloisa Helena A. Andrade^c, José Guilherme S. Maia^c

gmaia@museu-goeldi.br

^aUniversidade Federal do Maranhão, Departamento de Química, 65080-040 São Luis, MA, Brazil

^bUniversidade Federal do Pará, Departamentos de Química, 66075-900 Belém, PA, Brazil

^cMuseu Emílio Goeldi, Departamento de Botânica, CP 399, 66040-170, Belém, PA, Brazil

Aniba canelilla is a high tree known as “casca-preciosa” (precious bark) and an important and historical species in the Amazon region because it was confused with cinnamon-trees during the voyage of Pizzaro and Orellana from the Andes to the Amazon estuary about 1540, and during the Humbolt and Bonpland’s 1800 expedition to find the “famous cinnamon” of the Orinoco River. This work was part of a project to identify biogeochemical markers of mineral soils in the Amazon, based on essential oil analysis (Taveira, 1991).

The essential oils of leaves, stem bark and trunk wood of *Aniba canelilla*, collected in the rainy and dry season from different soil types, were obtained by hydrodistillation and analysed by GC-MS.

The odoriferous principle of leaf, bark and trunk wood of *Aniba canelilla*, responsible for the cinnamon odour, is 1-nitro-2-phenylethane, rare in the nature. We observed that methyleugenol is also an important volatile constituent in the oil of *A. canelilla*. The percentage content of these two compounds was depending of the season time. In the rainy period the 1-nitro-2-phenylethane reach values near 95%, while methyleugenol remain below 18%. By contrast, in the dry period 1-nitro-2-phenylethane decrease to 39%, while methyleugenol reach 45%. The leaf oils produced by specimens collected at different soil types in the dry season showed the lower percentage contents for 1-nitro-2-phenylethane (39.0%) and methyleugenol (0.5%). On the other hand, the mono- and sesquiterpene compounds present in the same oils showed the higher percentage contents (13.5% and 43.2%, respectively). The presence of small quantities of benzaldehyde, benzene acetaldehyde, benzonitrile and benzene acetonitrile identified mainly in the essential oils of leaves of *A. canelilla* is unusual and noteworthy, and they could be the end-products of phenylalanine degradation. This latter, probably the origin of 1-nitro-2-phenylethane and methyleugenol and both compounds seems to be biologically interchangeable depending on the impact of extrinsic factors.

Acknowledgements: supported by MCT-PPG7/European Community.

Taveira FSN (1991) Plantas Aromáticas da Serra de Carajás Como Prováveis Marcadores Biogeoquímicos de Solos Mineralizados. Chemistry M.Sc Thesis, Pará University, 155p, Belém.

P. 78. *Stachys sylvatica* L.: secreting trichomes and essential oil

L. Maleci Bini ^a, A. Paolillo ^a, M. Antonelli ^a, B. Tirillini ^b

maleci@unifi.it

^a Dipartimento di Biologia Vegetale, Via La Pira 4, 50121 Firenze, Italy

^b Istituto di Botanica, Università di Urbino, Via Bramante 28, I.61029 Urbino, Italy

Stachys sylvatica L. is a perennial herb belonging to the *Stachys* section (Ball, 1972) and is widespread in all the Italian peninsula in mountain and moist places. The whole plant emanates an unpleasant smell and is covered by non glandular and glandular trichomes, which give it a velvety look. The plant is used for medicinal purposes in Italian folk medicine (Gastaldo, 1987).

In this work the different types of trichomes present on the plant (leaves, stem and flowers) and the essential oil composition (determined for the first time), are reported.

The trichome study was carried out using light and electron microscopy (S.E.M and T.E.M). Peltate trichomes, which are considered characteristic of Labiatae species secreting essential oil (Werker, 1993; Hallahan, 2000), are lacking. Instead, capitate trichomes occur as three types, as follows:

1) short capitate, characterised by one cell stalk, two secreting cells and a small subcuticular space;

2) long capitate, characterised by a stalk constituted of two or three cells, four secreting cells and a large subcuticular space containing the secretion;

3) very long capitate, characterised by a stalk constituted of three or more cells, six or more secreting cells, each of them bearing a small subcuticular space on the apex. This type of trichomes is probably characteristic of the *Stachys* genus. Indeed, it has been described only for some species of this genus (Falciani *et al.*, 1995). The number of the secreting cells, which constitute the head, depends on the species considered; however, the trichome morphology is the same in all the species examined. These trichomes usually are present on the inflorescence.

Concerning the substances secreted by the different types of trichomes, the histochemical analyses indicate that essential oils are present in all the trichomes. Moreover, in the short capitate trichomes a polysaccharidic secretion is present too.

The ultrastructure observations are consistent with the histochemical data. Indeed, both types of long capitate trichomes show plastids with dense stroma and numerous tubules containing electrondense material and smooth endoplasmic reticulum, i.e. typical organelles of a lipophile secretion (Bourett *et al.*, 1997; Ascensao *et al.*, 1998).

The oil analysis was carried out, separately on leaves and inflorescences, by Gas Mass Spectrometry. Preliminary results indicate a similar oil composition for both parts of the plant analysed. Germacrene D is the main component.

Ascensao L, N Marques, MS Pais (1997) *Int. J. Plant Sci.* 158: 249-258.

Ball PW (1972) *Stachys* L. In: Tutin T.G. *et al.* (eds.) *Flora Europaea* Vol. 3 Cambridge, University Press.

Bourett MT, RJ Howard, DP O'Keefe, DL Hallahan (1994) *Int. J. Plant Sci.* 155: 623-632.

Falciani L, L Bini Maleci, M Mariotti (1995) *Bot. J. Linn. Soc.* 119: 245-256.

Gastaldo P (1987) *Compendio della Flora Officinale Italiana*. Piccin. Padova.

Hallahan DL (2000) *Advances in Botanical Research. Plant trichomes*. 31: 77-120. Academic Press. San Diego.

Werker E (1993) *Flavour Fragr. J.* 8: 249-255.

P. 79. Essential oils from *Pothomorphe umbellata* and *Pothomorphe peltata*

K. Shiroma^a, A. F. Ferri^a, V. R. Vasconcellos^b, J. P. F. Teixeira^a, M. Moraes^c, J. I. Gomes^d,
Márcia Ortiz M. Marques^a

mortiz@cec.iac.br

^a Laboratório de Produtos Naturais, Instituto Agronomico- Brazil, ^b Lasefi – DEA/FEA – Universidade Estadual de Campinas- Brazil, ^c Departamento de Farmácia, Universidade Federal do Pará- Brazil, ^d CPATU- Embrapa – Amazonia Oriental - Brazil

The genus *Pothomorphe* Miq. belongs to the Piperaceae family and is composed by two species: *Pothomorphe peltata* (L.) Miq. and *Pothomorphe umbellata* (L.) Miq. These medicinal plants occur in Brazil, being *P. peltata* from the western and Amazon region, and *P. umbellata* from the southern region of the country. *P. peltata*, whose synonyms are *Hecheria peltata* Kunth (L.) and *Piper peltata*, is commonly named “caápeba do Norte” or “malvarisco” (Yunker, 1973); there is a popular use of its leaves as an anti-inflammatory, cholagogue, choleric, diuretic, vermifugal and anti-malarial (Milliken, 1997). *P. umbellata*, also known as *Hecheria umbellata* (L.) Kunth and *Piper umbellatum* L. is commonly named “caápeba do Sul” or “paripoba” (Yunker, 1973), and is traditionally used as cholagogue and choleric being recommended by the Brazilian Pharmacopoeia (Silva, 1929). Due to the similarity in their morphology, these plants are often mistaken for one another. However, morpho-anatomic studies report the following differences between them: in the insertion of the petiole, and in the occurrence and distribution of trichomes and stomata (Moraes, 1999).

Phytochemical studies of the essential oil from the leaves from *P. umbellata* collected in Brazil shows the occurrence of phellandrene, asarone and dillapiole (Hammer and Johns, 1983). The chemical composition of *P. peltata*, from the city of Belém, Pará State, showed the predominance of sesquiterpenes (Moraes, 1999). We describe herein the chemical composition of the leaf essential oils from *P. peltata* and *P. umbellata* as a contribution to the characterization of these species and in order to improve their use as medicinal plants.

P. peltata and *P. umbellata* were grown at the experimental farm at Instituto Agrônômico in Campinas city - São Paulo State, southern Brazil. The *P. peltata* seeds came from the Amazon region (city of Belém Pará State). The essential oils were extracted by steam distillation of the leaves for ca 75 minutes. The analyses were performed using a GC-MS unit (Shimadzu, QP-5000), equipped with a fused silica capillary column DB-5 (J & Scientific, 30 m x 0.25 mm x 0.25 µm). The carrier gas was Helium (1.7 mL/min), with the following temperature program: 50°C - 280°C at 4°C/min. The injector and detector temperatures were 240°C and 230°C, respectively. The identification of the oil components was made by comparison of their mass spectra with those from an instrument library (Nist62.lib), and/or from literature along with the retention indices.

The chemical profile from these two plants were quite different; *P. umbellata* showed the predominance of monoterpenes (92,3%) while *P. peltata* possessed ca.100% of sesquiterpenes. The most abundant components in the oil from *P. umbellata* were β-myrcene (4.2%), α-phellandrene (2.8%), Δ-3-carene (77.7%) and terpineole (3.2%); the main sesquiterpenes were β-caryophyllene (0.9%), δ-cadinene (0.9%) and γ-cadinene (0.5%). The major components of the oil from *P. peltata* were β-caryophyllene (49,1%), germacrene-D (30,2%) and trans-β-guaiene (7,5%).

Hammer MLA, EA Johns (1983) *J. Ethenopharm*, **40**: 53-75.

Milliken W (1997) *Econ. Bot*, **51**: 212-237.

Moraes, M (1999) PhD thesis, Universidade Estadual Paulista, São Paulo, Brazil.

Silva RAD. (1929) *Pharmacopeia dos Estados Unidos do Brasil*, São Paulo, Editora Nacional, 649.

Yunker TG. (1973) *Hoehnea*, São Paulo, Brazil, **3** : 29-284.

P. 80. Essential oil constituents as taxonomic markers in *Ocimum* spp.

Priscila C. Fernandes ^a, Márcia Ortiz M. Marques ^a, João P. F. Teixeira ^a, Pedro R. Furlani ^b, Vanessa R. Vasconcellos ^a

mortiz@cec.iac.br

^a Laboratório de Produtos Naturais, Instituto Agronomico de Campinas (IAC), Brasil

^b Centro de P&D de Solos e Recursos Agroambientais, IAC, Brasil

The species *Ocimum minimum* L. and *O. basilicum* L. are related to each other and commercially known as cultivars of a same species, *O. basilicum*, popularly referred to as basil. But papers such as from Grayer *et al.* (1996a) and Grayer *et al.* (1996b) reported that in crossing experiments of these two materials, there was a production of no viable seeds, evidencing the fact that these plants should be two different species, both rich in essential oil. The cross-pollination, very frequent in *Ocimum* spp, leads to a large number of sub-species and great genetic variability (Marotti *et al.*, 1996). Plant secondary metabolites can be influenced by genetic, physiological and environmental factors. The aim of this work is to contribute to the chemosystematic of the *Ocimum* genus, and for this purpose, essential oils profiles were determined for the leaves of both species (*O. minimum* and *O. basilicum*) cultivated under two conditions (organic mineral substrate and hydroponics system) and collected during the inflorescence phase, at two different dates.

The essential oils were extracted by steam distillation for 90 minutes. The extracts were analyzed by GC-MS (Shimadzu, QP-5000), with a fused silica capillary column DB-5 (J & Scientific, 30m x 0.25mm x 0.25 μ m). Helium was used as carrier gas (1.0 mL/min), and the oven temperature was programmed to increase from 60°C to 280°C at 3°C/min. The injector and detector temperatures were 220°C and 240°C, respectively. The identification of the oil constituents was confirmed by GC-MS, through the comparison of the instruments library (Nist62.lib), data from literature (McLafferty and Stauffer, 1989) and retention indexes (Adams, 1995). The statistical analysis was processed using Minitab- Statistical Software – 13 Demo.

The essential oils presented linalool as the major component, followed by α -trans-bergamotene, germacrene D, γ -cadinene, and cubenol. Among the cited substances, it is observed a significant difference in the quantity of linalool and α -trans-bergamotene, between the studied species. The linalool content in *O. minimum* varied from 44.3 to 54.3%, while in *O. basilicum* varied from 22.7% to 37.4%. For α -trans-bergamotene, the inverse result is observed: 6.6% to 9.5% for *O. minimum*, and 13.2 to 19.7% for *O. basilicum*. The variations within a same species were due to differences in the collecting time and growing system. The stepwise regression analysis, $\alpha=0.15$, showed that the difference in the α -trans-bergamotene content explains 81.4% of the differences between the studied species and adding the linalool and γ -cadinene contents, it is possible to explain 89.84% and 96.60%, of the difference between the species, respectively.

These results, associated to the phenotypic differences between *O. minimum* and *O. basilicum*, support the conclusion that linalool, α -trans-bergamotene and γ -cadinene can be used as taxonomical markers for the studied species.

Adams RP (1995) Identification of essential oils by gas chromatography/mass spectroscopy, Carol Stream, Allured Pub. Co. USA, 469p.

Grayer RJ, GC Kite, FJ Goldstone, SE Bryan, A Patont, E Putievisky (1996a) Phytochemistry 43 (5): 1033-1039.

Grayer RJ, SE Bryan, NC Veitech, FJ Goldstone, A Patont, E Wollenwebber (1996b) Phytochemistry 43 (5): 1041-1047.

Marotti M, R Piccaglia, E Giovanelli (1996) J. Agricult. Food Chem. 44: 3926-3929.

McLafferty FW, DB Stanffer (1989) Registry of Mass Spectral Data, New York. John Wiley - Interscience Pub.

Teixeira JPF, MOM Marques, PR Furlani, R Facanali (2000) Horticultura Brasileira 18: 982-983.

P. 81. Comparative Study of the Essential Oils of Three *Stachys* Species Grown Wild in Iran

Shiva Masoudi^a, Abdolhossein Rustaiyan^b

shmasoudi @ Yahoo.com

^aDepartment of Chemistry, Islamic Azad University, Central Tehran Branch, Tehran, Iran

^bDepartment of Chemistry, Islamic Azad University, North Tehran Branch, Tehran, Iran

The genus *Stachys* (Lamiaceae) is found in mild regions of the Mediterranean and in Southwest Asia. The genus *Stachys* consists of about 200 species, 34 are described in the flora of Iran, of which 13 are endemic (Mozaffarian 1996). *Stachys* species have several folkloric uses, e.g. the leaves of *S. officinalis* L. Trev. are used as a cholagogic, carminative and to relieve headaches. *Stachys* species have also been studied in biosystematic and chemotaxonomic studies in which flavonoids, quinones, iridoids, phenolic acids and diterpenoids were reported. The oils of the genus *Stachys* have been the subject of only a few studies.

As part of our research on aromatic flora of Iran, we investigated the chemical composition of the oils isolated by hydrodistillation from the aerial parts of *Stachys pilifera* Benth., *S. acerosa* Boiss. and *S. benthamiana* Boiss., which are endemic to Iran, by means of GC/MS in combination with retention indices.

In *S. pilifera* oil, 30 components representing 88.9% of the total oil were identified. *cis*-Chrysanthenyl acetate (25.2%) and *trans*-verbenol (19.7%) were the major compounds.

S. acerosa oil contained also *cis*-chrysanthenyl acetate (41.0%) and linalool (23.5%) among the 20 constituents characterised comprising 92.1% of the total components detected. Germacrene-D (16.8%), linalool (16.6%) and β -caryophyllene (11.0%) were the main constituents among the 20 characterised comprising 91.2% of the total components detected in the oil of *S. benthamiana*.

As can be seen the above information two oils (*S. pilifera* and *S. acerosa*) were rich in oxygenated monoterpenes, while the oil *S. benthamiana* consisted mainly of sesquiterpenes.

Mozaffarian V (1996) *A Dictionary of Iranian Plant Names*, P. 522-523 Farhang Moaser, Tehran, Iran.

P. 82. The effects of external factors on the quality and quantity of the camomile essential oil

Abdolnasser Massoudi

nasser_massoudi@yahoo.com

Dept. of R&D, KAF Co., P. O. Box 15815-3478, Tehran, IRAN

German camomile [*Chamomilla recutita* (L.) Rauschert] is one of the most important plants that are used widely in pharmaceuticals and cosmetics.

This research was performed to obtain the optimum growth conditions, the effects of soil texture and nutrient on flower crop and essential oil. In addition influence of IAA and different climates and different harvesting times on the quality and quantity of chamomile oil, was investigated.

The results show that the best soil texture for camomile growth is loam. This soil type leads to a flowering time decreasing of 29 days and a flower crop increasing of 376.9 %, in relation to plants cultivated under standard conditions.

IAA affects on plant growth and the quality of oil show that the flowering time was decreased in 20 days and essential oil yield and chamazulene content increased 58.3% and 33 %, respectively.

Cultivation of the plant in different climate influences affected also the camomile essential oil yield.

Different harvesting time (07:00; 12:00; 19:00 hours) showed that the best time for harvesting is at 12:00 hours.

P. 83. Composition of the essential oil from the leaves of *Pittosporum undulatum* from the Azores

J. R. Medeiros^a, H. Medeiros^b, L. B. Davin^c, N. G. Lewis^c

medeirosjmr@hotmail.com

^aCentro de Investigação de Recursos Naturais, Universidade dos Açores, 9502 Ponta Delgada, Açores, Portugal.

^bInstituto de Inovação Tecnológica dos Açores, Estrada de São Gonçalo, 9504-540 Ponta Delgada, Açores, Portugal.

^cInstitute of Biological Chemistry, Washington State University, PO Box 646340, Pullman, WA 99164-6340

The composition of the essential oil obtained by hydrodistillation from the leaves on the flowering phase of *Pittosporum undulatum* Vent. (Sjogren, 1984) growing on San Miguel island (Azores) was investigated, with constituents being identified by GC/MS analysis (Van Den Dool, 1962; Davies, 1990).

The oil was found to contain monoterpenes, sesquiterpenes, diterpenes and alkanes, of which the sesquiterpenes calamenene (41.4%), farnesol (10.9%), spathulenol (5.6%) and β -selinene (5.2%) and the diterpene (8 β ,13 β)-kaur-16-ene (10.7%) were the major components.

Tests for the detection of antithrombin activity (Medeiros, 2000) of the crude oils also revealed that these had even better activity than heparin.

Balasubrahmanyam VR, Rawat AKS (1990). *Econ. Bot.*, 44, 529–530.

Dato'Yaacob KB, Ariffin S (2000). *J. Essent. Oil Res.*, 12, 205–206.

Davies NW (1990). *J. Chromatog.*, 503, 1–24.

Medeiros JR, Macedo M, Constanca J, Nguyen C, Cunningham G, Miles DH (2000). *J. Ethnopharmacol.*, 72, 157–165

Ramanandraibe V, Rakotovo M, Andriamaharavo RN, Bessiere JM, Ravaonindrina N, Ramanoelina ARP. (2000). *J. Essent. Oil Res.* 12, 650–652.

Sjogren E (1984). *Açores, Flores*, Faial, Açores, Portugal.

Van Den Dool H, Kratz PD (1962). *J. Chromatogr.*, 11, 463–471

P. 84. Composition and antimicrobial activity of the essential oils from leaves and flowers of *Hedychium gardnerianum* from the Azores

J. R. Medeiros^a, L. B. Campos^b, S. C. Mendonca^b, L. B. Davin^c, N. G. Lewis^c

medeirosjmr@hotmail.com

^a Centro de Investigação de Recursos Naturais, Universidade dos Açores, 9502 Ponta Delgada, Açores, Portugal

^b Instituto de Inovação Tecnológica dos Açores, Estrada de São Gonçalo, 9504-540 Ponta Delgada, Açores, Portugal

^c Institute of Biological Chemistry, Washington State University, PO Box 646340, Pullman, WA 99164-6340, USA

The composition of the essential oils, obtained by hydrodistillation, from the leaves and flowers of *Hedychium gardnerianum* Sheppard ex Ker Gawler (Sjogren, 1984) growing on San Miguel island (Azores) was investigated. The compounds were identified by GC-MS analyses (Adams, 1995). There are studies about the composition and characteristics of the essential oils of various *Hedychium* species (Gottlieb, 1959; Haggag, 1972), but only the composition of oils from *H. gardnerianum* rhizomes has been examined (Weyerstahl, 1998). The oils in the leaves and flowers were rich in α -pinene, β -pinene and α -cadinol. Their potential antimicrobial activities were tested against *Staphylococcus aureus*, *S. epidermis* and *Pseudomonas aeruginosa* by the agar diffusion method (Tanaka, 1992). The oils had the highest activity against *S. aureus* and *S. epidermis* but had no activity against *P. aeruginosa*.

Adams R (1995) *Identification of Essential Oils by Gas Chromatography/Mass Spectroscopy*. Allured Pub. Co., Carol Stream, Illinois, USA

Gottlieb OR, MT Magalhães (1959) *Quim.*, 18: 179-180.

Haggag MY, AM El-Shary (1972) *Egypt. Pharm. Sci.*, 18 (4): 465-476.

Sjogren, E (1984) *Açores flores*. Horta, Açores, Portugal.

Tanaka Y (1992) In: *The Search for Bioactive Compounds from Microorganisms*, Omura S, Ed., Springer-Verlag, 30-44.

Weyerstahl P, H Marschall, K Thefeld, GC Subba (1998) *Flavour Fragr. J.*, 13 (6), 377-388.

P. 85. Glycosidic derivatives of volatile components for cosmetic applications

Chantal Menut, Alain Leydet, Huguette Agnanié, Jean Louis Montero

cmenut@univ-montp2.fr

Laboratoire de Chimie Biomoléculaire UMR 5032, ENSCM, 8, rue de l'École Normale 34296 Montpellier Cedex, France

Years after the first positive identification of geraniol β -D-glucoside in rose flowers (Francis and Allcock 1969), glycosidically bound forms of several volatile components have been identified in various other natural sources: these structures may be regarded as non-volatile flavour and fragrance precursors.

On the other hand use of natural ingredients in cosmetic preparations are more and more appreciated. For example, essential oils are not only used in cosmetics for perfuming purpose but also for their biological properties (antioxidant, antielastasic...) (Etienne, 1998).

Synthesis of glycosidic derivatives of the three main constituents of the "rose scent": citronellol, geraniol and 2-phenylethanol is described. L-Rhamnose was selected for its ability to target keratinocyte lectins.

These components should find interesting applications in cosmetic formulation, owing to a better hydrophilic lipophilic balance as well as a progressive release of "the rose scent" by cutaneous microflora action.

Etienne JJ (1998) *20th IFSCC Congress*, September 14-18, Cannes (France).

Francis MJO, Allcock C (1969) *Phytochemistry*, **8**, 1339.

P. 86. Antioxidant activity of some *Thymus* species grown in PortugalM. Falcato Simões^a, M. G. Miguel^b, A. C. Figueiredo^c, José G. Barroso^c, Luis G. Pedro^c, L. Carvalho^a

mgmiguel@ualg.pt

^a Universidade de Trás-os-Montes e Alto Douro, Quinta de Prados, 5000 Vila Real, Portugal^b Faculdade de Engenharia de Recursos Naturais, Universidade do Algarve, Campus de Gambelas, 8000-117 Faro, Portugal^c Centro de Biotecnologia Vegetal, Dep. de Biologia Vegetal, FCL, C2, Campo Grande, 1749-016 Lisbon, Portugal

Lipid peroxidation in biological membranes leads to cellular damage, which is involved in the development of chronic diseases that are life limiting. These include atherosclerosis, heart diseases, carcinogenesis and ageing processes. The oxidation of lipids in foodstuffs also affects their quality due to the volatile compounds formed, limiting the shelf life of fresh and processed food. Synthetic antioxidants such as butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), gallats and *tert*-butylhydroquinone (TBHQ), and natural antioxidants such as tocopherol and ascorbate derivatives, are largely used in food processing. In addition to these compounds essential oils from various aromatic plants have also been the subject of an increasing interest as antioxidants more health friendly. In this work, the chemical analysis and the antioxidant activity of the essential oils isolated from *Thymus mastichina*, *Th. zygis*, *Th. caespititius* and *Th. camphoratus* were analysed. The antioxidant index percentage was calculated using the formula $(1-t/c) \times 100$, where *c* is the absorbance value obtained in a fully oxidised control assay and *t* is the absorbance of the test sample, when a modified thiobarbituric acid reactive species (TBARS) assay was used. The antioxidant activity was carried out in the absence and presence of the radical inducer 2,2'-azobis(2--amidinopropane) dihydrochloride (ABAP). Different concentrations of essential oils (62.5, 125, 250 and 500 ppm) were used. Egg yolk was used as substrate. In the absence of ABAP, BHT was found to be the most efficient antioxidant, at all the tested concentrations. The antioxidant index percentages obtained with the essential oils, in concentrations of 250 ppm and 500 ppm, isolated from *Th. caespititius* and *Th. zygis* revealed to be as important as α -tocopherol and even superior when compared with BHA. The essential oils of *Th. mastichina* showed the less effective antioxidant effect. In the presence of ABAP, α -tocopherol and BHA were the most effective antioxidants, with similar antioxidant index percentages at 500 ppm. Within the essential oils, only *Th. zygis* showed the better antioxidant activity, reaching the highest AI% at 500 ppm, but significantly below the values achieved with α -tocopherol and BHA. The oils from *Th. mastichina* in a concentration range of 62.5 to 250 ppm and those from *Th. caespititius* at a concentration of 62.5 ppm showed pro-oxidant activity (Table 1).

Table 1. Antioxidant index percentage (AI%) of the essential oils, BHT, BHA and α -tocopherol using TBARS assay with and without ABAP

		500 ppm	250 ppm	125 ppm	62.5 ppm
Without ABAP	<i>Thymus mastichina</i>	39	32	17	10
	<i>Thymus zygis</i>	83	73	57	11
	<i>Thymus caespititius</i>	76	73	43	3
	<i>Thymus camphoratus</i>	52	38	16	14
	BHT	88	82	79	74
	BHA	66	65	61	60
	α -tocopherol	74	75	71	0
With ABAP	<i>Thymus mastichina</i>	5	-10	-15	-20
	<i>Thymus zygis</i>	53	33	11	0
	<i>Thymus caespititius</i>	42	26	2	-36
	<i>Thymus camphoratus</i>	37	34	9	7
	BHT	69	66	51	42
	BHA	83	74	64	55
α -tocopherol	82	76	67	51	

P. 87. Antioxidant activity of the essential oils of *Thymbra capitata*, *Thymus mastichina* and *Th. camphoratus* using sunflower and peanut oils as substrates

M. Falcato Simões^a, M. G. Miguel^b, A. C. Figueiredo^c, José G. Barroso^c, Luis G. Pedro^c, L. Carvalho^a

mgmiguel@ualg.pt

^a Universidade de Trás-os-Montes e Alto Douro, Quinta de Prados, 5000 Vila Real, Portugal

^b Faculdade de Engenharia de Recursos Naturais, Universidade do Algarve, Campus de Gambelas, 8000-117 Faro, Portugal

^c Centro de Biotecnologia Vegetal, Dep. de Biologia Vegetal, FCL, C2, Campo Grande, 1749-016 Lisbon, Portugal

The autooxidation of lipids in foodstuffs affects their quality, as it is responsible for the development of rancidity. Butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), gallats and *tert*-butylhydroquinone (TBHQ) are synthetic antioxidants commonly added to foods to prevent autooxidation. Nevertheless, these substances have been questioned because they show some toxicity and carcinogenicity. The demand for natural antioxidants is growing in order to render food products safer for mankind. Essential oils are receiving attention as potential antioxidants, due to their importance in food preservation.

In the present work, the chemical composition of the essential oils isolated from the leaves of *Thymus mastichina*, *Thymus camphoratus* and *Thymbra capitata* grown in Portugal and collected during the vegetative phase of the plants, was examined by GC and GC/MS. The essential oils (0.1 g/100 g) were also tested in order to evaluate their antioxidant ability on different substrates (peanut and sunflower oils). BHT and BHA were used as reference compounds for comparison. A blank assay was prepared under the same conditions, without adding any antioxidant. The rate of oxidation was followed by periodic determination of the peroxide index in the oils stored at 60 °C, for 78 days.

The essential oils isolated from *T. capitata* were dominated by carvacrol (71.9 %), whereas 1,8-cineole (49.4 %) was the major component of the essential oils from *Th. mastichina*. In contrast, linalool (16.6 %), linalyl acetate (15.2 %) and 1,8-cineole (10.8 %) were the three main compounds detected in the oils from *Th. camphoratus*.

For peanut oil, BHT was found to be the most effective antioxidant leading to a peroxide index value of 52.3 meq O₂/kg at the end of the assay. Concerning the essential oils, during the same time, values ranging from 142.3 meq O₂/kg (*Th. camphoratus*) to 152.0 meq O₂/kg (*T. capitata*) were found. For sunflower oil, considering the end of the experiment (78th day), neither BHT (193.0 meq O₂/kg) nor BHA (204.3 meq O₂/kg) revealed to be effective antioxidants. The lowest peroxide values, 170.7 and 172.3 meq O₂/kg, were found for oils of *Th. mastichina* and *Th. camphoratus*, respectively. It is also noteworthy the better performance of all tested essential oils in preventing the oxidation of peanut oil, when compared with that recorded for sunflower oil, that, to some extent, can be related to the lipid composition of these two substrates (Table 1).

Table 1. Antioxidant effect (peroxide index, meq O₂/kg oil) of the essential oils, BHT, BHA and blank on peanut and sunflower oils, stored in the dark at 60°C, at day 78th.

	Peanut oil	Sunflower oil
<i>Thymus mastichina</i>	145.0	170.7
<i>Thymus camphoratus</i>	142.3	172.3
<i>Thymbra capitata</i>	152.0	181.7
BHT	52.3	193.0
BHA	155.7	204.3
Blank	172.3	200.7

P. 88. Chemical composition and radical scavenger capacity of fennel (*Foeniculum vulgare* Mill.) fruits essential oil and fractions

Neda Mimica-Dukić^a, Sebastijan Kujundžić^b, Olga Tzakou^c, Maria Couladis^c

dukaned@eunet.yu

^a Institute of Chemistry, Faculty of Natural Sciences, University of Novi Sad, Trg Dositeja Obradovića 3, 21000 Novi Sad, Yugoslavia

^b DSP Chromatography, Bulevar AVNOJ-a 44a, 11070 Belgrade, Yugoslavia

^c Department of Pharmacognosy and Chemistry of Natural Products, School of Pharmacy, University of Athens, Panepistimiopolis Zografou, 157 71 Athens, Greece

Fennel (*Foeniculum vulgare* Mill, Apiaceae) is a well-known medicinal plant, native to the Mediterranean, but extensively cultivated all over the world (Wichtl, 1994). In phytotherapy the fruits (*Foeniculi fructus*) and essential oil (*Foeniculi aetheroleum*) are used in catarrhs of the upper respiratory tract, as a secretomotor, secretolytic, and antiseptic expectorant, as a spasmolytic and carminative in mild digestive disorders. Externally it is used as an eye lotion (decoction) and in functional visual disorders (Wichtl, 1994; Blumenthal, 1998). In the present study we investigated the radical scavenger capacity (RSC) of fennel fruit essential oil and fractions obtained from it.

The ground fennel fruits were submitted to steam distillation in a Clevenger apparatus. The obtained Essential Oil (F) was submitted to liquid chromatography in a glass column with teflon stop (60 cm; 3 cm i.d.) filled with silica gel. The sample was chromatographed with 200 ml pentane, 200 ml pentane: Et₂O (1:1), and 200 ml Et₂O. Fractions of 50 ml were collected, obtaining 12 fractions. The fractions were analysed by TLC on silica gel sheets, and similar fractions were put together obtaining 6 fractions (F1, F2, F3, F4, F5 and F6). The chemical composition of the oil and its fractions was analysed by GC/MS. The oil and the obtained fractions were investigated on their RSC measuring spectrophotometrically the disappearance of DPPH^{*} at 515 nm according to the procedure described by Soler-Rivas *et al.*, 2000.

According to the results of the GC/MS analysis, the fennel fruit essential oil (F) comprise phenylpropanoids (76.1%), oxygenated monoterpenes (16.8%) and monoterpene hydrocarbons (6.2%), with (*E*)-anethole (72.3%), fenchone (16.4%) and methyl chavicol (3.8%) as the main compounds. The obtained fractions showed the following chemical profile: F1- monoterpene fraction with limonene as the main compound; F2 – phenylpropanoid fraction with (*E*)-anethole as the main compound; F3 – same as fraction F2; F4 – aromatic fraction with *p*-anisaldehyde as the main compound; F5 – fraction with similar amounts of oxygenated monoterpenes, benzyl derivatives and phenylpropanoids, with (*E*)-anethole and *p*-anisaldehyde as the main component; F6 – phenylpropanoid fraction with (*E*)-anethole as the main compound. The essential oil of fennel fruits showed very low RSC reducing the DPPH^{*} radical for 3.00% at the concentration of 125 µg/ml, while (*E*)-anethole as the main compound had even a smaller RSC. The fractions tested at the same concentration levels had RSC of 0.00, 0.00, 1.03, 20.20, 30.14 and 21.57% for F1, F2, F3, F4, F5 and F6, respectively.

According to these results it can be concluded that the carriers of RSC in the Fennel essential oil are the oxygenated monoterpenes, and that the reduction of the phenylpropanoid and benzyl derivatives content has a positive impact on RSC.

Blumenthal M (ed.) (1998) *The Complete German Commission E Monographs*. American Botanical Council, Austin, Texas.

Soler-Rivas C, JC Espin, HJ Wichers (2000) *Phytochemical Analysis* 11: 330-338.

Wichtl M (1994) *Herbal Drugs and Phytopharmaceuticals*. Medpharm Scientific Publishing, Stuttgart.

P. 89. Essential oil of *Stellaria media* L. as possible contact dermatitis allergen

Neda Mimica-Dukić^a, Maria Couladis^b, Marina Jovanović^c, Boza Pal^d, Bisererka Mihajlović^e, Andrea Tot^a

dukaned@eunet.yu

^a Institute of Chemistry, Faculty of Natural Sciences, University of Novi Sad, Trg D.Obradovica 3, YU-21000 Novi Sad

^b Department of Pharmacognosy and Chemistry of Natural Products Panepistimiopolis Zografou, 157 71 Athens, Greece

^c Clinic for Dermatovenerologic Diseases. Clinical Center, Novi Sad, Medical Faculty, Hajduk Veljkova 1-3. YU-21000 Novi Sad

^d Institute of Biology, Faculty of Natural Sciences, University of Novi Sad

^e Public Health Institute, Medical Faculty, University of Novi Sad, YU-21000 Novi Sad

Stellaria media (L.) Vill (Caryophyllaceae) is a plant widespread all over the world, and considered as one of the commonest weeds. In folk medicine is used as diuretic, astringent, secretolytic and cough alleviating (Fournier, 1948). Although the medicinal properties of this plant is known for a long time, there are only limited reports on its chemical composition, mainly related to the flavonoid compounds (Budzianowski and Pakulski, 1988, 1991). In recent clinical study some cases of contact allergic dermatitis, type of erythema multiform, associated with this plant are described (Jovanović and Poljački, 2001). Plant contact dermatitis, is primary provoked by the formation of the hapten (electrophiles)-protein (nucleophiles) conjugate. Among natural haptens, sesquiterpene lactones are of particular significance, responsible for allergenic reaction of Compositae plants. However, the other terpenes such are limonene, Δ^3 -carene, terpineol, α - and β -pinene, could also act as contact dermatitis allergens (Benezra et al., 1985). Taking into account that lower terpenoids, abundantly presented in essential oils, could be responsible for allergenic reaction of *Stellaria media*, in the present study the chemical composition of essential oil was investigated.

The essential oil, isolated by hydrodistillation from the dried aerial parts of *Stellaria media*, collected in Vojvodina (North Serbia), was obtained in a yield of 0.02%, and analysed by TLC and GC/MS. Forty-three compounds were identified, representing 93.9% of the total oil, monoterpene being the major components. Among them, 1,8-cineole was dominant (25.3%), but considerable ratio of *p*-cymene (12.4%), nepetalactone (13.4%) and camphor (9.1%) was also found. Among sesquiterpenes, (*E*)-caryophyllene and (*E*)-caryophyllene oxide were the most abundant ones. The obtained results suggest that monoterpenes, especial 1,8-cineole and the iridoide nepetalactone, could be responsible for the allergenic properties of *Stellaria media*. A detailed dermatological study using the essential oil and target compound is in the course.

Benezra C, G Ducombs, Y Sell, J Fousseran (1985) Plant Contact Dermatitis. B.C.Decker, ed., The C.V. Mosby Comp. Saint Lous, Toronto, London.

Budzianowski, J, G Pakulski (1988) Planta Med. 54: 576-577

Budzianowski J, G Pakulski (1991) Planta Med. 57: 290-291

Fournier P (1948) Le Livre des Plantes Medicinales Et Veneneuses de France. (III). Encyclopedia Biologique . P. Lecevalier Edt.Paris, pp. 453-454.

Jovanović M, M Poljački (2001) Societatea Romana de Dermatologie 46: 25-6.

P. 90. Morphometry of glandular trichomes and yield of essential oils from *Lippia alba* (Mill) N. E. Br. ex Britt & Wilson (Verbenaceae) in function of the season and location of the leaves on the stem

Dulce M. Castro^a, Lin C. Ming^a, Márcia O. M. Marques^b, Silvia R. Machado^c

dulce@saloonet.com.br

^a Department of Vegetable Production - FCA/Unesp / Botucatu - SP

^b Center of Genetics, Molecular Biology and Fitoquímica of the Agronomic Institute of Campinas (CGBMFq - IAC)

^c Departamento de Botany-I.B. /Unesp/Botucatu

The present work had as objective verifies possible variations in the morphometrical of the trichomes main glandular revenue and essential of oil *Lippia alba* in function of the station make year and position of the leaves any branch. The parameters seasonal variation were considered (spring, summer, autumn and winter) and the location of the leaves in the axis stem (apical, medium and basal). Extractions of essential oil were accomplished through hydrodistillation. For the structural characterisation of the glandular trichomes, samples of the blade were included in resin glycol methacrylate and processed conventionally for analysis in through light (LM). The analysis morphometrical of the glandular trichomes was accomplished starting from eletronmicrografic obtained in scanning electron microscopy (SEM), digitate and analysed through the system of analyses KS300. The obtained data were submitted to statistical analyses (Tukey 5%). The largest revenue of the oil was obtained in the summer, in the leaves of the part apical of the branch; in these leaves, in the same station, the number of trichomes peltate was significantly larger, in both surfaces foliate; independent of the station and of the location of the leaves, the abaxial epidermis presented larger amount of glandular trichomes. They happened variations with relationship to the diameter and area of the head of the trichomes peltate between the different stations and location of the leaves, however, such variations were not significant. In the conditions of this work, there was a positive correlation between the amount of peltate trichomes and the revenue of essential oil.

P. 91. Composition of the essential oil of *Lonicera nummularifolia* Jaub & SpachR. Miri^a, K. Javidnia^a, A. Jafari^b, R. Sabet^a

mirir@sums.ac.ir

^aFaculty of Pharmacy, Shiraz university of Medical Sciences, Shiraz, Iran.^bCentral Research of Natural Resource and Animal Husbandry, Yasuje, Iran

The genus *Lonicera* belonging to the Caprifoliaceae family is represented by six species grown in Iran. *L. nummularifolia* (syn: *L. persica*) is distributed in Anatolia, Iraq, Afghanistan and many parts of Iran (Rechinger, 1982). There are many reports on the pharmacological activity of *L. japonica*. The aqueous extract of *L. japonica* showed anti-inflammatory effects (Lee, 2001). Polyphenolic compounds isolated from *L. japonica* thurb inhibited platelet activation and showed cytoprotective effect on hydrogen peroxide-induced cell injury (Chang, 1992).

In this study the essential oil of *Lonicera nummularifolia* Jaub & Spach was analysed by GC/MS. The yield of the oil was 0.04% (w/w). Eighty-three compounds represent 98.83% of the total oil. The main monoterpene component was linalool (2.9%), which can be seen in *L. japonica*, as the main component (Ji, 1990). The essential oil of plant consisted mainly of sesquiterpenes (80%). The main constituents of the oil were spathulenol (21.6%), α -muurolol (14.1%), E-caryophyllene (6.8%), α -cadinol (5.4%) and bicyclogermacrene (5.1%).

Chang WC, FL Hsu (1992) Prostaglandins leukot Essent Fatty Acids, 45: 307-12.

Ji L, J Pan, Z Xu (1990) *Zhongguo Zhong Yao Za Zhi.*, 15: 680-2.Lee JH, WS Ko, YH Kim, HS Kang, HD Kim, BT Choi (2001) *Int. J. Mol., Med.*, 1: 79-83.Rechinger KH (1982) *Caprifoliaceae*. In: Edit., K.H.Rechinger. Austria, *Flora Iranica*, Vol. 68, 4-16. Akademische Druck-u. Verlagsanstalt, Graz.

P. 92. Metabolism of (+)- and (-)-limonenes to respective carveols and perillyl alcohols by CYP2C9 and CYP2C19 in human liver microsomes

Mitsuo Miyazawa^a, Masaki Shindo^a, Tsutomu Shimada^b

miyazawa@apch.kindai.ac.jp

^a Department of Applied Chemistry, Faculty of Science and Engineering, Kinki University, Kowakae, Higashiosaka, Osaka 577-8502, Japan (M.M., M.S.)

^b Osaka Prefectural Institute of Public Health, Nakamichi 1-chome, Higashinari-ku, Osaka 537-0025, Japan (T.S.)

Limonene, a monocyclic monoterpene, is present in orange peel and other plants and has been shown to have chemopreventive activities. (+)- and (-)-limonene enantiomers were incubated with human liver microsomes and the oxidative metabolites thus formed were analysed using gas chromatography-mass spectrometry. Two kinds of metabolites, (+)- and (-)-*trans*-carveol (products by 6-hydroxylation) and (+)- and (-)-perillyl alcohol (products by 7-hydroxylation), were identified and the latter metabolites were found to be formed more extensively than the former ones with liver microsomes prepared from different human samples. Sulfaphenazole and flavoxamine and antibodies raised against purified liver cytochrome P450 (P450 or CYP) 2C9, that inhibit both CYP2C9- and 2C19-dependent activities, significantly inhibited microsomal oxidation of (+)- and (-)-limonene enantiomers. The limonene oxidation activities correlated well with contents of CYP2C9 and activities of tolbutamide methyl hydroxylation in liver microsomes of 62 human samples, while these activities did not correlate with contents of CYP2C19 and activities of S-mephenytoin 4-hydroxylation. Of 11 recombinant human P450 enzymes (expressed in *Trichoplusia ni* cells) tested, CYP2C8, 2C9, 2C18, 2C19, and CYP3A4 catalysed oxidation of (+)- and (-)-limonenes to respective carveols and perillyl alcohols. Interestingly, human CYP2B6 did not catalyse limonene oxidations, whereas rat CYP2B1 had high activities in catalysing limonene oxidations. These results suggest that both (+)- and (-)-limonene enantiomers are oxidised at 6- and 7-positions by CYP2C9 and CYP2C19 in human liver microsomes. CYP2C9 may be more important than CYP2C19 in catalysing limonene oxidation in human liver microsomes, since levels of the former protein are more abundant than CYP2C19 in these human samples. Species-related differences exist in the oxidation of limonenes in CYP2B subfamily in rats and humans.

P. 93. Volatile components from the roots of *Scrophulia ningpoensis* Hemsl

Mitsuo Miyazawa, Yoshiharu Okuno

miyazawa@apch.kindai.ac.jp

Department of Applied Chemistry, Faculty of Science and Engineering, Kinki University, Kowakae, Higasi-osaka, Osaka 577-8502, Japan

Scrophulariae radix (Genzin in Japanese) is the dry roots of *Scrophulia ningpoensis* HEMSL. (Scrophulariaceae) and is cultivated as a medicinal plant in China. The plants have been used for the treatment of fever, swelling, constipation, pharyngitis, neuritis and laryngitis in the Chinese traditional medicine. Various iridoid glycoside and iridoido-related aglycones were isolated from *S. ningpoensis* and identified.¹⁻³⁾ In our previous papers, we isolated four cinnamic acid derivatives as suppressive compounds of SOS-inducing activity, which is one of the DNA repair systems, from this plant⁴⁾. The volatile components of several Chinese crude drugs have been investigated in our research on the flavour compounds or flavour ingredients.⁵⁻⁸⁾ However so far, there is no report of the volatile components from *S. ningpoensis*, and odour has not yet been reported for this Chinese crude drug. In this report, the composition of volatile components from the radix of *S. ningpoensis* was investigated.

The composition of the volatile oil from the radix of *Scrophulia ningpoensis* has been investigated by capillary GC, GC-MS. The oil contained 103 volatile components of which 3.5 % were terpenoids. The main constituents were fatty acids, such as palmitic acid (25.4%), linoleic acid (10.0%), α -linolenic acid (6.1%), γ -linolenic acid (4.8%), *cis*-palmitoleic acid (3.9%), *trans*-oleic acid (3.5%), *cis*-oleic acid (2.7%), and *trans*-palmitoleic acid (1.2%).

1. Kitagawa I, Nishimura T, Takei M, Yoshioka I. *Chem. Pharm. Bull.* 1967; **15**: 1254-1256.
2. Kajimoto T, Hidaka M, Shoyama K, Nohara T. *Phytochemistry* 1989; **28**: 2701.
3. Qian J, Hunkler D, Rimpler H. *Phytochemistry* 1992; **31**: 905-911.
4. Miyazawa M, Okuno Y, Nakamura S, Kameoka H. *J. Agri. Food Chem.* 1998; **46**: 904-910.
5. Miyazawa M, Minamino Y, Kameoka H. *Flavor Fragr. J.* 1996; **11**: 57-60.
6. Miyazawa M, Minamino Y, Kameoka H. *Flavor Fragr. J.* 1997; **12**: 15-17.
7. Miyazawa M, Yamamoto Y, Kameoka H. *J. Essent. Oil Res.* 1992; **4**: 227-230.
8. Miyazawa M, Shimamura S, Kameoka H. *Natural Product Letters* 1997; **9**: 245-248.

P. 94. Chemical analysis and antifungal activity of the essential oil of *Calea clematidea* (Asteraceae)

Ademir F. Morel^{a*}, Adriana Flach^a, Ubiratan F. da Silva^a, Emilia C. S. Dessoay^a, Euclésio Simionatto^a, Carlos E. B. Linares^b, Sydney H. Alves^b

*afmorel@base.ufsm.br

^a Departamento de Química da Universidade Federal de Santa Maria (UFSM), Campus-Camobi, 97.105-900, Santa Maria, RS, Brazil


^b Departamento de Análises Clínicas e Toxicológicas da Universidade Federal de Santa Maria, Santa Maria, RS, Brazil

The genus *Calea* comprises ca100 species growing in the tropical and sub-tropical America, as herbs and shrubs. *Calea clematidea* Baker, Asteraceae, is a native shrub found in the southern region of Brazil and in Uruguay (Burkart, 1974). In Rio Grande do Sul, it is traditionally used by local people as anti-influenza agent and for the treatment of catarrh. In this work, we describe the study of the essential oils of the leaves and flowers of *Calea clematidea* from the state of Rio Grande do Sul, Brazil, a plant which has not been investigated previously.

The chemical composition of the essential oils of *Calea clematidea* Baker obtained by hydrodistillation of the leaves and flowers was analysed by GC and GC/MS and assayed for their antifungal activities. The essential oil of the leaf had a high content of a new natural epoxy-terpenoid, named clemateol (ca 70%) (**1**), with minor amounts of *o*-vanillin (6.5%), spathulenol (4.2%), α -terpinene (4.0%), germacrene-B (2.9%), yomogi alcohol (1.8%), (*E*)-caryophyllene (1.7%), *m*-cymene (1.6%), and α -gurjunene (1.5%), while the essential oil of the flowers was characterized by a higher content of thymol methyl ether (ca. 80%), with minor amounts of clemateol (4.8%) and *o*-cymene (4.7%). The antimicrobial activity of the oils was also evaluated against dermatophytes for their possible use in pharmaceutical preparations for topical applications. The oil of the leaves showed a moderate antifungal activity (MIC>3 mg/mL) against *Trichophyton tonsurans*, *Trichophyton rubrum*, *Trichophyton menthagrophytes* var. *interdigitale*, *Epidermophyton floccosum*, *Microsporium gypseum*, *Microsporium canis* and *Microsporium nanum*.

On alkaline hydrolysis, clemateol (**1**) afforded 2,5,5-trimethyl-2,3-epoxy-hept-6-en-4-ol (**2**), identified by co-TLC, IR, ¹H- and ¹³C-NMR. A Horeau determination of the absolute configuration indicated (*R*) configuration for the hydroxyl-bearing C-4 of compound **2**. To determine the absolute configuration of C-3 of **1**, the alcohol **2** was initially converted to the corresponding methyl ether **3** which was subsequently subjected to TCNE-catalysed alcoholysis in methanol to give **4** as a single stereoisomer (Masaki, 1993). The absolute configuration of C-3 of compound **4** was also determined as (*R*) by the horeau's method. Therefore, the stereochemistry of **1** was determined as (3*R*) and (4*R*).

In general, the essential oils have been reported to inhibit fungal growth. The results show that *Calea clematidea* essential oil obtained from the leaves, clemateol and compound **3** exhibited a moderate fungistatic and fungicidal property against dermatophytes. The oil of the flowers had no effect on the microorganisms tested.



1. Burkart A. (1974) Flora Ilustrada de Entre Rios. ISAC eds. Entre Rios, Argentina 394 - 397

2. Masaki Y, T Miura, M Ochiai. (1993) Synlett 847- 849

P. 95. Essential Oils of Two Marchantiales Liverworts: *Targionia lorbeeriana* and *Asterella blumeana*

Marta Neves ^a, Manuel Fernandes-Ferreira ^b, Rui Morais ^c

marta@esb.ucp.pt

^aExtensão da Escola Superior de Biotecnologia em Caldas da Rainha - Universidade Católica Portuguesa, Rua Mestre Mateus Fernandes 2500-237 Caldas da Rainha, Portugal

^bUniversidade do Minho, Departamento de Biologia, 4300 Braga, Portugal

^cEscola Superior de Biotecnologia - Universidade Católica Portuguesa, Rua Dr. António Bernardino de Almeida, 4200-072 Porto, Portugal

Liverworts are known to be a rich source of terpenoids and phenolic compounds, which constitute the oil bodies. From these plants, a high diversity of substances has been isolated, including new structural types, bioactive compounds and unusual fragrances (Asakawa 1995).

Since only a few species of liverworts have been chemically investigated (less than 5% of known species), these plants constitute a phytochemical treasure where many interesting substances remain to be isolated. However, advances on the study of the chemistry of liverworts are hindered by the difficulty in collecting a sufficient quantity of pure material of certain species. Their small size and the fact that bryophytes usually grow mixed with other species and strongly attached by their rhizoids to the soil, make their purification a difficult and a time consuming procedure (Asakawa, 1995).

In vitro cultures of liverworts can be a good alternative to obtain biomass for the phytochemical analyses, since several studies with different species, suggested that the ability of *in vitro* grown liverworts to produce secondary metabolites, is similar to those found in field grown plants (Becker, 1994).

Recently we reported bioactive sesquiterpene lactones from the wild and *in vitro* grown *Targionia lorbeeriana* (Neves *et al.* 1999 and 2001) and triterpenoids from *in vitro* cultures of *Asterella blumeana* (Neves *et al.* 1998).

In this communication we describe the research on essential oils of *Asterella blumeana* and *Targionia lorbeeriana* by GC and GC-MS analysis of liverworts hydrodistillates.

The composition of essential oils of wild and *in vitro* grown *Targionia lorbeeriana* was compared. In both cases the monoterpenyl ester *cis*-2-methylene-3-(1-methylethenyl)-cyclohexanylacetate constituted the major compound. A significant number of compounds produced in the open were maintained in *in vitro* cultures, although a considerable number of other ones were not detected. Instead, under *in vitro* conditions, some new compounds were found which do not accumulate under wild conditions.

Essential oils produced by *in vitro* grown gametophytes, *calli* and suspension cultures of *Asterella blumeana* were evaluated. Monoterpene hydrocarbons constituted the majority of essential oils of plantlets, followed by oxygenated monoterpenes and oxygenated alkanes. Undifferentiated cultures were found to be unable to produce volatile terpenoids.

Asakawa Y (1995) In. *Progress in the Chemistry of Organic Natural Products*. Vol. 65. W Herz, GW Kirby, RE Moore, W Steglich & Ch Tamm (Eds.). Springer - Verlag Wien New York.

Becker H (1994) *Journal of Hattori Botanical Laboratory* 76: 283-291.

Neves M, M Ferreira, C Terreaux, K Hostettmann, R Morais (2001) *Zeitschrift für Naturforschung*, 56c, 726-730.

Neves M, R Morais, S Gafner, H Stoeckli-Evans, K Hostettmann (1999) *Phytochemistry* 50, 967-972.

M Neves, R Morais, S Gafner, K Hostettman (1998) *Phytotherapy Research* 12, S21-S24.

P. 96. Comparative aroma compound analysis of different essential oils of *Lippia rugosa* from Cameroon using GC-FID, GC-MS and olfactometry

Martin Benoit Ngassoum^a, Leopold Tatsadjeu^a, Leopold Jirovetz^b, Gerhard Buchbauer^b, Manochehr Shahabi^b

ngassoum@caramail.com

^a Department of Applied Chemistry, ENSAI-IUT, University of Ngaoundere, BP 355, Ngaoundere, Cameroon

^b Institute of Pharmaceutical Chemistry, University of Vienna, Althanstrasse 14, A-1090 Vienna, Austria

Lippia rugosa A. Chev. (syn. *L. adoensis* F.TW.A., *L. nigeriensis* Moldenke) is a robust woody perennial plant of the *Verbenaceae* family, up to 12 feet high, with large oblong-lanceolate bluish-green leaves, pleasant aromatic flowers, small, whitish in branched inflorescence. The upper surface of the leaves are rough to the touch, venation prominently rugosa, reticulate and the plant is found in the Savannah area of the Adamaoua province (Cameroon). *Lippia multiflora*, another species of the about 200 different *Lippia* species is common in western Africa, differs from *L. rugosa* by the upper surface of the leaves, which is smooth to the touch and venation obscure. The plant, locally named "Gossolhi", is used for medicinal purposes against many diseases, such as indigestion, rheumatism, fever, cough and jaundice.

Five essential *L. rugosa* oils from Cameroon were obtained by steam distillation and evaluated by professional perfumers for their aroma as follows: fresh old leaves: sweet, floral; fresh young leaves: sweet, floral (strong geraniol note); fresh flowers: sweet, floral with herbal and spicy side-notes; dried leaves: fresh-fruity (strong limonene note), sweet, floral, weak woody and stems: sweet, floral, fresh-fruity side-notes.

The composition of the five essential oils of *Lippia rugosa* from Cameroon was investigated using GC-FID, GC-MS and olfactometry. As main compounds especially monoterpenes and sesquiterpenes were identified. Linalool, geraniol, geranial, limonene, myrcene, neral and linalool oxides as well as β -caryophyllene, *trans*- β -farnesene, germacrene-D, T-murolol and caryophyllene oxides were found in higher concentrations.

A correlation of aroma compounds responsible for the characteristic odor impressions of each sample is additionally given.

P. 97. Contribution to the *Hypericum* L. genus chemotaxonomy in the mainland Portugal

Teresa Nogueira^a, Fernanda Duarte^a, Regina Tavares^a, M. J. Marcelo Curto^a, Carlo Bicchi^b, Patrizia Rubiolo^b, Mário Lousã^c

teresa.nogueira@ineti.pt

^aInstituto Nacional de Engenharia e Tecnologia Industrial, Estrada do Paço do Lumiar, Edifício F, 1649-038 Lisbon, Portugal

^bDipartimento di Scienza e Tecnologia del Farmaco, Via Pietro Giuria 9, I-10125 Turin, Italy

^cDepartamento de Proteção de Plantas e Fitoecologia, Instituto Superior de Agronomia, Tapada da Ajuda, 1349-017 Lisbon

In recent years *Hypericum perforatum* L. (*Guttiferae*) has been studied worldwide. Pharmacological activities, including antidepressant, antiviral and antibacterial, provide supporting evidence for several of the established traditional uses reported (Barnes *et al.*, 2001). In Portugal three species of *Hypericum* L. genus (*H. androsaemum* L., *H. perforatum* L. and *H. undulatum* Schousb. ex. Willd.) have been used for a long time.

Following previous phytochemical analysis of this genus (Nogueira *et al.*, 1998, 1999), a comparative chemotaxonomical study of the *taxa* from the mainland Portugal is presented. This study relies on taxonomical characters, analysis of the aromas and chemical composition of essential oils of the following species: *H. androsaemum* L., *H. pulchrum* L., *H. montanum* L., *H. tomentosum* L., *H. pubescens* Boiss., *H. elodes* L., *H. perfoliatum* L., *H. linarifolium* Vahl., *H. humifusum* L., *H. undulatum* Schousb. ex. Willd., *H. perforatum* L., *H. calycinum* L. and *H. hircinum* subsp. *majus* (Aiton) N. Robson.

The analysis of the aromas was performed by olfactometry, using a sensor array detection device equipped with a multi-element array of 32 chemical sensors (Hodgins, 1997). The chemical characterization of essential oils was carried out by gas chromatography (GC) and GC coupled to mass spectrometry. The results were treated by multivariate analysis, classification and ordination techniques, namely cluster and principal components analysis.

Significant differences were found in the aromas and in the major components of *H. androsaemum* essential oils, by comparing to other *taxa*. *H. tomentosum* appears also as a separate group based alone on the essential oils characters. The group including all *taxa* but *H. androsaemum* and *H. tomentosum* reveals some compositional diversity of aromas and essential oils well correlated with morphological taxonomical characters.

Barnes J, LA Anderson, JD Phillipson (2001) *J. Pharm. Pharmacol.* 53: 583-600.

Hodgins D (1997) *Techniques for Analyzing Food Aroma* R Marsili Ed. Marcel Dekker New York 11: 331-371.

Nogueira T, F Duarte, F Venâncio, R Tavares, M Lousã, C Bicchi, P Rubiolo (1998) *Silva Lusitana* 6: 55-61.

Nogueira T, F Duarte, R Tavares, MJ Marcelo Curto, J Capelo, AC Freitas (1999) *Flavour and Fragrance Journal* 14: 195-199.

P. 98. Stereoselective formation of (1*R*, 2*S*, 4*R*)-(+)-*p*-menthane-2,8-diol from α -pinene by *Aspergillus niger*


Yoshiaki Noma^a, Mai Furusawa^a, Toshihiro Hashimoto^b, Yoshinori Asakawa^b

ynoma@tokushima.bunri-u.ac.jp

^aFaculty of Domestic Sciences, Tokushima Bunri University, Yamashiro-cho, Tokushima 770-8514 Japan

^bFaculty of Pharmaceutical Sciences, Tokushima Bunri University, Yamashiro-cho, Tokushima 770-8514 Japan

The biotransformation of pinenes has been investigated from view point of environmental sciences between microorganisms and terpenoids in cedar and pine forests in Japan and production of functional substances (Noma & Asakawa 1994, 1995, 1998, 2000). *Aspergillus niger* converted (-)- α -pinene (**1a**) to (1*R*, 2*S*, 4*R*)-(+)-*p*-menthane-2,8-diol (**6a**) via (-)-*trans*-sobrerol (**2a**), (-)-8-hydroxycarvotanacetone (**3a**) and (+)-8-hydroxycarvomenthone (**4a**), together with (-)-vervenol, (-)-verbenone and (-)- α -terpineol. On the other hand, (+)- α -pinene (**1b**) was biotransformed enantio- and diastereoselectively except for the formation of (-)-8-hydroxyisocavomenthone (**5b**).



The present paper deals with the enantio- and diastereoselectivity of (-)-, (+)- and racemic *trans*-sobrerols (**2a**, **2b** & **2ab**) and (-)-, (+)- and racemic *cis*-sobrerols, (-)-, (+)- and racemic 8-hydroxycarvotanacetones (**3a**, **3b** & **3ab**) and (-)-, (+)- and racemic 8-hydroxy carvomenthone (**4a**, **4b** & **4ab**) by *A. niger* in order to clarify the stereoselective formation of **6**. It was clarified that the stereoselective formation of **6** from **1** is due to the high substrate specificity of reductase towards the reduction of **4a**.

Noma Y, Asakawa Y (1994) In: *Biotechnology in agriculture and forestry. Medicinal and aromatic plants VII* (Bajaj YPS ed.) Vol. 28, p. 185-202. Springer Berlin, Heidelberg.

Noma Y, Asakawa Y (1995) In: *Biotechnology in agriculture and forestry. Medicinal and aromatic plants VIII* (Bajaj YPS ed.) Vol. 33, pp. 62-96. Springer Berlin, Heidelberg.

Noma Y, Asakawa Y (1998) In: *Biotechnology in agriculture and forestry. Medicinal and aromatic plants X* (Bajaj YPS ed.) Vol. 41, p. 194-237. Springer Berlin, Heidelberg.

Noma Y, Asakawa Y (2000) *Current Topics in Phytochemistry* Vol. 4, 63-78, Research Trends, Trivandrum, India

P. 99.**Aromatic and medicinal plants of Sudan**Amal M. Nour^a, Zeinab A. Mohammed^b, A. E. Abdalla^b, H. S. Khalid^b

amal_mukhtar@hotmail.com

^a Dept. of Pharmacognosy University of Khartoum P.O. Box 11496 Khartoum – Sudan^b Medicinal & Aromatic Plants Research Institute P. O. Box 2404 Khartoum – Sudan

Sudan is the largest country in Africa with a land area of 2.5 million squares Km. It has three different climatic zones namely northern deserts, semi-arid central region and the tropical swamp and rain forest region of the south. Therefore Sudan enjoys a vast, diverse flora consisting of 3173 species of flowering plants belonging to 170 families and 1280 genera.

Medicinal and Aromatic plants and their derivatives are on sale in special shops called "Atareen" but there is a very little cultivation of aromatic plants in Sudan. They are collected from plants that grow wildly and therefore supply is irregular.

The poster covers examples of essential oils-containing or fragrance-bearing species. It is presented in a form table indicating the plant family, plant name (Latin, English and Arabic), plant habit and habitat (photograph or line drawing for the species) part used and type of the essential oil, fragrance or flavouring agent present in this plant part.

The main objective of this poster is to give an idea of the potentiality of investing on aromatic plants and their cultivation and utilisation in Sudan.

Andrews, F. W. (1950, 1952 and 1956), The Flowering Plants of Aglo-Egyptian Sudan. (Three Volumes)

**P. 100. Investigation of essential oil content of crude drugs from lovage
(*Levisticum officinale* Koch.)**

I. Novák, G. Székely, Zs Pluhár

inovak@omega.kee.hu

Szent István University, Faculty of Horticultural Sciences, Department of Medicinal and Aromatic Plants, Budapest, Hungary,
P.O. Box 53., H-1518

The essential oil content of species of the *Apiaceae* family and among them the essential oil content of *Levisticum officinale* is influenced by different biological and technological factors (Bylaite et al., 1998; Stahl-Biskup and Witchmann, 1991; Szebeni and Galambosi, 1992; Székely et al., 2001). Our investigations were focused on determining the role of some factors on the essential oil content and composition of different crude drugs of lovage. These factors were the followings: plant organ, age of plant, phenophase and harvesting time.

The experimental populations were grown under open field conditions at the Experimental Station of the Department in Budapest. We investigated one-, two-, and three-year-old populations of *Levisticum officinale* KOCH. of unknown origin from our gene-bank collection in 2001. Average leaf samples were collected in ten days' interval from July till November in case of one- and two-year-old populations and root samples were taken from the end of August till the end of October in ten days' interval from each population. Seed samples were taken in the two-year-old population in five different phenophases (green fruit, milky fruit, waxy fruit, ripe fruit and over ripe fruit stage). Essential oil content of crude drugs was determined by hydrodistillation. The main components of the oil were analysed by capillary gas-chromatography using standards for identification.

Our results proved the significant effect of the age of the plantation and a difference in oil accumulation also between plant organs, which is in harmony with former findings (Bylaite et al., 1998; Szebeni and Galambosi, 1992). In case of root samples the highest essential oil content was found in the three-year-old population (0,5-0,7% d.w.), compared with the two- and one-year-old populations (0,2-0,5% d.w.; 0,1-0,3% d. w., respectively). In case of leaf samples the effect of the age was not significant (between 1,3-2,4%). The highest volatile oil contents were found in the leaves in the second decade of August. In case of the roots the maximum was reached in the end of October (one- and three-year-old stands) and in the end of August (two-year-old stands), which requires further investigations. As for the fruits, the highest volatile oil content was found at green stage (5,9% d. w.).

The effect of harvesting time, age of plantation and phenophase on the accumulation of the main essential oil components (phtalide component in root, α -terpinyl-acetate in leaves and β -phellandrene in the seeds) could not be proved.

Bylaite E, Venskutonis RP, Roozen JP (1998) Influence of Harvesting Time on the Composition of Volatile Components in Different Anatomical Parts of Lovage *J. Agr. Food Chem.* 46: 3735-3740.

Stahl-Biskup E, Witchmann EM (1991) Composition of the Essential Oils from Roots of some *Apiaceae* in Relation to the Development of their Oil Duct System *Flavour Fragr. J.* 6: 249-255.

Szebeni-Galambosi Zs, Galambosi B (1992) Growth, Yield and Essential Oil of Lovage Grown in Finland. *J. Ess. Oil Res.* 4: 375-380.

Székely G, Bernáth J and Zámboriné Németh É (2001) Floral-Biological Characteristics and Fruit Development of Fennel World Conference on Medicinal and Aromatic Plants, 8-11 July, Budapest, Hungary, Abstracts: 125.

P. 101. Comparison of fruit development features from fennel populations of various age

Gabriella Székely, J. Bernáth, I. Novák
gbszekely@hotmail.com

Department of Medicinal and Aromatic Plants, Faculty of Horticultural Sciences, Szent István University, Budapest, Hungary, P.O. Box 53. H-1518

Fennel (*Foeniculum vulgare*. Mill) is one of the important species of the Apiaceae, which is used in large amount in phytotherapy and food industry worldwide and in Hungary as well.

In this respect fennel cultivar (*Foeniculum vulgare* subsp. *capillaceum* var. *vulgare*) 'Soroksári' and a population of genebank origin ('F86') were analysed on individual level. In the course of 2001 fruit development of plant stands being in the first, second and third vegetation cycles was examined.

In order to characterise the fruit development processes, fruits from the 1st range umbels were taken and analysed, continuously. The sampling was conducted from the start of the fruit development (from the end of fertilisation) until the full ripening.

Essential oil content of the fruits was determined by hydrodistillation in Clevenger-apparatus according to the method of Ph. Hg. VII. Relative percentage of the oil constituents was calculated from the GC (Shimadzu GC-B14) peak areas in percent of the total area.

It is a characteristic phenomenon of the essential oil accumulation, that its quantity and quality in the fruits change in relation to the increase of total dry matter during the ontogenesis. The essential oil yield is higher at the beginning of fruit development, then decreases along with the formation of endosperm (Bernáth *et al.*, 1999, Gleisberg, 1960).

The essential oil content of fruits either collected in the first, or in the second vegetation cycles is the highest in green seed stage and it shows a decreasing tendency in case of waxy and ripe seed stages, in close agreement with the data published in the literature (Kattaá, 1996, Bernáth *et al.*, 1999). However, in case of investigated populations, the maximum oil content in the third vegetation years was measured in full ripe stage.

According to our investigation, the content of anethole and methyl chavicol shows an increasing tendency during the fruit development, while that of fenchone, limonene, α -pinene and β -pinene decreases.

Our results on the content of anethole, methyl chavicol are partially identical to those reported in the literature, mainly because the accumulation level of the components highly depends on the taxa.

Acknowledgements: The research was supported by Hung. Nat. Sci. Found (OTKA T-034289).

Bernáth J., Németh, É. Petheő F., Mihalik, E. Kálmán, K. Franke R. (1999): Regularities of the essential oil accumulation in developing fruits of fennel (*Foeniculum vulgare* Mill.) and its histological background, J. Essent. Oil Res. 11:431-438.

Kattaá A., 1996: Morphological and production-biological study of fennel (*Foeniculum vulgare* Mill.) and caraway (*Carum carvi* L.) populations of different origin, PhD. Dissertation, Univ. Hort. and Food Industry, Budapest, pp. 60.

Gleisberg, W., R. Hartrott, 1960: Die Reifungsvorgang in der Fenchelfrucht Garten- bauwissenschaft, 7: 245-218.

P. 102. Determination of the growing location of marjoram (*Origanum majorana* L.) samples by comparison of essential oil profiles

Johannes Novak^a, Friedrich Pank^b, Jan Langbehn^b, Wolf-Dieter Blüthner^c, Carla Vender^d, Léon Van Niekerk^e,
Wolfram Junghanns^f, Chlodwig Franz^a

Johannes.Novak@vu-wien.ac.at

^aInstitute for Applied Botany, University of Veterinary Medicine, Veterinaerplatz 1, A-1210 Vienna, Austria

^bInstitute of Horticultural Crops, Federal Centre for Breeding Research on Cultivated Plants, Neuer Weg 22/23, D-06484
Quedlinburg, Germany

^cN.L.Chrestensen Erfurter Samen- und Pflanzenzucht GmbH, Postfach 854, D-99092 Erfurt, Germany

^dISAFSA, Piazza Nicolini 6, I-3805 Villazzano di Trento, Italy

^eDarbonne, 6, bd. Joffre - B.P.8, F-91490 Milly-La-Forêt, France

^fMAWEA, Majoranweg 21, D-06449 Aschersleben, Germany

A canonical discriminant function analysis (CDA) was applied on essential oil compounds profiles to determine the growing location of marjoram samples.

Twelve out of 21 essential oil compounds were integrated into two discriminant functions as the best predictors for geographical differences. The stepwise integration allowed the determination of the essential oil compounds mainly responsible for the geographical variation, namely myrcene, terpinene-4-ol, γ -terpinene and *p*-cymene.

The two discriminant functions themselves showed a strong association between the locations and the predictors. The two functions allowed a correct classification of 95% of samples (crossvalidated).

The high positive classification rate demonstrates the CDA's ability as quality assurance tool enabling authenticity control.

P. 103. Volatile components of the New Zealand liverwort *Hymenophyton flabellatum* (Hymenophytaceae)

M. Toyota^a, I. Omatsu^a, Y. Asakawa^a, J. Braggins^b


phiku_usi@ph.bunri-u.ac.jp

^aFaculty of Pharmaceutical Sciences, Tokushima Bunri University, Yamashiro-cho 180, Tokushima 770-8514, Japan

^bSchool of Biological Sciences, The University of Auckland, Private Bag, 92019, New Zealand

The New Zealand liverwort *Hymenophyton flabellatum* (Hymenophytaceae) grows on shaded wet soil, humus and old logs in forest, usually in shade, and on banks beside streams. It contains 1-(2,4,6-trimethoxyphenyl)-but-2-en-1-one (**1**), responsible for the characteristic hot taste of this liverwort, together with β -caryophyllene (Asakawa *et al.* 2001). Further fractionation of the volatile components of the ether extract of *H. flabellatum* resulted in the isolation of five 1-(2,4,6-trisubstituted phenyl) but-1-one derivatives (**2-6**) and three (**7-9**) (Toyota *et al.* 2002). The structures of the newly isolated compounds were elucidated by ¹H and ¹³C NMR spectral analysis.

Compound **1** has been found in the Japanese fern, *Arachinoides standishii* (Tanaka *et al.* 1980), and compound **1** and **5** in the aquatic Angiosperm, *Dysophyla verticillata* (Chakrabarti & Chakraborty 1988). As far as we are aware this is the first record of the isolation of phenylbutenones and chromones in liverworts.



Asakawa Y, Toyota M, Oiso Y, J Braggins (2001) *Chem. Pharm. Bull.*, **49**, 1380-1381.

Chakrabarti A, Chakraborty DP (1988) *Phytochemistry* **27**, 3683-3684.

Tanaka N, Maehashi H, Saaito A, Murakami T, Saiki Y, Chen C-M, Iitaka Y (1980) *Chem. Pharm. Bull.* **28**, 3970-3077.

Toyota M, Omatsu I, Asakawa Y (2002) *Chem. Pharm. Bull.* (to be submitted).

P. 104. The essential oil of *Cyclotrichium longiflorum* Leblebici

T. Özek^a, B. Demirci^a, K. H. C. Başer^b, Z. Aytaç^c

tozek@anadolu.edu.tr

^aMedicinal and Aromatic Plant and Drug Research Centre (TBAM), Anadolu University, 26470 Eskişehir, Turkey

^bFaculty of Science and Letters, Gazi University, 06500 Ankara, Turkey

The genus *Cyclotrichium* (Boiss.) Manden. et Scheng. is represented in Turkey by six species (Davis 1982, 1988). *Cyclotrichium longiflorum* Leblebici was described in the 1st supplement of the Flora of Turkey as a new record for Turkey (Davis 1988). The plant material was collected from South Eastern Turkey.

Water-distilled essential oil from the air-dried aerial parts of *Cyclotrichium longiflorum* Leblebici was analysed by GC and GC/MS. Isopinocampone (34.8%), β -pinene (17.3%) and pinocampone (8.4%) were found as major constituents.

Davis PH, *Flora of Turkey and the East Aegean Islands*. Vol. 7, p. 346-9, University Press, Edinburgh (1982).

Davis PH, *Flora of Turkey and the East Aegean Islands*. Vol. 10, p. 208, University Press, Edinburgh (1988).

P. 105. Seasonal variation in the chemical composition of *Cistus albidus* L. from Spain

Jesús Palá-Paúl^a, M^a José Pérez-Alonso^a, Arturo Velasco-Negueruela^a, J. Sanz^b

Quibey@bio.ucm.es

^a Dpto. Biología Vegetal I (Botánica), Facultad de Biología, Universidad Complutense de Madrid, 28040-Madrid, Spain

^b Instituto de Química Orgánica, Juan de la Cierva nº 3, 28006-Madrid, Spain.


The genus *Cistus* L., belongs to the Cistaceae family and, it comprises 12 species in Spain. This genus is widely distributed in Mediterranean area and their species are considered as pioneers in disturbed environment. *Cistus albidus* L., is an evergreen woody shrub up to 1.5 meters. It grows in calcareous and siliceous soils and it is well represented all throughout Spain although it is more common in the mid South (1).

The terpene content and emission of *Cistus albidus* have been previously studied under field conditions (2), but as far as we know this is the first report about the seasonal variation on the chemical composition of the oil of this species extracted by hydrodistillation and analysed by Gas-Chromatography (GC) and GC-Mass spectrometry.

We have analysed the chemical composition of two samples of the same population of *Cistus albidus* gathered in Spain, one on winter and the other one on spring season. The samples were divided in leaves, stems, seeds and flowers according to the season of the year. The seed oils of this species were the part of the plant with less yield and constituents. All the samples were richer in sesquiterpenes than in monoterpenes. We have found quantitative but not qualitative differences.

The principal compounds of the winter leaves and stems oils were identified as α -zingiberene (14.8%), aromadendrene (10.5%), *ar*-curcumene (10.5%) and guaiol (6.5%) while the essential oil from the spring leaves and stems showed as main constituents aromadendrene (9.2%), *ar*-curcumene (8.8%), α -zingiberene (8.0%), guaiol (7.0%), (*E*)-caryophyllene (2.4%) and α -*trans*-bergamotene (1.8%). Finally the flower oils also contained these components but *epi*- α -muurolol (8.4%) and γ -selinene (5.4%) were identified as the principal ones.

Main constituents



1.- Castroviejo, S.; Aedo, C.; Cirujano, S.; Laínz, M.; Mntserrat, P.; Morales, R.; Muñoz Garmendia, F.; Navarro, C.; Paiva, J.; Soriano C. 1993. *Flora Ibérica, plantas vasculares de la Península Ibérica e Islas Baelares*. Real Jardín Botánico, C.S.I.C. Madrid.

2.- Llusia, J. and Puñuelas J. 2000. *Seasonal patterns of terpene content and emission from seven Mediterranean woody species in field conditions*. American Journal of Botany. 87, 133-140.

P. 106. Essential oil composition of the seasonal heterophyllous leaves of *Eryngium vesiculosum* Labill from Australia

Jesús Palá-Paúl^a, Joseph J. Brophy^b, Lachlan M. Copeland^c, M^a José Pérez-Alonso^a, Arturo Velasco-Negueruela^a

Quibey@bio.ucm.es


^a Dpto. Biología Vegetal I (Botánica), Facultad de Biología, Universidad Complutense de Madrid, 28040-Madrid, Spain

^b School of Chemistry, The University of New South Wales, Sydney NSW 2052, Australia

^c Botany, School of Rural Science & Natural Resources, University of New England, Armidale NSW 2351, Australia

Eryngium vesiculosum Labill., commonly named prostrate blue devil or prickfoot, is a perennial herb of coastal sands, lake margins and river beds in Australia and New Zealand (1). It has different leaves according to the seasons of the year. Summer leaves are laminoid and prickly while winter ones are linear entire and fascicular (2). Two samples of each Australian season of the same population have been gathered from the Northern Tablelands of New South Wales, Australia. The essential oil of the flowers and both kinds of leaves have been analysed by GC and GC-MS. We have found quantitative but not qualitative differences. The principal compounds of the winter leaves were identified as β -caryophyllene (20.3%), germacrene-D (19.2%) and α -humulene (8.8%) while the summer ones showed bicyclogermacrene (22.2%), β -caryophyllene (15.6%), germacrene-D (15.8%) and α -humulene (8.1%) as major constituents. Bicyclogermacrene has changed its minor percentage composition in winter leaves oil (2.4%) to a principal compound in the other ones (22.2%). The flower oil contained practically the same composition of the summer leaves. Although we have previously reported the essential oils of some *Eryngium* species (3), this is the first report about the chemical composition of the flower and the different sort of leaves of *E. vesiculosum*.

Main constituents



- 1.- Harden G. J. (1992) *Flora of New South Wales*, Royal Botanic Gardens Sydney. Volume 3. NSWU Press, Sydney.
- 2.- Webb C. J. (1984) *Heterophylly in Eryngium vesiculosum (Umbelliferae)*, New Zealand Journal of Botany, 22: 29-33.
- 3.- Brophy J. J., Goldsack R. J., Copeland L. M. and Palá-Paúl J (2002) *Essential Oil of Eryngium L. species from New South Wales (Australia)*, Journal of Essential Oil Research 2002 (in press).

P. 107. *Agastache foeniculum* (Pursh) O. Ktze. essential oil composition and its antibacterial activity

Anamarija Partl^a, Nikola Blažević^b, Božidar Stilinović^c, Dragomir Brkić^a
^apartl@croatia.botanic.hr

^a Zavod za farmaceutsku botaniku FBFa, Schrottova 39, HR-10000 Zagreb, Hrvatska

^b IREKS Aroma, Radnička cesta 37, HR-10000 Zagreb, Hrvatska

^c Botanički zavod PMFa, Rooseveltov trg 6, HR-10000 Zagreb, Hrvatska

Agastache foeniculum (Pursh) O. Ktze. is a perennial herb native to North America which is also cultivated as a decorative plant or used as a spice due to its anise-like aroma. Leaves are opposite, almost glabrous, whitish underneath, flowers purplish in terminal spike-like inflorescence.

It is closely related to east Asian (China, Japan, Korea) species *A. rugosa* (Fisch et Mey.) O. Ktze. which is one of the main medical herbs in Chinese traditional medicine and was scientifically surveyed as such, for example as antimycotic and anti-HIV drug (Blaszczyk *et al.*, 2000; Kim *et al.*, 1999, 2001).

The aim of this study is to determine the main essential oil components of *A. foeniculum* grown in Croatia and to test the antibacterial activity of the essential oil. Test bacteria were: *Bacillus subtilis*, *Staphylococcus aureus*, *Klebsiella* sp., *Serratia marcescens*, *Escherichia coli*, *Citrobacter* sp. and *Proteus mirabilis*. The plates were incubated 24 h at 37°C except for *S. marcescens*, which was incubated at room temperature. Blind control was used to monitor normal bacterial growth under the same conditions.

The oil was obtained from dried plant material by hidrodistillation using a modified Clevenger apparatus for 4 hours. The analysis of the essential oil by GC and coupled GC-MS showed the main components to be estragole (92.5%) and limonene (4.2%) while all other components were present in less than 0.5%. In some former studies (Fuentes-Granados *et al.*, 2000) myrcene was found as the third main component, present in our sample only in 0.05%. Total oil yield of the flowering plant was somewhat less than 2% of the dry plant weight.

Preliminary antibacterial study, performed using the technique of antibiograms, showed strong activity against *B. subtilis*, some activity against *S. aureus* and *Klebsiella* sp., discoloration and growth inhibition of *S. marcescens* and strong growth inhibition, without specific zones of inhibition, against *E. coli*, *Citrobacter* sp. and *P. mirabilis*. From these results, it can be concluded that the volatile components strongly inhibit the growth of different bacteria, rather by completely killing the bacteria due to the diffusion through the substrate. Further studies on other bacteria, but also some pathogenic fungi, would be interesting.

Blaszczyk T, J Krzyzanowska, E Lamer-Zarawska (2000) *Phytotherapy Res.* 14 (3): 210-212.

Fuentes-Granados RG, MP Widrlechner, LA Wilson (2000) *J. Essent. Oil Res.* 12 (5): 581-594.

Kim HK, HK Lee, CG Shin, H Huh (1999) *Arch. Pharm. Res.* 22 (5): 520-523.

Kim MH, WT Chung, YK Kim, JH Lee, HY Lee, B Hwang, YS Park, SJ Hwang, JH Kim (2001) *J. Essent. Oil Res.* 13 (3): 214-218.

P. 108. Comparison of essential oil composition and antibacterial activity between some Croatian wild growing plants and commercially available oils

Anamarija Partl^a, Marijana Brekalo^a, Nikola Blažević^b, Božidar Stilinović^c

apartl@croatica.botanic.hr

^a Zavod za farmaceutsku botaniku FBFa, Schrottova 39, HR-10000 Zagreb, Hrvatska.

^b IREKS Aroma, Radnička cesta 37, HR-10000 Zagreb, Hrvatska

^c Botanički zavod PMFa, Rooseveltov trg 6, HR-10000 Zagreb, Hrvatska

Three common aromatic species growing wild and commonly planted in Croatia - Dalmatian sage, rosemary and lavender - were chosen to compare their essential oil yield and composition against some commercially available oils of the same or related species.

Wild growing plants were collected in May/June on the south Croatian island Korčula; lavender and sage were collected during flowering period. Lavender was determined as *Lavandula latifolia* Med. which grows wild in Croatia but is very rare. Rosemary was determined as *Rosmarinus officinalis* L., which grows wild on some south Croatian islands in the typical rosemary garigue. Previous literature data state that although the same species, rosemary can vary significantly in the oil composition (Bego 1995; Schnaubelt, 1998, 1999). Dalmatian sage, *Salvia officinalis* L., is native to east Adriatic coast where it grows on rocky grasslands, but was spread throughout the whole Mediterranean. Oils were obtained by hidrodistillation using a modified Clevenger apparatus.

Two sets of commercially available oils were used for comparison of the essential oil composition - one set of oils which stated they are produced from Adriatic plants and another set of imported oils. The oils were analysed by GC to screen main components.

Antibacterial activity of the oils from wild growing plants was tested using the imported oils as a comparison and control for monitoring normal bacterial growth. The test bacteria were *Escherichia coli* and *Staphylococcus aureus*, and all the plates were incubated for 24 h at 37°C.

Wild growing plants showed the following oil characteristics: lavender yielded 5.5% essential oil from dried flowers and the main components were linalool (30.8%), borneol (18.4%) and 1,8-cineole (17.0%), while the total esters were only 6.7%, from which 5.6% was linalyl-acetate; rosemary had an oil yield of 2.7%, having as main components 1,8-cineole (34.8%), borneol (13.8%) and α -pinene (12.3%); and sage had an oil yield of 2.0%, with thujones (34.5%), 1,8-cineole (20.1%) and camphor (11.3%) as main components. Commercial samples of lavender oil showed the following results: imported lavender, declared as *L. angustifolia*, had very low linalyl-acetate (less than 6.0%) and the major components were 1,8-cineole (16.0%), linalool (40.0%), borneol (10.0%) and camphor (4.0%), while the Adriatic lavender (lavandine) had 8.0% linalyl-acetate, with major components being linalool (47%), 1,8-cineole (8.5%), borneol (8.5%), terpinen-4-ol (7.5%) and only 2.0% camphor. Adriatic sage oil was constituted of 44.0% thujones, 14.0% camphor and 11.0% 1,8-cineole, and imported sage of 29.0% thujones, 20.5% camphor and 14.5% 1,8-cineole. Adriatic rosemary oil showed 46.0% 1,8-cineole, 12.0% α -pinene, 6.5% borneol and 7.0% camphor, while the imported oil had 46.0% 1,8-cineole, 12.0% α -pinene, 6.0% borneol and 6.8% camphor, which are practically the same results.

Antibacterial tests showed higher activity for imported rosemary oil on *E. coli*, but lower on *S. aureus*; almost the same activity against both bacteria by lavender and the strongest bactericidal activity of both sage oils, especially against *S. aureus*.

This study reveals the necessity of perpetual testing of each oil batch against target bacteria.

Bego GV (1995) *What's essential about essential oils*. Lakin Publ., Lausanne, Switzerland.

Schnaubelt K (1998) *Advanced Aromatherapy*. Healnih Art Press Rochester, Vermont.

Schnaubelt K (1999) *Medical Aromatherapy*. Frog Ltd. Berkeley, California.

P. 109. Composition of the essential oil of *Crithmum maritimum* L. grown in Turkey

Musa Özcan ^a, Luis G. Pedro ^b, A. Cristina Figueiredo ^b, José G. Barroso ^b

luis.pedro@fc.ul.pt

^a Department of Food Engineering, Faculty of Agriculture, Selçuk University, 42031 Konya, Turkey.

^b Centro de Biotecnologia Vegetal, Dept. Biologia Vegetal, FCL, C2, Campo Grande, 1749-016 Lisbon, Portugal

Crithmum maritimum L. (Apiaceae), the only species of the genus, can be found along the coast of Europe, Africa, Asia and Australia. In Turkey fresh leaves and young branches are used as a condiment and pickle and, in popular medicine, as an appetizer and diuretic (Baytop, 1984; Özcan, 2000). According to Guenther (1950), early studies on *C. maritimum* oil showed that all parts of the plant contain volatiles, but the yield and the chemical composition of the oils varied considerably depending on the site of growth and soil conditions. A study of the seasonal variation in the composition of the essential oil of *C. maritimum* showed major fluctuations in the relative amounts of several components, mainly sabinene (7-42%) and γ -terpinene (26-55%) (Barroso *et al.*, 1992). Other reports on essential oils from *C. maritimum*, collected in Italy and Portugal, showed also different compositions (Flamini *et al.*, 1999; Pateira *et al.*, 1999).

The essential oil isolated from plants collected at Bolu-Abant, in Turkey, showed sabinene (27%), limonene (24%), γ -terpinene (19%) and terpinene-4-ol (9%) as the main components (Baser *et al.*, 2000). Senatore *et al.* (2000) reported on the composition of the oils from plants collected at Antalya and at Mersin, Turkey. The former showed β -phellandrene (30%), methylthymol (25%), *cis*- β -ocimene (14%) and *p*-cymene (13%) to be the major components while the latter was characterised by γ -terpinene (24%), dillapiole (21%), β -phellandrene (14%) and sabinene (12%). In another study, Özcan *et al.* (2001) found sabinene, γ -terpinene, methylthymol, terpinen-4-ol, dillapiole and *p*-cymene as major components.

The young leaves and the branches of *C. maritimum* were collected, during the first week of June 2001, at late vegetative phase, in icel (Sipahili and Yesilovacik), at sea level. The plant material was submitted to hydrodistillation using a Clevenger-type apparatus and the oil yields were 0.20% for Sipahili (S) and 0.23% for Yesilovacik (Y). The monoterpene hydrocarbon fraction was dominant in both oils, attaining 90% (S) and 80% (Y) of the total oils. The main components were γ -terpinene (S- 36%; Y- 88%), β -phellandrene (S- 21%; Y- 22%), sabinene (S- 13%; Y- 9%) and *p*-cymene (8% in both oils). The oxygen-containing monoterpenes attained 10 and 9%, respectively, and were dominated by methylthymol (S- 8%; Y- 9%). The only sesquiterpene detected in both oils was germacrene D (S- 0.2%; Y- 0.3). Dillapiole, the only component in the phenylpropanoid fraction, was detected in trace amounts in the oil from Sipahili but attained 10% in the oil from Yesilovacik.

Barroso JG, LG Pedro, AC Figueiredo, MSS Pais, JJC Scheffer (1992) *Flavour Fragr. J.* 7: 147-150.

Baser KHC, T Özek, B Demirci, Y Saritas (2000) *J. Essent. Oil Res.* 12: 424-426.

Baytop T (1984) *Treatment with plants in Turkey*, Publ. n° 3255, Istanbul University, Istanbul, Turkey.

Flamini G, E Mastrorilli, PL Cioni, I Morelli, L Panizzi (1999) *J. Essent. Oil Res.* 11: 788-792.

Guenther E (1950) *The Essential Oils*, Vol. IV. D. van Nostrand Company, New York.

Özcan M (2000) *Z. Eur. Food Res. Technol.* 210: 424-426.

Özcan M, A Akgul, KHC Başer, T Özek, N Tabanca (2001) *Nahrung/Food* 45: 353-356.

Pateira L, T Nogueira, A Antunes, F Venâncio, R Tavares, J Capelo (1999) *Flavour Fragr. J.* 14: 333-343.

Senatore F, F Napolitano, M Özcan (2000) *Flavour Fragr. J.* 15: 186-189.

P. 110. Essential oils of *Origanum vulgare* subsp. *hirtum* growing wild in Turkey

Musa Özcan ^a, Luis G. Pedro ^b, A. Cristina Figueiredo ^b, José G. Barroso ^b

luis.pedro@fc.ul.pt

^a Department of Food Engineering, Faculty of Agriculture, Selçuk University, 42031 Konya, Turkey.

^b Centro de Biotecnologia Vegetal, Dept. Biologia Vegetal, FCL, C2, Campo Grande, 1749-016 Lisbon, Portugal

Wild marjoram (*Origanum vulgare* L.) is a perennial bushy herb, which grows wild in Western and Southern Turkey and in other Mediterranean countries (Davis, 1982).

Origanum species have been used in medicine and as spice and condiment since antiquity, mainly because of their essential oils, which contain considerable amounts of carvacrol and thymol. The volatile aromatic compounds are employed in the food industry as a flavoring. The oil is used in perfumery for its spicy herbaceous notes. *O. vulgare* is well known as a plant with a medicinal value and as such is accepted officially in a number of countries.

The aerial parts of *O. vulgare* subsp. *hirtum* were collected at two different locations, Büyükeceli (İçel; sea level) and Delikkaya (İçel; 400 m). The essential oils, isolated by hidrodistillation from the air-dried aerial parts, using a Clevenger-type apparatus, were obtained in yields of 3.0% and 4.7% (v/w), respectively. GC and GC/MS analyses lead to the identification of twenty-eight components, accounting for more than 99% of the total oils. Monoterpenes constitute the main fraction, amounting to 98% in both oils, with carvacrol (51% and 63%), linalool (28% and 2%), γ -terpinene (6% and 10%) and *p*-cymene (5% and 9%) as the main components, respectively.

As reported previously (Fleischer and Sneer, 1982; Sezik *et al.*, 1993; Başer *et al.*, 1994; Özcan and Chalchat, 2002) carvacrol (up to 79%), *p*-cymene (20%) and γ -terpinene (up to 16%) are the main components of *O. vulgare* subsp. *hirtum* oil.

Our results fit into the above-mentioned reported data, except for the content of linalool (28%) in the oil isolated from the plants collected at Büyükeceli.

Başer KHC, T Özek, M Kürkcüoğlu, G Tümen (1994) *J. Essent. Oil Res.* 6: 31-36.

Davis PH (1982) *Flora of Turkey and the East Aegean Islands*. Vol.7, Univ. Press, Edinburgh.

Fleischer A, N Sneer (1982) *J. Food Agric.* 33: 441-446.

Özcan M, J C Chalchat (2002) *J. Spices Aromatic Crops* (in press).

Sezik E, G Tümen, N Kırırmer, T Özek, KHC Başer (1993) *J. Essent. Oil Res.* 5: 425-431.

P. 111. Essential oils from *Pterospartum tridentatum*

Ana L. Pereira^a, Generosa Teixeira^b, Pedro A. G. Santos^c, A. Cristina Figueiredo^c, José G. Barroso^c

ana.pereira@fc.ul.pt

^a Centro de Biologia Ambiental, Dep. de Biologia Vegetal, FCL, C2-Piso 1, Campo Grande, 1749-016 Lisboa, Portugal

^b Centro de Biologia Ambiental, Faculdade de Farmácia da Universidade de Lisboa, Av Forças Armadas 1649-019 Lisboa, Portugal

^c Centro de Biotecnologia Vegetal, Dep. de Biologia Vegetal, FCL, C2, Campo Grande, 1749-016 Lisboa, Portugal

Pterospartum tridentatum WK. & Lge is a small genus of the *Papilionoideae* subfamily included in the *Cytiseae* tribe, subtribe *Genistinae* (Talavera *et al.* 1999). It is a shrub up to 100cm, with yellow flowers, alternate branches and coriaceous winged stems, which shows a characteristic anatomy and interesting epidermal features (Teixeira and Pereira 2001).

This plant is widely distributed in central west Iberian Peninsula and it is used in popular medicine in Portugal, namely for the control of *diabetes mellitus* type 2. Phytochemical studies performed on the aqueous extract of the flowered stem apices of *P. tridentatum* led to the identification of flavonoids (Vitor and Paulo 2001), which revealed some *in vitro* antioxidant activity (Vitor, pers. comm.).

The essential oils of *P. tridentatum* were isolated by hydrodistillation, from aerial parts of the plant collected during the flowering phase, to determine the oil yield, and by distillation-extraction to determine the constituents percentage composition. The analyses of the essential oils were performed by GC and GC-MS.

An yellowish oil was obtained from *P. tridentatum* in a yield of <0.05%. *cis*-Theaspirane (13%), *trans*-theaspirane (12%), octen-3-ol (11%), geraniol (5%), *n*-heptanal (5%), *n*-nonanal (5%), *cis*-rose oxide (4%) and linalool (4%) were the most representative components of the oil.

Talavera S (1999) *Flora Iberica* vol. VII(I): 578.

Teixeira G, A Pereira (2001) *Biology of the Cell* 93: 338.

Vitor RF, A Paulo (2001) *Abstracts of Int. Symp. Phytochem. Soc. Europe* P049.

P. 112. Investigations on drug production and essential oil properties of thyme (*Thymus vulgaris* L.) in different growing areas of Hungary

Zsuzsanna Pluhár, Eszter Dienes

zpluhar@omega.kee.hu

Department of Medicinal and Aromatic Plants, Faculty of Horticultural Sciences, Szent István University, Budapest, Hungary,
P.O. Box 53, H-1518

Three populations of thyme (*Thymus vulgaris* L.), located in two regions of Hungary (in Budapest and in Transdanubia) were studied in detail regarding morphological, production biological and chemical properties during two vegetation periods (2000-2001).

The aim of our studies was to find individuals possessing high dry herbal mass and/or high level of essential oil yield including increased total amount of phenolic compounds (thymol + carvacrol), which may serve as a basis for developing new, highly productive Hungarian varieties in the future.

In the eastern Transdanubian region a younger and an older population were grown, while in Budapest we examined a population of the same age as the older Transdanubian population, parallelly. Twenty plants per each population were investigated individually. Each population existed on loose sandy soils differing mainly in the proportion of calcium carbonate (being the lowest in Budapest) and in macro element content (the young Transdanubian population was the best supplied).

Regarding morphological features, plant height changed by age, by harvesting period and by growing site. Higher values could be measured in Budapest, at first harvest and in younger population. Plant size diameter was affected only by growing site: Budapest appeared to be the better area in this respect.

The variability regarding individual dry mass production was considerable influenced by growing region, by age of the plants and by cutting period. We established that at the time of the first harvest the yields were about two-fold if compared with the second cutting.

Based on the Hungarian and international standards we have been found that most of the individuals examined in Budapest fulfil the required minimal essential oil content of 1.2 ml/100g. Moreover, our selection purpose, regarding 2 ml/100g essential oil level in average, could be reached or exceeded by some plants grown in Budapest, either. We detected the most constant essential oil composition in this site as well, where the total amount of phenolic compounds and that of the thymol were highly outstanding. The proportion of thymol and carvacrol changed considerably in the Transdanubian growing region not only individually, but also by cutting period. The essential oil yield was mainly affected by the dry mass.

According to high productivity and high accumulation level of essential oil, containing appropriate content of phenolic monoterpenes, several individuals has been selected in Budapest for further evaluation.

P. 113. In vitro evaluation of the antioxidant effect of the volatile compounds from *Satureja montana* L.

Mladen Milos, Ani Radonić

radonic@ktf-split.hr

Faculty of Chemical Technology, University of Split, Teslina 10, 21000 Split, Croatia

Widely used artificial antioxidants, such as butylated hydroxytoluene and butylated hydroxyanisole are very effective in their role as antioxidants (Madsen *et al.* 1995). However, their use in food products has been failing off due to their instability, as well as due to a suspected action as promoters of carcinogenesis (Namiki 1990, Pokorny 1991). For this reason, there is a growing interest in the studies of natural additives as potential antioxidants. The antioxidant properties of many aromatic herbs are reported to be effective in this role mainly due to the presence of hydroxyl groups in their phenolic compounds (Herrmann *et al.* 1981, Brraco *et al.* 1981, Kramer, 1985).

Savory (*Satureja montana* L.) is an aromatic plant growing wild in the Mediterranean region of Croatia. With regard to the presence of phenolic compounds in its essential oil, savory is known to possess some biological, especially antibacterial, activity. As a part of an investigation of natural antioxidants from Dalmatian aromatic plants, in this paper we report the detailed study of chemical composition and antioxidant activity of the savory volatile compounds in order to find the fraction or component with the highest protection against lipid oxidation. The analyses of the total essential oil, as well as of its fractions, were made by capillary gas chromatography coupled to mass spectrometry (Hewlett-Packard GC/MS system). The evaluation of antioxidant power was performed *in vitro* by the β -carotene bleaching method (Pratt, 1980).

The antioxidant activity of volatile compounds from *Satureja montana* L. was compared with that of pure compounds. Antioxidant power decreased in the order BHT > α -tocopherol > CHO fraction > total essential oil > phenolic fraction > thymol > carvacrol > CH fraction. The CHO fraction was more potent than total essential oil and other fractions, but less effective than synthetic reference standard BHT and comparable in activity with natural antioxidant α -tocopherol. The fact that CHO fraction was more effective as antioxidant than phenolic fraction or its pure constituents thymol and carvacrol suggests that synergy among minor oxygen containing compounds plays a crucial role for antioxidant power of the savory essential oil. Also, the concentration of the oil influenced its antioxidant power, too. The control sample without addition of antioxidant oxidised most rapidly and descending bleaching rates were demonstrated for the increased concentration.

Brraco U, J Loliger, J Viret (1981) *Journal of American Oil Chemical Society* 58: 686-690.

Herrmann K, M Schutte, H Muller (1981) *Deutsch Lebensm-Rdsch* 77: 134-138.

Kramer RE (1985) *Journal of American Oil Chemical Society* 62: 111-113.

Madsen HL, G Bertelsen (1995) *Trends in Food Science & Technology* 6: 271-277.

Namiki M (1990) *Critical Reviews in Food Science & Nutrition* 29: 273-300.

Pokorny J (1991) *Trends in Food Science & Technology* 9: 223-227.

Pratt DE (1980) *Autoxidation in food and biological systems*, Plenum Press, New York, 283-292.

P. 114. Marquesas Islands sandalwood concrete and biodiversity conservation of a forest species

Jean-François Butaud^a, Phila Raharivelomanana^b, Jean-Pierre Bianchini^b, Vincent Baron^c

raharive@upf.pf

^aDépartement Forêt et Gestion de l'Espace Rural, Service du Développement Rural, BP 100, 98713 Papeete, Tahiti, French Polynesia

^bLaboratoire de Chimie Analytique Appliquée, Université de la Polynésie Française, BP 6570, 98702 Faaa, Tahiti, French Polynesia

^cCentre de Coopération Internationale en Recherche Agronomique pour le Développement, BP 467, 98713 Papeete, Tahiti, French Polynesia

Santalum insulare (Bertero ex A. DC.) is an indigene sandalwood species from Marquesas Islands where 2 varieties have been distinguished by taxonomists: *S. insulare* var. *marchionense* and var. *deckeri* (Fosberg & Sachet 1985).

During the last 2 centuries, sandalwood has been heavily logged at the point it has disappeared from some islands and became quite rare on the others (Denning 1980).

Recently, a project to safeguard and multiply this natural resource in the field of forestry was initiated by the cooperation of the Territory of French Polynesia (Rural Development Service, SDR) and the International Cooperation Center in Agronomic Research for the Development (CIRAD). Concurrently, chemical analysis of wood samples gathered from 4 islands of marquesan archipelago have been done at the chemistry laboratory of the French Polynesia University (UPF).

Concrete from heartwood of 45 trees spared on 4 islands from Marquesas archipelago were analyzed by GC/MS method in order to estimate the quality of marquesan sandalwood and to put in relief eventual composition variability amongst sandal from different populations or islands.

Thus, multivariate analysis on the composition of the concrete volatile part put in evidence notable chemotype differences, depending on the geographical distance between islands but also inside the same island. Concentrations of main compounds (Alpha *et al.* 1996, 1997) were investigated and can be illustrated by variations of α - and β -santalol proportion between 15 and 60% and these of (*Z*)-nuciferol from 1 to 17% of the concrete.

So, before further genetic research, SDR initiated a program to develop sandalwood nurseries and plantations in order to conserve specificity of each sandalwood provenance. Following this rule, separated plantations from different chemotypes are done on each island.

Moreover, on islands where the sandalwood resource is very scarce, it might not be possible to multiply and replant local provenance. In this case, sandalwood from the closest chemotype will be provided to the island inhabitants.

Alpha T, Raharivelomanana P, Bianchini JP, Faure R, Cambon A (1997) Bisabolane sesquiterpenoids from *Santalum austrocaledonicum*. *Phytochemistry* **44** (8): 1519.

Alpha T, Raharivelomanana P, Bianchini JP, Faure R, Cambon A, Joncheray L (1996) α -santaldiol and β -santaldiol, two santalane sesquiterpenes from *Santalum insulare*. *Phytochemistry* **41** (3): 829.

Denning G (1980) Islands and Beaches. Discourse on a silent land: Marquesas 1774-1880. University Press of Hawaii. Honolulu.

Fosberg FR, Sachet MH (1985) *Santalum* in Eastern Polynesia. *Candollea* **40**: 459-470.

P. 115. *Origanum virens* L. essential oil preservation by encapsulation in starch

Carlos Ribeiro^a, Maria Luísa Beirão-da-Costa^b and Margarida Moldão-Martins^b

carlos.ribeiro@esab.ipbeja.pt, mmoldao@isa.utl.pt

^a Escola Superior Agrária de Beja, Rua Pedro Soares, Apartado 158, 7801-902 Beja Portugal

^b Center for Microbiology and Agriculture Industries (CMIA) DAIAT, Tapada da Ajuda, 1349-017 Lisboa Portugal

Flavour encapsulation protects less-stable flavour components against oxidative and photochemical degradation. Stability of encapsulated flavours depends on the flavour constituents and on the encapsulating agent and process (Anandaraman and Reineccius, 1986).

Encapsulation flavour in starch seems to be an efficient method to preserve aroma compound. According Boutboul *et al.* (2002) aroma-starch interactions mainly result from an adsorption phenomenon involving hydrogen bonds and not from inclusion complexes.

The spherical aggregates produced by spray dryer of starch granules dispersions contain spaces in the form of interconnecting cavities that provide extensive porosity that can be used to protect aroma compounds. This kind of encapsulation is suitable to preserve aromas to be incorporated in foods intending to be subjected to processing at high temperatures.

More than 90% of the encapsulated flavouring on the market is produced by spray drying (Nussinovitch, 1997).

The aim of the present study is the encapsulation of the aroma of *Origanum virens* in a rice starch matrix produced by spray drying.

Starch spherical aggregates are produced by spray drying the dispersion of starch granules in water containing different bonding agents.

Essential oil is produced by steam distillation conducted for 60 minutes at atmospheric pressure. Essential oil was sprayed at different concentrations in the spherical aggregates of starch.

Retention and stability of the main aroma compounds were analysed by GC and GC-MS.

Main compounds of studied *Origanum* were: carvacrol (about 68 %), *p*-cymene (about 4.5 %) and myrcene (about 2.3 %). From the bonding agents tested, carboxymethylcellulose (CMC) produces a smaller size and a higher number of spheres being a good retention medium for essential oil compounds. From the preliminary results it is possible to conclude that the encapsulated essential oil in the starch matrix has a good stability.

Anandaraman S, Reineccius GA (1986) *Food Tech.* Nov. 88: 91-93.

Boutboul A, Giampaoli P, Feigenbaum A, Ducruet V (2002) *Carbohydrate polymers* 47: 73-82.

Nussinovitch A (1997) In: *Hydrocolloid Application. Gum Technology in the food and other industries*, Blackie Acad. & Profs. London: 247-264.

P. 116. Aroma volatiles of rhizomes and roots of cultivated *Rhodiola rosea* L. from Norway

Jens Rohloff^a, Grete Rakvaag^a, Steinar Dragland^b

jens.rohloff@chembio.ntnu.no, steinar.dragland@planteforsk.no

^a The Plant Biocentre, Department of Botany, Norwegian University of Science and Technology NTNU, N-7491 Trondheim, Norway

^b The Norwegian Crop Research Institute (Planteforsk), Apelsvoll Research Centre, Division Kise, N-2350 Nes på Hedmark, Norway

In a 3-year project, 5 different clones of rose root (*Rhodiola rosea* L.) collected from coastal and mountain regions of Norway have been cultivated at the Apelsvoll Research Centre, Division Kise^b. Rhizomes and roots of female and male plants have been analysed by headspace solid-phase microextraction procedure (HS-SPME)^{1,2,3} and gas chromatography coupled with mass spectrometry. As reported earlier, terpenic and aliphatic volatiles were detected as the main aroma compounds contributing to the characteristic rose-like fragrance of the underground parts of *R. rosea*^{1,2}. (*E*)-Pinocarveol (21.8 %) was detected as the main terpenic volatile in the roots of 3-year old plants, as well as cumic alcohol and myrtenol, whereas geraniol, cinnamaldehyde and cinnamyl alcohol showed highest concentrations in the tuberous rhizomes. These results were confirmed by recent HS-SPME analyses (May 2002) of fresh samples of cultivated rose root from Mid-Norway⁴ with (*E*)-pinocarveol and myrtenol as the main compounds in the roots (22.6 % and 19.6 %, respectively). The volatiles of rhizomes were dominated by geraniol (42.1 %) and decanol (36.9 %). Although the drying process (48 h at 35°C) resulted in significantly reduced concentrations, the relative distribution of aroma volatiles was almost unaffected.

Only slight differences were observed when comparing 3-year old female and male plants from Apelsvoll Research Centre analysed by HS-SPME. Higher amounts of phenyl propanoids (cinnamyl alcohol, cinnamaldehyde), oxygen-containing terpenes (cumic alcohol, cuminaldehyde) and aliphatic volatiles (octanol and decanol) were detected in rhizomes and roots of *R. rosea*, whereas higher concentrations of characteristic volatiles such as myrtenol, geraniol and phenylethyl alcohol were observed in female plants.

The essential oil analysis of steam-distilled samples from dried plant material of 2-year old clones (2001) from Apelsvoll Research Centre showed that rhizomes consisted mainly of oxygen-containing terpenes such as geraniol (51.9 %), myrtenol (10.4 %) and the phenyl propanoid cinnamyl alcohol (9.1 %). Straight-chain aliphatic alcohols and acids occurred in minor amounts (10.8 %). These results will be supplemented by further investigations of male and female rose root plants in the summer of 2002, to gain further information about rose root essential oil as a potential source for the production of fragrance and perfume essences.

¹ Rohloff, J. (1999) Monoterpene composition of essential oil from peppermint (*Mentha x piperita* L.) with regard to leaf position using solid-phase microextraction and gas chromatography/mass spectrometry analysis, *J.Agric. Food Chem.*, 47:3782-3786.

² Rohloff, J., Skagen, E.B., Steen, A.H. & Iversen, T.-H. (2000) Production of yarrow (*Achillea millefolium* L.) in Norway: Essential oil content and quality, *J.Agric. Food Chem.*, 48:6205-6209.

³ Rohloff, J. (2002) Essential oil composition of sachalinmint from Norway detected by solid-phase microextraction and gas chromatography/ mass spectrometry analysis, *J.Agric. Food Chem.*, 50:1543-1547.

⁴ Rohloff J (2002) Volatiles from rhizomes of *Rhodiola rosea* L., *Phytochem.*, 59: 655-661.

⁵ Belov VN, Lavrova TV, Vashkevich NG, Mikhailov AY (1994) Extraction of essential oils from plant raw material by steam distillation, *Russ.J.Appl.Chem.*, 67:154-156.

P. 117. Analysis of essential oils and flavonoids of *Dittrichia viscosa* subsp. *revoluta* wild grown and micropropagated plants

M. Costa^a, J. M. F. Nogueira^b, M. G. Miguel^a, A. Romano^b

aromano@ualg.pt

^a Faculdade de Engenharia de Recursos Naturais, Universidade do Algarve, Campus de Gambelas, 8000-817 Faro

^b Departamento de Química e Bioquímica and CCMM, Faculdade de Ciências da Universidade de Lisboa, Campo Grande, Ed. C8, 3º Piso, 1749-016 Lisboa

Dittrichia viscosa W. Greuter (Compositae) is a sub shrub widespread in Europe, especially on Mediterranean areas (Valdés, 1987). *D. viscosa* subsp. *revoluta* (Hoffmanns & Link) P. Silva & Tutinis is an endemic species of Portugal, only growing in the south of Portugal (Pinto da Silva and Tutin, 1973). This sub shrub can be found in waste places and it has been used for years in folk medicine topically as an anti-scabies, anti-inflammatory, and wound-healing agent (Font Quer, 1973). Previous studies reported that this plant has also an anti-ulcerogenic effect, related to its flavonolic components (Alarcon de la Lastra *et al.*, 1993), as well as an anti-inflammatory effect due to the terpene composition of the essential oils (Mañez *et al.*, 1999).

The present contribution reports on the main components of the essential oils from field-grown plants (FP), *in vitro* shoot cultures (InV) and micropropagated plants (MP), obtained from the same clone, determined by capillary GC and GC-MS. This work also describes *in vitro* vegetative propagation of *D. viscosa* subsp. *revoluta*, an endemic Portuguese species, from mature field-grown plants.

The yield of essential oil from *D. viscosa* subsp. *revoluta* isolated from the original field-grown plant (FP) was 0.18% (v/wt). Yields lower than 0.05% (v/wt) were obtained from InV and MP of the same clone. The oils were shown to be a complex mixture of several components. Only 24 constituents were identified, representing 39.9, 42.0 and 49.5% of the total oils from FP, InV and MP samples, respectively. The major constituents of the essential oils were 1,8-cineole (11.0, 5.1 and 5.1%), δ -cadinene (5.0, 7.0 and 6.6%); *E*-nerolidol (4.4, 9.7 and 11.9%); T-cadinol (7.1, 14.4 and 17.4%) and α -cadinol (6.7, 8.4 and 9.4 %), for FP, InV and MP, respectively. Other constituents were found in minor amounts or even at trace levels. The flavonoid fraction was mainly constituted by quercetin with concentrations ranging from 0.3 mg/100 mg FW, in FP, to 48.5 mg/100 mg FW, in InV samples.

Alarcon de la Lastra C, A Lopez, V Motilva (1993) *Planta Medica* 59: 497-501.

Mañez S, M Recio, I Gil, C Gomez, R Giner, J Waterman, J Ríos (1999) *Journal of Natural Products*, 4: 601-604.

Font Quer P (1973) *Plantas Medicinales*. Labor, Barcelona, Vol. 3: 788-9.

Pinto da Silva AR, TG Tutin (1973) *Botanic Journal of the Linnean Society* 67: 290

P. 118. Essential oil of *Johrenia ramosissima* Mozaffarian from Iran

Abdolhossein Rustaiyan^a, Maryam Nikusokhan^a, Zohreh Habibi^b

rustaiyan@excite.com

^a Dept. of Chemistry, Science and Research Campus, Islamic Azad University, Tehran, Iran

^b Dept. of Chemistry, Shahid Beheshti University, Tehran, Iran

Five species of the genus *Johrenia* are found in Iran, all of them endemic. The aerial parts of *J. ramosissima* Mozaffarian, a new endemic species in Iran, were collected from Marzan Abad-Chalous in the Province of Mazandaran in the north region of Iran, in July 1999. The dried aerial parts were hydrodistilled, for 3h, using a Clevenger-type apparatus to yield 0.1% w/w of oil. The oil was analysed by GC/MS. Thirty-five compounds, representing 89% of the total oil, were identified, with bornyl acetate (14.8%), *trans*-pinocarveol (12.5%) and α -campholenol (9.5%) as main constituents.

P. 119. Steam distillation of *Thymbra spicata* oilSerap Sonsuzer^a, Serpil Sahin^a, Levent Yilmaz^b, Mine Kurkcuoglu^c, K. Husnu Can Baser^c

serp@metu.edu.tr

^a M.E.T.U. Food Eng. Dept. 06531 Ankara-TURKEY^b M.E.T.U. Chem. Eng. Dept. 06531 Ankara-TURKEY^c A.U. TBAM 26470 Eskisehir-TURKEY

Thymbra spicata is a thyme-like plant widely grown and used in Turkey. It is usually between 10-50 cm long and it has a black-green colour. Flowers are pink-purple. Leaves have 10-15 mm length and 2-3 mm width. It has various aroma compounds in it. Generally carvacrol is the main compound but thymol chemotypes also exist. Oil of *Thymbra spicata* is used for many purposes such as flavoring of all kinds of food products, as antioxidant and antimicrobial for foods, preparation of some liqueur, for scenting of the soaps and lotions, medical purposes, as an antiseptic, antiplasmatic and antimicrobial.

The spice essential oils have for long been used in seasoning formulations for their obvious advantages over ground spices (Heath and Pharm, 1978). Essential oils are generally uniform in flavor quality. They are sterile and free from all extraneous contaminants. They can readily be incorporate into liquid concentrates and emulsions.

Steam distillation is a special method used to separate high-boiling mixtures or to separate a material from a non-volatile impurity. In this process, temperature of volatilization is lowered by the injection of live or open steam into the charge (Pratt, 1967). Steam distillation is used to separate mixtures at a temperature lower than the normal boiling points of their constituents. In this way, it is similar to vacuum distillation and finds its main use in separating or purifying heat sensitive materials. The principle advantage of using steam is that the volatiles are easily condensed in water (Teranishi *et al.*, 1977). A further advantage of this process is that the steam displaces atmospheric oxygen and hence protects the material from oxidation (Krell, 1963).

The objective of the study is to study the effect of steam flow rate and particle size on yield and composition of essential oil obtained from *Thymbra spicata*.

For all conditions, during the first period of the distillation there is a rapid increase in the amount of oil, but then it remained almost constant. For both flow rates, size was found to be an effective parameter on yield. The higher oil yield was reached with higher flow rate. The composition of oil did not vary with time, size and flow rate.

Heath HB, MBE Pharm (1978) *Flavor Technology: Profiles, Products, Applications*. AVI Publishing Company Inc., USA:231.

Krell E (1963). *Handbook of Laboratory Distillation*. Elsevier Publishing Company, New York.

Pratt HRC (1967) *Countercurrent Separation Processes*. Elsevier Publishing Company, New York.

Teranishi R, EL Murphy, RT Mon (1977) *Journal of Agricultural and Food Chemistry*. 25 (3): 464-470.

P. 120. Antifungal activity of *Thymus* oils on species of Dermatophytes

E. Pinto^a, A. Palmeira^a, L. Salgueiro^b, C. Cavaleiro^b, M. J. Gonçalves^b, C. Pina-Vaz^{c,e}, A. Rodrigues^{c,e},
S. Oliveira^c, C. Tavares^c, J. Oliveira^d

ligia@ff.uc.pt

^a Dept. Microbiology, School of Pharmacy/CEQOFF, R. Aníbal Cunha 4050 Porto, Portugal

^b Lab. Pharmacognosy, Faculty of Pharmacy/CEF, R. do Norte 3000 Coimbra, Portugal

^c Dept. Microbiology, School of Medicine, Alameda Prof. Hernani Monteiro 4200 Porto, Portugal

^d Dept. Obstetrics/Gynecology, School of Medicine, Alameda Prof. Hernani Monteiro 4200 Porto, Portugal

^e IPATIMUP, University of Porto, R. Dr. Roberto Frias 4200 Porto, Portugal

Dermatophytoses are common superficial infections that can be found all over the world. Industrialized drugs used against these affections by official academic medicine are of high economical cost, have undesired secondary effects and fungi strains show each time greater resistance towards them.

Previous research of our team demonstrated the antifungal activity and some of the mechanism of action of some essential oils (EO) (Pina-Vaz *et al.* 2001) against *Candida* species. The aim of this study is to evaluate the antifungal activity of the EO of *Thymus vulgaris*, *T. zygis* subsp. *zygis* and *T. mastichina* subsp. *mastichina* (*Thymus* oils), used in folk medicine, in order to support its application as therapeutical agents in the treatment of dermatophytoses.

The composition of the oils was investigated by gas-chromatography and gas-chromatography/mass spectroscopy (Cavaleiro *et al.* 2001). The Minimal Inhibitory Concentration (MIC), determined according to the NCCLS protocol, M 38-P, was used to evaluate the antifungal activity against dermatophytes strains (*Microsporum canis*, *Trichophyton rubrum* and *Epidermophyton floccosum*). The antifungal activity of their major components was also evaluated. *T. vulgaris* and *T. zygis* oils showed similar antifungal activity, higher than *T. mastichina*. This is probably due to their chemical composition, (*T. vulgaris* mainly composed by carvacrol and *p*-cymene; *T. zygis* mainly composed by thymol and *p*-cymene; *T. mastichina* mainly composed by 1,8-cineole).

The MIC values were determined against three dermatophytes. Results are shown in the table.

EO/Compound	MIC µl/ml (v/v)		
	<i>Epidermophyton floccosum</i>	<i>Trichophyton rubrum</i>	<i>Microsporum canis</i>
<i>Thymus vulgaris</i>	0.078	0.078-0.156	0.078-0.156
<i>Thymus zygis</i>	0.078	0.078	0.078
<i>Thymus mastichina</i>	0.625	0.625	0.625-1.250
1,8-Cineole	5.0	2.5-5.0	5.0
<i>p</i> -Cymene	0.020-0.039	0.020-0.039	0.039
Carvacrol	0.078	0.078	0.039
Thymol	0.156	0.156	0.078

This study showed the antifungal activity of these EO on dermatophytes, particularly *T. vulgaris* and *T. zygis*, which may contribute to the scientific basis for supporting future therapeutical trials on cutaneous mycoses due to dermatophytes.

Acknowledgements: FCT, POCTI and FEDER (POCTI/40167/ESP/2001) for financial support.

Cavaleiro C, Rezzi S, Salgueiro L, Bighelli A, Casanova J, Proença da Cunha A (2001) *Biochem Syst Ecol* 29: 1175-1183

Pina-Vaz C, Rodrigues AC, Pinto E, Costa de Oliveira S, Martinez de Oliveira A, Salgueiro L, Cavaleiro C, Martinez de Oliveira J (2001) *Mycoses* 44 Supp 1: 59

P. 121. Hairy root cultures of *Levisticum officinale*: growth and essential oil production in a bioreactor system

Pedro A. G. Santos^a, A. C. Figueiredo^a, M. M. Oliveira^b, J. G. Barroso^a, L. G. Pedro^a, S. G. Deans^c, J. J. C. Scheffer^d

psantos@fc.ul.pt

^aCentro de Biotecnologia Vegetal, DBV, FCUL, C2, Campo Grande, 1749-016 Lisbon, Portugal

^bITQB / IBET, Quinta do Marquês, Aptd. 12, 2781-901 Oeiras, Portugal

^cDept. of Food Science & Technol., SAC, Auchincruive, Ayr KA6 5HW, Scotland, UK

^dDiv. of Pharmacognosy, LACDR, Leiden University, Gorlaeus Labs, PO Box 9502, 2300 RA Leiden, The Netherlands

Levisticum officinale (lovage) is an Apiaceae herbaceous plant known for its aromatic properties. Lovage hairy root cultures, established as described by Santos *et al.* (2000) and maintained either in SH (Schenk and Hildebrandt 1972) or B5 (Gamborg *et al.* 1968) liquid medium, were inoculated in bubble-column bioreactors containing 1 litre of the same media. Each bioreactor was inoculated with 0.5g fresh weight of hairy roots (corresponding to about 24mg and 15mg dry weight, for SH and B5, respectively), and the experiments were performed in the dark at 24°C. The hairy roots were harvested after 7, 14, 21, 28 and 35 days, and their growth determined in terms of fresh and dry weight. The essential oils from the hairy root samples were isolated by distillation-extraction and analysed by GC and GC-MS.

Hairy roots grown in SH medium attained a maximum fresh weight of 59g.L⁻¹ (4g.L⁻¹ d.w.), at the end of 28 days, while hairy roots grown in B5 medium showed a maximum fresh weight of 23g.L⁻¹ (1g.L⁻¹ d.w.), at the end of 35 days.

Falcarinol was the dominant component of the essential oils from both the SH- and the B5-grown hairy root cultures (10-53% and 10-72%, respectively), but differences were found as to the other major components. *trans*-β-Farnesene (2-20%), *Z*-ligustilide (9-17%), *n*-octanal (2-9%), γ-elemene (2-9%), *Z*-3-butylidenephthalide (1-6%) and pentylcyclohexadiene (traces-5%) were present in large relative amounts in the oils from the SH-grown hairy roots. Apart from falcarinol, the essential oils from the B5-grown hairy root cultures were dominated by *Z*-ligustilide (4-13%), *n*-octanal (2-11%) and limonene (traces-6%).

Acknowledgements: the authors acknowledge the Fundação para a Ciência e a Tecnologia for a PhD Fellowship granted to PAG Santos (PRAXIS XXI/BD/18302/98)

Gamborg OL, RA Miller, K Ojima (1968) *Exp. Cell Res.*, **50**: 151-158.

Santos PAG, AC Figueiredo, MM Oliveira, JG Barroso, LG Pedro, SG Deans, JJC Scheffer (2000) *PSE Meeting - Future Trends in Phytochemistry*, Rolduc, The Netherlands, 15.

Schenk UR, AC Hildebrandt (1972) *Can. J. Bot.*, **50**: 199-204.

P. 122. Biotransformation of terpenes – preliminary studies of microbial activity

S. Schäfer, J. Schrader, D. Sell
silvia.schaefer@dechema.de

Karl-Winnacker-Institut der DECHEMA e.V., Theodor-Heuss-Allee 25, 60486 Frankfurt, Germany

Terpene hydrocarbons can be considered as industrial waste due to their low flavour intensity and their toxicity to the environment. Special biological systems are able to convert terpenes into defined, valuable aroma and fragrance compounds with potential use for the food, cosmetics and pharmaceutical industries. Some of the microorganisms, especially the higher fungi of the ascomycetes and basidiomycetes, are known to have a marked capacity for the biotransformation and bioconversion of terpene precursors.

No terpenoids are produced biotechnologically on an industrial scale despite their often unique organoleptic properties. The main reasons lie in the chemical nature of the substrates and the target molecules: the low water solubility, high volatility and cytotoxicity of terpenes and terpenoids impede processing by conventional bioprocesses. Moreover, due to their microbial metabolic versatility one precursor molecule is very often converted into a wide variety of derivatives, partly representing useless or not readily separable by-products.

For this reason, to achieve an economically viable utilization of terpene resources, the main goal of this interdisciplinary research with different research institutes and industry is to develop an innovative bioprocess capable of reducing or even overcoming the aforementioned constraints. Diverse process features are being investigated, such as activity-controlled substrate feeding and in situ product removal strategies, to obtain long-term optimal growth conditions and thus maximum productivity.

The bio dry weight cannot be used to determine the activity of submerged fungi, either under normal growing conditions or under transformation conditions, due to the formation of pellets. A bioelectrochemical sensor was applied to determine the activity of basidiomycetes. During transformation the microorganisms show different activity depending on the substrate fed. The oxygen uptake rates under different conditions were measured and they can be used as an indicator of fungal activity.

Various fermentations have been performed to optimize the gassing technique and the feeding strategy in order to fulfil the demands of transforming fungi and to optimize the quality of the processed essential oil to be used in the food and cosmetics industry.

P. 123. Fast analysis of natural products by SPME-GC/MS

Karina Schoenefeld, Christine Bartzsch, Karl-Heinz Feller, Joerg Weber

Karina_Schoenefeld@web.de

Department of Medical Engineering, University of Applied Sciences Carl-Zeiss-Promenade 2, 07745 Jena, Germany

Fragrances and flavours are important to natural raw materials based industries. Quantitative limitations and cultivation factors of plant and other natural resources that are difficult to control as well as consumers' preference for natural products and ingredients are driving forces for developing new production methods.

In such cases the qualitative and quantitative analysis of the ingredients of natural products is very important as well from the side of the residual material limitations as from the taste and flavour of food products.

Often there are used complex methods for such purposes. Sampling and preparation and other handling steps are very time consuming. Search of alternative methods is very important especially in industry. There fast results of analysis are required. Additional aspects are that the quantitative analysis of single compounds in complex mixtures as usually done under such condition is not the solution of the problem mentioned.

In such unusual application SPME is an easier practicable method, which gives mainly a profile of the component mixture in natural products together with the information about the relative composition of the product components. Further advantages of SPME are their easy handling, no limit of volume, fast results, long-run investigation and anywhere you can use it. One more advantage of using SPME is the high sensitivity of this method (De la Calle Garcia *et al.*, 1997; Weber *et al.*, 1999). In these cases the variety recognition is the main application field in industrial use.

An automation of considerably investigation is leading to reproducible results.

SPME-GC/MS is applicable to investigate natural products. Furthermore the multiple extraction by SPME can be used for quantitative determination.

Here methodical aspects are shown together with examples from various natural products analysis.

De la Calle Garcia D, M Reichenbacher, K Danzer, C Hurlbeck, C Bartzsch, K-H Feller (1997) *J. High Resol. Chromatogr.* 20: 665.

Weber J, M Beeg, C Bartzsch, K-H Feller, D De la Calle Garcia, M Reichenbacher, K Danzer (1999) *J. High Resol. Chromatogr.* 22: 322.

P. 124. Rapid determination of volatile components in phytopharmaceuticals and cosmetics by headspace SPME-GC analysis

D. Distler, Hartwig Schulz
H.Schulz@bafz.de

Federal Centre for Breeding Research on Cultivated Plants, Institute for Plant Analysis
Neuer Weg 22/23, D-06484 Quedlinburg, Germany

Since 1993 solid phase microextraction (SPME) is available as a commercial product, mainly used for environment and food determinations [Pawlizyn 1999, Schulz 2000]. Only few applications were published in the pharmaceutical field of research. There is a very high demand for analysing active compounds in various phytopharmaceutical products. Therefore the aim of this study was to develop rapid and solvent free applications in order to analyse the volatile components in selected drugs (thyme, camomile, peppermint and marjoram) and related pharmaceutical preparations. Saving of time in sample preparation results in a considerable reduced analysing time especially in case of series determinations.

The results of SPME experiments performed for the plant species mentioned above were determined and compared at different conditions. The influence of the following parameters was studied: quantity of the sample, sample preparation (powdered/not powdered; addition of water, variation of sodium chloride content), temperature of the sample (35° C, 45° C, 55° C, 65° C) and extraction time (5, 10, 15, 20, 25, 30, 40, 60 minutes). Different SPME-fibres were tested at these parameters (fibre coatings: PDMS, PDMS/DVB, polyacrylate, CAR/PDMS, CW/DVB, DVB/CAR/ PDMS).

In order to obtain high reproducibility with respect to equilibration time, temperature as well as homogeneity of the sample, an automated headspace-SPME device was used. This instrument provided also a high sample through-put which is especially important in context with quality control and support of breeding experiments. Generally headspace technique was used in the study which guaranteed extended life-time of the SPME fibres and improved adsorption conditions; this prohibits the adsorption of undesired plant compounds such as sugar components. The valuable plant substances were determined both qualitatively and quantitatively. An internal standard method was applied for quantification of selected volatile substances.

The results of SPME experiments received under different conditions were compared with those obtained by usual GC analysis of the isolated essential oils. The correlation between the SPME methods and the usual GC assays is described by a statistical approximation.

J Pawlizyn (1999) Applications of solid phase microextraction: *RSC Chromatography Monographs* (Series Editor: R.M.Smith), Cambridge UK.

Schulz H, H Krüger, N Herchert, ERJ Keller (2000) Vorkommen flüchtiger Sekundärmetabolite in ausgewählten *Allium*-Wildtypen: *J. Appl. Bot.* **74**, 119-121.

P. 125. Composition and chemical variation during daytime of constituents of the essential oil of *Pogostemon patchouli* Pellet leaves

Magnólia A S. da Silva^a, Polyana A. D. Ehlert^a, Márcia O. M. Marques^b, Lin C. Ming^a

^a Department of Plant Production – Horticulture Sector, College Science, Botucatu, São Paulo, Brazil. P.O Box 237, 18603-970.

^b Phytochemistry Section, Campinas Agronomical Institute, Campinas, SP, Brazil CEP 13075-630.

The patchouli (*Pogostemon patchouli* Pellet) is used in herbal medicine as an aphrodisiac, antidepressant, and antiseptic. Patchouli essential oil is used in aromatherapy to treat skin complaints. It is also employed for headaches and fever. Its essential oil is constituted of benzoic aldehyde, eugenol, cinnamic aldehyde and sesquiterpene patchoulol.

To verify the composition and the chemical variation of its essential oil along 24 hours, leaves of patchouli of medicinal collection plants of the Department of Vegetable Production/Horticulture of UNESP-Botucatu – SP, were collected every three hours during one day (of the 06.00 to 03:00 h), between November, 25 and 26, 2000. The leaves had its oil isolated by hydrodistillation, in Clevenger apparatus during three hours with two repetitions. The chemical composition analysis of the extracted essential oils was carried out by gas chromatography/mass spectroscopy (GC-MS, Shimadzu, QP-5000). The chemical representatives identification was made through the comparative analysis of the mass of spectra of the oil components with the database of the system GC-Em (nist 62.lib), literature and retention index.

The chemical composition of the essential oils was not affected by the schedule of the harvest. The five more abundant components were: a major sesquiterpene (59.44 to 65.22%) not identified yet, α -bulnesene (5.37 to 7.66%), α -guaiene (3.80 to 5.73%), δ -patchoulene (2.54 to 3.36%) and another sesquiterpene (5.58 to 7.44%) also not identified, with retention time quite near to the main component. In bibliographical rising was detected divergences in relation to the majority component (Massada, 1976) and to patchoulol (Betts, 1994), an alcohol sesquiterpene. In our oil samples, it was not detected the presence of eugenol and patchoulol.

Betts, TJ 1994. Evaluation of a Chiral-Val Capillary for the gas-Chromatography of Volatile Oil Constituents, Including Sesquiterpenes in Patchouli Oil, *Journal of Chromatography A*, 664: (2), 295-300.

Massada, Y. 1996. Analysis of Essential Oils by Gas Chromatography and Mass Spectrometry, New York, John Wiley Sons, Inc.

P. 126. Influence of the organic manuring on the content and composition of ginger (*Zingiber officinale* Roscoe) essential oil

Magnólia A S. da Silva^a, Márcia O. M. Marques^b, Lin C. Ming^a

^aDepartment of Plant Production – Horticulture Sector, College Science, Botucatu, São Paulo, Brazil. P.O Box 237, 18603-970.

^bPhytochemistry Section, Campinas Agronomical Institute, Campinas, SP, Brazil CEP 13075-630.

Native to Asia, ginger (*Zingiber officinale* Roscoe) is grown throughout the tropics. Ginger is a well-investigated plant, and its therapeutic benefits are largely due to its volatile oil and oleoresin content. The powdered rhizomes are used in food and to flavour jams, sweets, and ginger ale, and as ingredients for curry. The spices, as the ginger, present variations in their chemical composition due to several factors such as origin area, of the natural hybridisation and of the agriculture-climatic variations (Bartley & Fley, 1994). The objective of this work was to verify the influence of the organic manuring of bird manure in the composition of the essential oil of the ginger.

The experiment was carried out in Department of Vegetable Production - Horticulture of University of Agronomic Sciences of UNESP in Botucatu-SP and in the laboratory of phytochemistry of IAC - Campinas-SP. The experimental area was of 240 m² and the used spacing was of 0,70 x 0,40 m, being 45 plants/plot, and adopted at random in blocks with 4 repetitions. The used treatments were 0, 1, 2, 4, 8 Kg/m² of chicken manure. The rhizomes used in the planting they were Hawaiian type that varied from 20 to 30 grams. The application of the fertiliser was accomplished together in December 2, 2000 and the planting in the 4 of that same month. The harvest was on July 1, 2001. The picked rhizomes were dried at to 30 °C. For the extraction of the essential oil the hydrodistillation method was used, in Clevenger apparatus. The time of distillation was of 1,5 hours. The analysis of the chemical composition of the extracted essential oils was led in gas chromatography/mass spectroscopy (GC-BAD, Shimadzu, QP-5000). The chemical identification was made through the comparative analysis of the mass spectra of the oil components with the database of the system GC-IN (Nist 62.lib) and literature.

The results showed the presence of 34 identified components. Sixty percent of the composition of the oil was formed for α -zingiberene, neral, geranial, β -phellandrene, camphene and β -sesquiphelladrene in all treatments. The treatment T2 showed higher content of the oil. The organic manuring (chicken manure) didn't show significant influence on the composition of ginger essential oil.

Bartely, J.P. & Foley, P. (1994) Supercritical fluid extraction of australian-grown ginger (*Zingiber officinale*) *Journal of the Science of Food and Agriculture*, v.66, n.3, p.365-371.

P. 127. Composition and biological activity of the essential oil from *Aloysia sellowii*

Euclésio Simionatto, Carla Porto, Ubiratan F. da Silva, Solange Cristina Hoenzel, Vinicius Ilha1, Emilia C. M. Dessoy, Ademir Farias Morel*

*afmorel@base.ufsm.br

Departamento de Química, NPPN, Universidade Federal de Santa Maria, 97045-100, Santa Maria, RS, Brazil.

Aloysia (Verbenaceae) is a genus that comprises about 100 species widespread all over the world, mainly in the South and Central America countries. *Aloysia sellowii* (Briq.) [= *Lippia affinis* (Briq.) Moldenke] (Pascual, 2001), known as "garupá", "cidrozinho" or "erva de sepultura" is a large aromatic shrub up to 2-4 meters in height, found throughout the greater part of Southern of Brazil (Rio Grande do Sul), Uruguay, Paraguay and Argentina. In Rio Grande do Sul it is traditionally used by local people as diuretic, stomachic agent and as a remedy for colds, gripe and bronchitis. In our continuing research on the essential oil of Rio Grande do Sul aromatic plants, we have investigated the essential oil of *A. sellowii* and its antibacterial activity.

The essential oils were obtained by hydrodistillation in 1.2-1.4 % yield of the aerial parts from two samples (A and B) of *Aloysia sellowii* from different collection sites of South of Brazil (Rio Grande do Sul), and were analysed by GC, GC/MS, and chiral phase gas chromatography (CPGC). Its constituents were identified by GC/MS using an electron-impact ionisation technique and by co-chromatography with authentic samples.

Fifteen compounds, constituting 87 % of the oil from Santa Maria (sample A), and nineteen compounds, constituting 80 % of the oil from Livramento (sample B), were identified. The major constituent of the sample A was 1,8-cineole (43 %), which is not present in sample B. The oil from sample B was richer in sesquiterpenoids than the oil from sample A, and was characterised by a high content of sabinene (12.6%) and β -Z-santalol (12.9%), which are not present in sample A. In addition, the configurations of the chiral monoterpene constituents were analysed on a dual column system, using two 25 m fused silica columns coated with modified cyclodextrins (2-6-Me-3-Pe- β -CD and 6-Me-2,3-Pe- β -CD) as the chiral stationary phase. Of the chiral monoterpenes identified in the oils of *A. sellowii*, the main constituents were: (-)-sabinene (100% ee), (+)- β -pinene (98% ee), (+)- α -pinene (93% ee), (+)-terpinen-4-ol (66% ee), (+)-limonene (91% ee) and (+)-linalol (76% ee).

The essential oils were screened for biological activity using the brine shrimps (*Artemia salina*) lethality test. Antibacterial activity was determined by the disc diffusion method and by the bioautography method (Homans and Fuchs, 1970). The essential oils of *A. sellowii* (samples A and B) displayed high toxicity (LC₅₀ = 2.85 μ g/mL) in the brine shrimp bioassay. The essential oil of sample A was more active against the tested Gram-positive and Gram-negative bacteria than the oil of sample B. The result suggests that the higher activity of the oil of sample A could be associated with the significant contribution of 1,8-cineole, which is known to possess antimicrobial activity (Sivropoulou, 1997).

1. Homans AL, A Fuchs (1970) *Journal of Chromatography* 51: 327-329
2. Pascual ME, K Slowing (2001) *Journal of Ethnopharmacology* 76: 201-214
3. Sivropoulou A, C Nikolaou, E Papanicolaou, S Kokkini, T Lanaras, M Arsenakis (1997) *Journal of agricultural and food chemistry* 45 (8): 3197-3201

P. 128.**Volatile constituents from *Helietta longifoliata***

Euclésio Simionatto, Neusa F. de Moura, Wellington de Abreu Gonzaga, Carla Porto, Solange C. S. Hoenzel, Ionara I. Dalcol, Ademir F. Morel*

*afmorel@base.ufsm.br

Departamento de Química, Universidade Federal de Santa Maria, RS, Brazil

Helietta longifoliata Britt (Rutaceae), locally called “canela-de-veado”, belongs to the Rutaceae, and is a plant that grows in South America (Southern Brazil, Uruguay, Paraguay and Argentine). It has been used in Brazilian folk medicine as a natural remedy, for the treatment of various diseases (Cruz, 1985).

The aerial parts of the plant (leaves) were submitted for 4 hours to hydrodistillation, using a modified Clevenger-type apparatus to yield 0.70% of yellowish oil $\{n_D^{25}: 1.51, [\alpha]_D^{25}: +22.8$ (in $\text{CHCl}_3, c = 0.01\}$. The essential oil was analysed by GC using a Varian 3800 Gas Chromatograph equipped with a capillary fused silica column (25 m x 0.25 mm) coated with SE-54. The GC conditions used were: carrier gas He (1 ml min⁻¹); on column injector 200 °C; FID 280 °C; column temperature 50 °C to 250 °C at 4 °C/min. GC-MS analyses were performed on a HP 5973-6890 GC-MSD system operating in the EI mode at 70 eV, equipped with a HP-5 cross-linked capillary column (30m x 0.25mm). Column and the injector temperatures were as above. The identification of the components was based on comparison of their mass spectra with those of a NBS Librarie (Massada, 1995) and/or published by Adams (Adams, 1995).

The oil was mainly composed by sesquiterpenes (ca 80%) and a lower amount of monoterpenes (ca 17,5%). Twenty five constituents were identified representing ca of 97,5% of the oil, limonene (17.50%), germacrene-D (16.60%), elemol (11.81%), bicyclogermacrene (11.67%), guaiol (11.53), and α -*epi*-bisabolol (7.24%) being the most abundant components.

The oil was bioactive against some Gram-positive and Gram-negative bacteria, as revealed by bioautography. The microorganisms used in the antibacterial assay *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Micrococcus luteus* (Gram-positive bacteria), *Klebsiella pneumoniae*, *Salmonella setubal* and *Escherichia coli* (Gram-negative bacteria), have been maintained at the Chemistry Department of Santa Maria - RS, Brazil. For the antibacterial assays, 10 μl of a solution corresponding to 100, 50, 25, 12.5, 6.25, 3.12 and 1.06 μg of the oil was applied to pre-coated TLC plates. The inoculum was prepared by culturing each organism in tryptone soya agar (TSA, Oxoid) at 37 °C to a turbidity equivalent to Mc Farland 0.5 standard (1.5×10^8 CFU/mL). One microliter of each diluted inoculum (10^4 - 10^6 CFU) was applied onto Mueller Agar (MHA-DIFCO) plates. Amoxicillin (0.2 μg) was used as positive control.

1. Adams R. (1995) Identification of Essential Oil Components by Gas Chromatography/Mass Spectroscopy. Allured Publishing, Co. Carol Stream, Illinois, 469pp
2. Cruz GL. (1985). Dicionário de Plantas Úteis do Brasil, 597. 3ª edição. Editora Civilização Brasileira S.A. Rio de Janeiro.
3. Massada Y. (1995) Analysis of Essential Oil by Gas Chromatography and Spectrometry. J. Wiley & Sons, New York,

P. 129. Characteristics of the essential oil from fennel flowers in two stages of development

Mirian B. Stefanini^a; M. O. M. Marques^b; L. C. Ming^a; R. Facanali^b; A. F. Ferri^b; M. A. A. Meireles^c

mstefanini@fca.unesp.br

^aDepartment of Plant Production -Horticulture of São Paulo State University -Campus of Botucatu, SP, Brazil

^bNatural Products Laboratory, Agronomical Institute, Campinas, SP, Brazil

^cLASEFI-DEA/FEA, UNICAMP

Foeniculum vulgare var. *dulcis*, a native plant original of Mediterranean, has medicinal uses as antispasmodic, appetite stimulant, stomachic, diuretic, anti-inflammatory, anti-diarrheic, against colics and as a lactation promoter. Several components of the essential oil from this plant have important applications, namely, fenchone is used as counterirritant; limonene is used as solvent, manuf resins, wetting and dispersing agent; trans anethole is used for flavoring agent in perfumary, in soap, pharmaceutical aid (flavor); metyl chavicol or estragole is used in perfumarie and as flavor in foods and liquers; α -Pinene, used in manufacture of comphor, insecticides, solventes, perfume bases.

The present work had as objective to identify the chemical composition of the essential oil of *Foeniculum vulgare* var. *dulcis* in two stages of development. The study was carried in the Lageado Experimental Farm (Agronomical Sciences College) of the Department of Plant Production, Horticulture Section, São Paulo State University, Campus of Botucatu, Brazil. The statistical design was complete randomized block with 2 treatments and 9 replications. Flowers buds and inflorescences in anthesis phase were collected. The essential oil was isolated by hydrodistillation from fresh in Clevenger apparatus; 100g of fresh plants parts were used in each extraction. The extraction proceeded for three hours. The analysis of essential oil was conducted by GC/MS. The means of yield (Tukey Test 5%) were 0,96% for inflorescences in anthesis phase and 0,39% for flowers buds. Statistical differences were observed only in myrcene and *trans*-anethole in essential oil of flowers buds.

Table 1. Relative amount of main components from the essential oil of the flower buds and inflorescences in anthesis from fennel

	Flower buds	Inflorescences in anthesis
<i>trans</i> -Anethole	66.75	59.77
Limonene	20.78	18.36
Fenchone	3.07	6.79
Metyl chavicol	2.47	2.23
α -pinene	1.56	2.12
γ - terpinene	1.05	tr
Myrcene		1.18

P. 130. Yield of essential oil in green seeds of fennel

Mirian B. Stefanini^a, L. C. Ming^a, M. O. M. Marques^b, A. F. Ferri^b, M. A. A. Meireles^c

mstefanini@fca.unesp.br

^a Department of Plant Production -Horticulture of São Paulo State University -Campus of Botucatu - SP - Brazil

^b Natural Products Laboratory, Agronomical Institute, Campinas, SP, Brazil

^c LASEFI-DEA/FEA, UNICAMP

The fennel (*Foeniculum vulgare* Mill.), it is an annual herbaceous plant, whose seeds are very used in the homemade medicine. The fennel seeds produce a yellow-clear aromatic essential oil, used in the production of several liquors drinks and on perfumery. They act as carminative and stimulant. An experiment was carried out at farm (Agronomical Sciences College) of the Department of Plant Production - Horticulture of São Paulo State University - Campus of Botucatu-Brazil. The statistical design used was complete randomized block with 3 replications and 8 treatments (collected of epoch). The seeds were sowings in summer of the 2001 (January 2001) and picked every 14 days, starting from fifteen days after the anthesis until the complete maturation during one year. The parameters evaluated in this work were essential oils yield of green seeds and chemical major compound of the essential oil of fennel. The essential oil was determined through hydrodistillation, in Clevenger apparatus; 50g of green seeds were used in each extraction. The extraction proceeded for three hours. For the Tukey Test (5%) there was statistical difference for the composts and the 5th, 4th, 6th and 3rd harvest epochs respectively, what corresponded between 186 to 228 days after sowing and presented the best results for fresh mass of green seeds. With relationship to the yield of the essential oil, for the Tukey Test (5%), there was not statistical difference among 3th, 4th, 5th, 6th and 8th harvest (Spring) in relationship the others. However the generality means of compounds majorities were: *trans*-anethole (74%), fenchone (15%), limonene (5%), methyl chavicol (3%), α -pinene (1%), γ -terpinene (0.9%) .

Table 1. Means of fresh matter of green seeds of fennel and yield of oil, in function of the collected epochs (treatments). UNESP- FCA/Botucatu – S.P, 2001.

Treatments (collected epoch)	Days after sowing	Seasons of Year	Fresh matter (g)	Yield of oil (ml)
1	158	Spring	308.47 D	0.0 B
2	172	Spring	532.67 CD	0.0 B
3	186	Spring	1018.40 BC	1.88 A
4	200	Spring	1338.13 AB	1.84 A
5	214	Spring	1615.63 A	1.83 A
6	228	Spring	1229.53 AB	1.68 A
7	242	Spring	430.97 D	0.45 B
8	256	Summer	197.33 D	1.53 A

Means followed by same letter in the column and in the line they don't differ significantly, at the level of 5% of probability for the Tukey Test.

P. 131. Supercritical fluid extraction of *Thymbra spicata*Serap Sonsuzer^a, Serpil Sahin^a, Levent Yilmaz^b, S. Gulum Sumnu^a, Temel Ozek^c, K. Husnu Can Baser^c

gulum@metu.edu.tr

^a M.E.T.U. Food Eng. Dept. 06531 Ankara-TURKEY^b M.E.T.U. Chem. Eng. Dept. 06531 Ankara-TURKEY^c A.U. TBAM 26470 Eskisehir-TURKEY

Supercritical fluid extraction (SFE) is a separation method that exploits the unique properties of gases above their critical points to extract soluble components from a raw material (Simandi *et al.*, 1998). SFE is an interesting technique for the extraction of flavor and fragrance compounds because relatively rapid reactions can be carried out under wild conditions and at low temperature and there is no need to evaporate organic solvents (Ozek *et al.*, 1988). The combined liquid-like solvating capabilities and gas-like transport properties of supercritical fluids make them particularly suitable for the extraction of diffusion-controlled matrices such as plant tissues (Scalia *et al.*, 1999).

Carbon dioxide is the most commonly used solvent as a supercritical fluid because of its low critical temperature and pressure ($T_c=31.1$ °C; $P_c=72.8$ atm), its non-toxic and non-flammable properties and its availability in high purity with low cost. It does not react with the food constituents. In addition, it is easily removable from the extract following decompression. Due to its relatively low critical temperature, thermal sample decomposition is reduced, supercritical CO₂ has good solvent properties for extraction of non-polar compounds such as hydrocarbons (Scalia *et al.*, 1999).

Thymbra spicata is a thyme-like plant widely grown and used in Turkey. It has various aroma compounds in it and carvacrol is the main compound.

The objective of this study is to study the effects of temperature, pressure and time on yield and composition of the oil in supercritical CO₂ extraction.

The yield decreased with increasing temperature and remained constant at high values. It was seen that when pressure was increased, yield increased. Monoterpenes were preferably extracted at the beginning of the process. The decrease in monoterpenes for higher levels of temperature, pressure and time may be due to the enhanced leaching of harder to extract substances (sesquiterpenes and oxygenated compounds). Both temperature and pressure positively affected *p*-cymene content in the extract. Sesquiterpene content of essential oil was found to be almost constant for shorter extraction times. However, for long time of extractions, its concentration increased as time increased. In addition, it was seen that sesquiterpene content increased with increasing pressure.

Ozek T, B Bozan, KHC Baser (1988) *Khim. Prir. Soedin.*, 746-751.

Scalia S, L Giuffreda, P Pallado (1999) *Journal of Pharmaceutical and Biomedical Analysis*, 21: 549-558.

Simandi B, M Oszagyan, E Lemberkovics, A Kery, J Kaszags, F Thyron, T Matyas (1998) *Food Research International*, 31 (10): 723- 728.

P. 132. Synergistic activity of components of *Thymus* oils on *Candida* species

C. Tavares^{a,b}, C. Pina-Vaz^{a,b}, E. Pinto^c, A. G. Rodrigues^{a,b}, S. Costa-Oliveira^{a,b}, L. Salgueiro^d, C. Cavaleiro^d,
M. J. Gonçalves^d, A. Palmeira^c, J. Martinez-de-Oliveira^e

cbrj@hotmail.com.

^a Dept. Microbiology, School of Medicine, Alameda Prof. Hernani Monteiro 4200 Porto, Portugal

^b IPATIMUP, University of Porto, R. Dr. Roberto Frias 4200 Porto, Portugal

^c Dept. Microbiology, School of Pharmacy/CEQOFF, R. Anibal Cunha 4050 Porto, Portugal

^d Lab. Pharmacognosy, Faculty of Pharmacy/CEF, R. do Norte 3000 Coimbra, Portugal

^e Dept. Obstetrics/Gynecology, School of Medicine, Alameda Prof. Hernani Monteiro 4200 Porto, Portugal

The raising of fungal infections with the consequent increase of resistance to antifungals urged the search of therapeutic alternatives. Previous research of our team demonstrated antifungal activity of the essential oils (EO) from different species of *Thymus*, in which carvacrol, thymol, *p*-cymene and 1,8-cineole are often the major components. The objective of this study is to evaluate the antifungal activity of each one of these components and the potential synergic effects of their combinations.

For this purpose, thymol/carvacrol, thymol/*p*-cymene, thymol/1,8-cineole, carvacrol/*p*-cymene, carvacrol/1,8-cineole and *p*-cymene/1,8-cineole combinations were studied by the checkerboard method for two clinical isolates of *Candida albicans* and *C. krusei*. The MIC of each component was compared with the MIC obtained with the combination and the fractional inhibitory index (FIX) was calculated. FIX<0.5 means synergism; FIX >0.5 and <4.0 means indifferent interaction; FIX>4.0 means antagonism.

C. albicans and *C. krusei* showed similar behaviour. No antagonism between the components were observed and the most synergic combinations were thymol/1,8-cineole and thymol/*p*-cymene, with FIX=0.125 (corresponding to a decrease of the MIC around 3 dilutions).

This study shows the advantages of the combination of some essential oils or some of their components by the MIC decreasing, and eventually side effects, important issues in future clinical trials on mucocutaneous candidosis.

Acknowledgement: This study was supported by FCT, POCTI and FEDER (POCTI/40167/ESP/2001)

P. 133. The effect of eleven individual essential oil constituents on the phytopathogenic bacterium *Clavibacter michiganensis* subsp. *sepedonicus*

K. Termentzik, K. Karamanolik, N. Trivara, H-I. A. Constantinidou

aterment@agro.auth.gr

Laboratory of Agricultural Chemistry, School of Agriculture, Aristotle University of Thessaloniki, Thessaloniki 54124, Greece

In the present study, the antibacterial activity of eleven monoterpenoids against the phytopathogenic bacteria *Clavibacter michiganensis* subsp. *sepedonicus* was examined. These terpenoids are α -pinene, limonene, 1,8-cineol, α -thujone, fenchon, pulegone, geraniol, carvacrol, α -terpineol, linalool and linalyl-acetate. They are common essential oil constituents of widely distributed Mediterranean ecosystems aromatic plants of *Labiatae* family, such as sage, lavender, rosemary and oregano. The target phytopathogen *C. michiganensis* subsp. *sepedonicus*, is a Gram positive bacterium, which is responsible for the bacterial ring rot of potato.

After determining the optimum growth conditions for the tested bacterial strain in the lab, two methods were applied to examine the antibacterial activity of the compounds in question. With respect to the cell structure (integrity of bacterial membranes), the sensitization of bacteria to lysis effects, after incubation with the tested compounds, was examined. With respect to the cell function (respiratory chain), the inhibitory activity of the terpenoids was tested on the respiration of bacterial cells.

Most of the tested compounds did not remarkably inhibit growth and function of *C. michiganensis* subsp. *sepedonicus* cells compared to their action against other microorganisms. Only carvacrol and geraniol showed a potent antibacterial activity. These oxygenated compounds strongly influenced bacterial respiration and caused enhanced lysis of bacterial cells. The two terpenoids could be further tested as of their probable use as biological antibacterial agents against this pathogen in the field.

P. 134.**Brazilian savannah: alive pharmacy in danger**Suzi Huff Theodoro³ Eliane Mendes Guimarães⁴

suzitheodoro@cds.unb.br; elimengui@hotmail.com

SAS Quadra 05 Bloco H Sala 205. Brasilia - DF/Brazil CEP 70070-914. Center of Sustainable Development – CDS/UnB

Brazil is the 5th country in territorial dimension, the 10th world economy and hold one of the biggest patrimony as far as planet biodiversity is concern. Although the country has this three important indicators Brazil also host strong socio-economic inequality. Beside the importance of the Amazon, the central region of Brazil is characterised by a Biome typical of region of Savannah with an immense natural resource. The vegetation of this region has a diverse use such as food source, medicinal and oil scent potential. The last 30 years this potential has been dramatically reduce by the agriculture in which to grow on the region need to take the natural vegetation losing important species.

Another aspect of this development model is related to the fact that the small land owners who used to live, know and depend on the resources were forced to leave their land and pushed to live on peripheral home in bigger cities. This fact makes them to loose the connection with the land source of food and health. The objective of this paper is to search for solution helping them to revert this situation and work with the low-income population in the surroundings of Brasilia the capital of Brazil. This study aim at recovering the traditional knowledge of needy people on medicine plants and, from that point, offer alternative health treatment to the community which lives in the periphery of the capital. The research consists on a survey of the best known medicine plants in the region of the Brazilian Savannah, identify them, and make their cultivation viable, within a public area where the community has the possibility of getting together and valuing the cultivation of alternative plants. To the survey of the plant species, as well as their therapeutic use, the research-action method was used in weekly meetings with the community, in which the information passed through the participants was noted down and registered. All together, around 50 species were catalogued, from which 30 were chosen for cultivation. The cultivation and production work of the seedling and phytotherapeutic, has been developed by researchers from the University of Brasilia and by the community involved. The soil fertilisation, which is of low quality, is being done under the Stonemeal (1) technique rules, which consists of the use of dust of determined rocks rich in macro and micro nutrients (Ca, Mg, P, V, Mo, among others). To this material, organic fertiliser is added, which has the role of breaking the present clay in the material, and therefore, accelerating the offer of micronutrients. This double fertilisation has the advantage of supplying a production free of agrototoxic, besides encouraging the community to consume better quality products. The results obtained so far demonstrate that the population's traditional knowledge, together with the use of alternative treatments, may be converted into a social insertion mechanism and the preservation of the region's biodiversity.

Theodoro, S. H. - Fertilização da terra pela terra: uma alternativa de sustentabilidade para o pequeno produtor rural. Tese de Doutorado defendida no Centro de Desenvolvimento Sustentável - CDS/UnB em 2000.

³ PhD in Sustainable Development

⁴ PhD student in Sustainable Development

P. 135. Constituents of flowers from *Solidago microglossa* DCP. H. O. Leda, T. C. B. Tomassini

tecoebarto@hotmail.com

Research and Technological Development Department Natural Product Laboratory (PN2)
Far-manguinhos – Fundação Oswaldo Cruz, Rua Sizenando Nabuco, 100 – Manguinhos
Zip code 21041-250 – Rio de Janeiro – Brazil

The genus *Solidago* (*Asteraceae*) comprises hundred twenty species some of them possessing several therapeutic properties (*S. virgaurea*, *S. gingantea*, *S. canadensis*). *Solidago microglossa* DC is a South American species, that in Brazil replaces *Arnica montana* L. for treatment of inflammatory diseases. Clerodane and labdane diterpene substances are present in roots of *S. microglossa* and the epigeal part shows antifungal diterpenes (Vila *et al.*, 2002), while some flavonoids were detected in leaves (Correia, 1998).

The main aim of this work is to isolate, identify and characterise biological active compounds from the ether extract of *S. microglossa* flowers. The extract was chromatographed by TLC, using toluene-ethyl acetate (7:3) as eluting system, and three spots were detected after spraying the plate with an acid anisaldehyde solution. The less polar constituent was re-chromatographed by preparative TLC using hexane-ethylacetate(8:2) as eluting system, affording a pure substance with the following data: *m.p.* 128-130°C; *m/e* = M⁺ 316 (2%), 192 (10%), 111 (20%), 82 (100%); ¹H-NMR in CDCl₃, δ 7.37 (s); 7.27 s; 6.28 (s); 5.71 (s); (CH); four methyl groups at 2.01 (s); 1.19 (s); 1.14 (s); 1.00 (s). These values suggest a diterpene compound that could be responsible for the reported anti-inflammatory activity (Singh, 1999; Demetzos *et al.*, 2001). An attempt to find out sesquiterpene lactones (STL), in *S. microglossa* has failed despite such compounds are abundant in *Arnica montana* L. which have been related to the anti-inflammatory properties of this species.

Correia, E. (1998) Aspectos da Propagação Sexuada e Vegetativa da Arnica Brasileira (*Solidago chilensis* Meyer Asteraceae), *Plantas Mediciniais, Aromáticas e Condimentares*, 2: 193-207.

Demetzos, C.; Dimas, K.; Hatziantoniou, S.; Anastasaki, T.; Angelopoulo, D. (2001) Cytotoxic and Anti-inflammatory of Labdane and cis-Clerodane Type Diterpenes, *Planta Med.*, 67:614-618.

Singh, M.; Pal, M.; Sharma, R.P. (1999) Biological Activity of the Labdane Diterpenes, *Planta Med.*, 65:2-8.

Vila, R.; Mundina, M.; Tomi, F.; Furlán, R.; Zacchino, S.; Casanova, J.; Cañigueral, S. (2002) Composition and Antifungal Activity of the Essential Oil of *Solidago chilensis*, *Planta Med.*, 68:164-167.

P. 136. Constituents of the bark oil of *Cedrelopsis grevei* H. Baillon from Madagascar

Jean François Cavalli^a, Félix Tomi^b, Antoine-François Bernardini^a, Joseph Casanova^b

tomi@vignola.univ-corse.fr

^a Université de Corse, Equipe chimie des produits naturels, UMR CNRS 6134, BP52 20250 Corte, France

^b Université de Corse, Equipe "Chimie et Biomasse", UMR CNRS 6134, Route des sanguinaires 20000 Ajaccio, France

Ptaeroxylaceae family, is present in southern Africa and Madagascar and is constituted by three genus: *Ptaeroxylon*, *Bottegoa* and *Cedrelopsis*. *Cedrelopsis grevei* H. Baillon is an endemic species from Madagascar. This tree, of the vernacular name Katafa or Katrafay (Katra: bitter and fay: juice), measures up to 15 meters high, has grey branches and smelling bark (cedar wood-like odour). This species is abundant in the dry dense forests or in the bush (South and West parts of the island), from the sea level to 900 meters of altitude.

As part of our on-going work on the characterisation of essential oils from Madagascar (1), we investigated the bark oil of *Cedrelopsis grevei*. A commercial sample of *C. grevei* from Madagascar was repeatedly chromatographed and the combined analysis of all the fractions by GC-RI, GC-MS and ¹³C-NMR let to the identification of 116 components. The major constituents were (E)-caryophyllene (9.3%), α -copaene (7.7%), α -selinene (5.8%), δ -cadinene (4.9%), β -selinene (4.5%), α -humulene (3.3%) β -bisabolene (2.8%) and ishwarane (1.5%). The analysis of five other commercial samples confirmed the pre-eminence of sesquiterpenes and exhibited a quantitative chemical variability of the main components ishwarane (17.4-1.0%), (E)-caryophyllene (12.5-1.3%), α -copaene (11.0-4.9%), β -elemene (9.6-0.2%) and α -selinene (9.4-1.1%).

1. Cavalli J.F., Ranarivelo L., Ratsimbason M., Bernardini A.F. and Casanova J. (2001) Constituents of the essential oil of six *Helichrysum* species from Madagascar *Flavour Fragr. J.*, 16, 253-256.

P. 137. Essential oil of *Santolina corsica* Jord. and Fourr. from CorsicaBernard Ferrari, Félix Tomi, Joseph Casanova

tomi@vignola.univ-corse.fr

Université de Corse, Equipe "Chimie et Biomasse", UMR CNRS 6134, Route des sanguinaires 20000 Ajaccio, France

Santolina corsica (*Asteraceae*) is an endemic species growing wild in altitude in the centre of Corsica and in Sardinia [1]. *Santolina* species are known for the pharmacological activity of their extracts containing bioactive compounds such as coumarins [2], or acetylenic compounds [3].

Continuing our research on essential oil bearing plants growing wild in Corsica we report here on the chemical composition of the essential oil from the aerial parts of *Santolina corsica* (plant collected in full blossom). Identification of the individual components was carried out by (i) the GC-retention indices (RI) on apolar and polar columns; (ii) comparison of the mass spectra with those of reference compounds; (iii) ¹³C-NMR spectroscopy according to an experimental procedure and a computerised method developed in our laboratory. In this procedure, the components are identified by comparison of the carbon chemical shift values in the mixture spectrum with those of reference spectra compiled in a computerised data bank [4].

After fractionation by column chromatography, 48 components accounting for 85.1% of the amount of the oil were identified. The main constituents were β-phellandrene (18.0%), β-myrcene (13.7%), santolinatriene (11.9%) and β-pinene (5,7%). The sample is characterised by the presence of 12 irregular monoterpenes. The most important are those bearing the santolinane skeleton: santolinatriene, santolina alcohol (0.2%), (Z)-lyratol (4.4%), (E)-lyratal (0.5%), lyratyl acetate (0.5%) and lyratyl propionate (0.1%). Other irregular compounds are artemisane monoterpenes: artemisia ketone and artemisia alcohol (1.4% and 0.2%, respectively), yomogi alcohol (2.7%). Finally, we identify lavandulol (0.2%). The simultaneous occurrence of the three irregular skeletons was never reported in an essential oil of *Santolina* species, while this profile is commonly encountered in *Artemisia* oils [5]. The composition of our sample differs drastically from that of *S. corsica* oil from Sardinia where camphor, artemisia ketone, borneol and aromadendrene were reported as main components [1].

1. Poli F., Bonsignore L., Loy G., Sachetti G., Ballero M. (1997) *Journal of Ethnopharmacology* 56: 201-208.
2. Maqua M.P., Vines A.C.G., Caballero E., Grande M.C., Medarde M., Bellido I.S. (1988) *Phytochemistry* 27: 3664-3667.
- 3 Christensen L.P. (1992), *Phytochemistry* 31: 7-49.
- 4 Tomi F., Bradesi P., Bighelli A. Casanova J. (1995) *J. Magn. Reson. Anal.* 1: 25-34,.
- 5 Stojanovic G., Palic R., Mitrovic J., Djokovic D. (2000), *J. Essent. Oil Res.* 12: 621-624.

P. 138. Essential oil composition from *Melaleuca quinquenervia* of New Caledonia

Bénédicte Trilles^a, Saliou Bouraïma-Madjebi^b

coustenoblefh@canl.nc

^a Unité de Production d'Espèces Végétales, Laboratoire de Biologie et Physiologie Végétales Appliquées, New Caledonia

^b Laboratoire de Biologie et Physiologie Végétales Appliquées, New Caledonia

Melaleuca quinquenervia (Cavanilles) S. T. Blake (Myrtaceae), common name Niaouli, is an indigenous species of New Caledonia Island.

Essential oil of Niaouli is produced in secretory glands located in the leaves. The essential oils were extracted by hydrodistillation.

In this study, 135 samples collected from 7 harvesting locations in New Caledonia were analysed by GC/MS and Principal Component Analysis (PCA – statistic method). The chemical composition of the essential oil of *M. quinquenervia* allowed the definition of six chemotypes. The oil yield is in ranged between 0% and 3%. The industrial mean of niaouli oil yield was 0.7%. The industrial essential oil selection criterion was the 1,8-cineol content above a concentration of 60% (yes is 60% it's a typing mistake).

The different characteristics of *M. quinquenervia* preserve by in vitro culture.

P. 139. Examination of antimicrobial activity of different medicinal plant extracts by impedimetry

M. Tulok^a, Cs. Mohácsi-Farkas^b, B. Balogh^a

mtulok@omega.kee.hu

^a Department of Medicinal and Aromatic Plants, SZIE University, P.O. Box 53 H-1502 Budapest, Hungary

^b Department of Microbiology and Biotechnology, SZIE University Sornlói út 14-16. H-1118 Budapest, Hungary

In food processing there is an increasing consumer demand on food products with the quality characteristics of fresh products, and foods containing only natural compounds. The extracts of medicinal or aromatic plants are one of the potential biopreservatives (Smid and Gorris 1999). There are various methods that are used to investigate antimicrobial activity of essential oils. One of the most successful method of the rapid and automated microbiological techniques is based on electrical measurements. Microbial metabolism usually results in an increase in both conductance and capacitance causing a decrease in impedance of the culture media. Inhibitory activity of essential oils and supercritical fluid extracts of *Origanum vulgare* ssp. *hirtum* (Link) Letswaart, *Satureja montana* L. and their main components was tested by impedimetry on a number of food-borne bacteria and fungi.

Greek oregano (*Origanum vulgare* ssp. *hirtum*, epithet of Department of Medicinal and Aromatic Plants) and winter savoury (*Satureja montana* "Bokroska", variety of Department of Medicinal and Aromatic Plants) grown at the Research Station of Department of Medicinal and Aromatic Plants of SZIE, Soroksár were used. *Essential oils* were produced by hydrodistillation in Clevenger apparatus according to VII. Phg. *Supercritical fluid extracts* (SCFE) were obtained by Isco SFX 2-10 laboratory-scale apparatus. Each sample was analysed by using Shimadzu GC-14B equipped with a flame ionisation detector.

Inhibitory activities were investigated by using a RABIT type automated impedimeter from Don Whitley Scientific (Shingley, U.K.). Stationary growth phase cultures of the test organisms were included in triplicates into Don Whitley impedimetric broth (for bacteria, direct technique) and yeast-extract peptone glucose broth (for fungi, indirect technique (Owens *et al.* 1989) and the conductance of the cultures were recorded automatically for 48 hours at 30°C without or after addition of essential oils or SCFEs in concentrations of 0.1 % and 0.05% (v/v). Viable cell counts of the cultures were estimated initially, the impedimetric detection time (TTD) were automatically determined by the instrument.

In comparison of herb extracts the highest antimicrobial activity was detected in case of *Origanum vulgare* subsp. *hirtum*. Results of GC/MS analyses show that the main component of herb extracts investigated is carvacrol. The results obtained demonstrated that using carvacrol as an antimicrobial agent had stronger antimicrobial activity than herb extracts.

Owens JD, DJ Thomas, RS Tompson *et al.* (1989) *Lett. Appl. Microbiol.*: 9, 245-249.

Smid EJ, GM Gorris (1999) *Handbook of food protection*, New York, Marcel Dekker

P. 140. Composition and antifungal activity of the essential oil from aerial parts and rhizomes of *Valeriana dioscoridis*

Olga Tzakou^a, Maria Couladis^a, Milica Pavlovic^b, Marina Sokovic^c

tzakou@pharm.uoa.gr

a Department of Pharmacognosy and Chemistry of Natural Products, School of Pharmacy, University of Athens, Panepistimiopolis Zografou, 157 71 Athens, Greece

b Faculty of Pharmacy, Department of Pharmacognosy, Vojvode Stepe 450, 11000 Belgrade, Yugoslavia

c Department of Plant Physiology, Institute for Biological Research "Sinisa Stankovic", 29 Novembra 142, 11000 Belgrade, Yugoslavia

Valeriana genus (Valerianaceae) comprises about 200 species and is represented in Europe by 20 species. All the members of the genus have a characteristic foetid odour.

Valeriana species are widely used in Western phytotherapy for the preparation of phytomedicines with mild sedative and antispasmodic properties (Wichtl, 1994; Blumenthal, 1998). The best known species is *V. officinalis*, whose oil composition has been the subject of considerable study. As part of on going study on the essential oils of *Valeriana* sp., in this work we report on the qualitative and quantitative analyses of the volatile constituents produced by the aerial parts and rhizomes of *Valeriana dioscoridis*, one of the several valerian species mentioned by Dioscorid.

Valeriana dioscoridis Sibth. & Sm. grows in the Balkan Peninsula, in rock-crevices, rocky woods and damp grassland. This species has a very short rhizome, with a cluster of fusiform tubers and stem solitary, slightly hairy, 25-75 cm high.

The oil analyses were performed by GC/MS and the identification of the compounds was based on comparison of their Kovats indices (KI), their retention times (RT) and mass spectra with those obtained from authentic samples and/or the NIST/NBS Wiley libraries.

The main components of the oil from the aerial parts and the oil from the rhizome were patchouli alcohol (14.3%, 16.3% respectively) and α -pinene (7.1%, 9.0% respectively). There were no substantial qualitative differences between these two oils.

The antifungal activities of *V. dioscoridis* rhizome oil and patchouli alcohol were analysed by modified microdilution technique (Daouk *et al.*, 1995). The results showed that the oil from rhizomes and the main component exhibited variable degrees of antifungal activity against the strains tested. The essential oil tested showed higher antifungal properties than patchouli alcohol. MICs of the oil were 1.0-10.0 μ l/ml, while MFCs were 1.5-15.0 μ l/ml. The most resistant fungal strains were *Aspergillus flavus*, *Trichoderma viride*, *Penicillium ochrochloron* and *Penicillium funiculosum*, while *Cladosporium cladosporioides*, *Fusarium tricinctum* and *Phomopsis helianthi* and dermatomycetes were the most sensitive ones.

Blumenthal M (ed.) (1998) The Complete German Commission E Monographs. American Botanical Council, Austin, Texas.

Bos R (1997) Analytical and Phytochemical studies of Valerian and Valerian-based Preparations. Thesis, State University of Groningen, The Netherlands.

Daouk KD, MS Dagher, JE Sattout (1995) Journal of Food Protection 58: 1147-1149.

Houghton PJ (1997) Valerian, the genus Valeriana. Harwood Academic, The Netherlands.

Wichtl M (1994) Herbal Drugs and Phytopharmaceuticals. Medpharm Scientific Publishing, Stuttgart.

P. 141. Chemical composition of the essential oils from *Thymus samius* and its related taxa *Thymus atticus* and *Thymus parnassicus*Olga Tzakou^a, Theophanis Constantinidis^b

tzakou@pharm.uoa.gr

^a Department of Pharmacognosy and Chemistry of Natural Products, School of Pharmacy, University of Athens, Panepistimiopolis Zografou, 157 71 Athens, Greece^b Institute of Systematic Botany, Department of Biotechnology, Agricultural University of Athens, Iera Odos 75, 11855 Athens, Greece

The genus *Thymus* L. (Lamiaceae) includes more than 350 taxa. *Thymus* species have been used for more than 2,000 years as medicinal herbs and many of them are still in use. *Thymus samius* is an endemic plant known only from the island of Samos (E. Greece). Morphologically, it resembles *Th. atticus* but has narrowly spatulate, pubescent leaves. It may be of hybrid origin, with *Th. cilicicus*, *Th. zygiformis* and *Th. parnassicus* as possible parents (Baden, 1991).

The aim of the present study was to compare the constituents of the essential oil from *Th. samius* with those from *Th. atticus* and *Th. parnassicus* growing in various localities of C. Greece. All samples collected during the flowering period. The dried aerial parts were subjected to hydrodistillation for 3h and the obtained oils were analysed by GC/MS. The identification of the compounds was based on comparison of their Kovats indices (KI), their retention times (RT) and mass spectra with those obtained from authentic samples and/or the NIST/NBS Wiley libraries.

The major components of *Th. samius* oil were germacrene-D (26.40%) and β -bisabolene (22.73%), two sesquiterpenes that arise from the precursor *E,Z*-farnesyl cation in the mevalonate pathway, as well as (*E*)-caryophyllene (15.48%) a sesquiterpene that arise from *E,E*-farnesyl cation.

The oils from all samples of *Th. parnassicus* oils were qualitatively similar but display some quantitative differences. Those from Mt. Pateras and Mt. Gerania samples were rich in (*E*)-caryophyllene (13.10%, 10.02% respectively). In the oil from Mt. Parnitha sample the main components were (*E*)-caryophyllene (8.48%), linalyl acetate (8.24%) and γ -terpinene (8.04%).

Th. atticus oil from Mt. Parnitha contained the monoterpenes 1,8-cineole (11.81%) and β -myrcene (7.32%), and the sesquiterpenes (*E*)-caryophyllene oxide (9.18%) and (*E*)-caryophyllene (6.66%) as main constituents. The oil from the same species grown on Mt. Parnassos was dominated by the sesquiterpenes (*E*)-nerolidol (17.49%), (*E*)-caryophyllene (15.84%), germacrene D (11.50%) and by the acyclic monoterpene β -myrcene (9.45%).

Another related taxon *Th. cilicicus* has been studied from Turkey (Akgul *et al.*, 1999). The oil from Beysehir region was found to contain α -terpineol (16.4%), camphor (9.7%), 1,8-cineole (7.8%) and α -pinene (6.9%); the sample from Silifke region had as main components α -pinene (16.7%), 1,8-cineole (10.4%), isoborneol (9.2%), *cis*-verbenol (8.2%), camphor (6.4%) and *trans*-verbenol (6.2%); the sample from Ermenek region revealed α -terpineol (33.44%) as the major component and camphor, citronellol and linalool (ca. 6-8% each). All samples of *Th. cilicicus* are rich in monoterpenes, whereas *Th. samius* and its related taxa *Th. atticus* and *Th. parnassicus* are rich in sesquiterpenes with the exception of *Th. atticus* sample from Mt. Parnitha. None of the studied oils belong to the most abundant phenol-rich chemotype.

Akgul A, M Ozcan, F Chialva, F Monguzzi (1999) *J. Essent. Oil Res.* 11:209-214.Baden C (1991) *Thymus*L. in *Flora Mountain of Greece*. Strid A & Kit Tan (eds), Edinburgh University Press, Edinburgh.Mabberley DJ (1997) *The Plant-Book*. 2nd edition, Cambridge University Press. Cambridge.

P. 142. Antimicrobial properties and geographical variation in essential oil composition of fever tea - *Lippia javanica* (Verbenaceae)

S. Subramoney^a, S. F. van Vuuren^a, A. M. Viljoen^a, B. Demirci^b, K. H. C. Başer^b

vanvuurensf@therapy.wits.ac.za

^a Department of Pharmacy and Pharmacology, Faculty of Health Sciences, University of the Witwatersrand, 7 York road, Parktown 2193, South Africa

^b Medicinal and Aromatic Plant and Drug Research Center (TBAM), Anadolu University, 26470-Eskişehir, Turkey

Lippia javanica is a widely spread woody shrub and the major traditional use is reflected in its vernacular name; fever tea and ‘koorsbossie’. An infusion of the leaves is also used as a decongestant for colds and coughs. Infusions may also be used topically to treat scabies and lice. This study shows that the essential oil chemistry varies dramatically both within and between natural plant populations and thus has a direct impact on variations of antimicrobial activity. The aerial parts were collected from individual plants in five natural populations. The hydrodistilled essential oils were analysed by GC-MS and a cluster analysis was performed on the essential oil dataset. From sixteen samples (representing 5 natural populations), five chemotypes were identified;

- a myrcenone-rich type (36 – 62%),
- a carvone-rich type (61 – 73%),
- a piperitenone-rich type (32 – 48%),
- an ipsenone rich-type (42 – 61%) and
- a linalool-rich type (>65%)

The myrcenone and linalool chemotypes have been mentioned in the literature but the carvone, ipsenone and piperitenone chemotypes have not previously been reported. The oil showed minimal activity against *Staphylococcus aureus*, *Escherichia coli*, and *Bacillus cereus*, and no apparent activity against *Pseudomonas aeruginosa* in a disk diffusion assay. The oil did however show activity against *Candida albicans* and *Cryptococcus neoformans*, in the same assay. Time kill studies were performed on three microbial respiratory isolates to document the scientific rationale of using *Lippia* for respiratory complaints in traditional herbal medicine. *Klebsiella pneumonia*, *Cryptococcus neoformans* and *Bacillus cereus* showed reduction in microbial populations with the strongest bacteriostatic effect observed for *Klebsiella pneumonia*.

P. 143. The antimicrobial activity and death kinetics of the essential oil and chemical components of the South African endemic, *Osmitopsis asteriscoides*

S. F. van Vuuren^a, A. M. Viljoen^a, E. Ernst^b, B. Demirci^c, T. Ozek^c, K. H. C. Başer^c

vanvuurensf@therapy.wits.ac.za

^a Department of Pharmacy and Pharmacology, Faculty of Health Sciences, University of the Witwatersrand, 7 York road, Parktown 2193, South Africa

^b College of Pharmacy, The University of Iowa, Iowa City, Iowa 52242, USA.

^c Medicinal and Aromatic Plant and Drug Research Center (TBAM), Anadolu University. 26470-Eskişehir, Turkey

The essential oil composition and anti-microbial activity of *Osmitopsis asteriscoides*, a medicinal plant used in traditional herbal preparations in South African is investigated. A preliminary screening was done using the disc diffusion method on nine bacterial and four fungal isolates. Time kill studies indicated a more detailed bacteriostatic progression with the antimicrobial determination of *Candida albicans*, *Pseudomonas aeruginosa* and *Staphylococcus aureus*. A strong fungicidal activity against *Candida albicans* was found at concentrations ranging from 0.5 to 2 %. Sixty-six compounds were identified by GC-MS in the hydro-distilled essential oil, which contained (-)-camphor and 1,8-cineole as the main constituents. The high concentration of these two terpenoids is presented as a possible explanation for the traditional use of *Osmitopsis asteriscoides* for treating microbe-related illnesses. A comparative antimicrobial time kill study of *Candida albicans* with 1,8-cineole and camphor standards as well as in combination were compared with the pure *Osmitopsis asteriscoides* essential oil. The time-kill method is presented as a superior assay to determine antimicrobial properties of essential oil and its constituents.

P. 144. Modelling of the lemongrass essential oil supercritical extraction process

Rubem M. F. Vargas^a, Eduardo Cassel^a, Luiz G. Longhi^a, Ana C. Atti-Santos^b, Marcia R. Pansera^b, Luciana Atti-Serafini^b

cassel@eq.pucrs.br

^a DEQ - PUCRS, Ipiranga 6681, Porto Alegre – RS, 90619-900, Brazil. FAX (0XX)51 33203625

^b Instituto de Biotecnologia – UCS, Francisco Getúlio Vargas 1130, 95070-560, Caxias do Sul – RS, Brazil

The extraction process of the lemongrass essential oil with supercritical CO₂ is studied in this work. We used plants of the experimental cultivated stands from Southern Brazil and the HP7680 apparatus to extract the essential oils. The oils were extracted from leaves and the following process parameters were used in this study: 40°C to 80°C, 100 bar, and 2mL/min (Table 1). The analysis was performed by GC and GC/MS.

Table 1 – Extraction Experimental Data (P = 100 bar)

Time	% mass concentration (g/g)				
	5 min	10 min	15 min	20 min	30 min
Temperature					
40°C	1.08	1.22	1.23	1.23	1.23
50°C	0.73	0.84	0.85	0.85	0.85
60°C	0.56	0.69	0.71	0.72	0.72
70°C	0.70	0.83	0.86	0.86	0.86
80°C	0.69	0.79	0.80	0.81	0.81

The aim of this work is simulate the lemongrass oil extraction process at supercritical condition. To perform this objective a mass transfer model is necessary and we guess the procedure proposed by Reverchon (1996). In this model two parameters are adjusted: the diffusivity of the solute in the solid matrix and the partition coefficient. Therefore the behaviour of these parameters is analysed in the function of the temperature and we present a correlation for the diffusivity and partition coefficient adjusted parameters. The extract yield curve is presented, comparing the experimental results with the model. The differences obtained between calculated and experimental data to solute concentrations profile, confirm that this model is efficient to represent the supercritical extraction of the lemongrass essential oil.

Reverchon, E. (1996) Mathematical Modeling of Supercritical Extraction of Sage Oil, *AIChE J.*, 42(6), pp. 1765-1771.

P. 145. African wormwood – essential oil composition, geographical variation and antimicrobial properties of a coveted traditional herbal remedy

T. Pelele^a, A. M. Viljoen^a, S. F. van Vuuren^a, B. Demirci^b, K. H. C. Başer^b

viljoenam@therapy.wits.ac.za

^a Department of Pharmacy and Pharmacology, Faculty of Health Sciences, University of the Witwatersrand, 7 York road, Parktown 2193, South Africa

^b Medicinal and Aromatic Plant and Drug Research Center (TBAM), Anadolu University. 26470-Eskişehir, Turkey

Artemisia afra (Jacq. Wild) also known as African wormwood, “umhlonyane”(Xhosa, Zulu), “lengana” (Sotho, Tswana) and Wildeals (Afrikaans) is an aromatic shrub belonging to the family of Asteraceae. It is widespread in South Africa extending from the mountainous regions of the South Western Cape, along the eastern coast to the Northern Province. Due to the popular use of *A. afra*, herbal tinctures have been prepared for commercial distribution. As the chemotypic variation remains unrecorded it has been impossible to standardise extracts containing *A. afra*. The aerial parts of 17 samples from four natural populations were hydrodistilled and the essential oil analysed by GC-MS and tested for antimicrobial property on a number of bacteria and fungi. The essential oil composition varied quantitatively and qualitatively within and between populations. With the aid of cluster analysis several chemotypes have been identified based on the presence and quantity of the following compounds; α -thujone (5.55-77.65%), β -thujone (1.37-57.73%), camphor (1.00-48.99%), 1,8 cineole (2.31-50.09%), Artemisia ketone (14.47-27.97%), Artemisia alcohol (9.31-27.76%) and santolinyl acetate (4.14-24.32%). The essential oil was active against all organisms except *P. aeurigosa* and *E. faecalis*. The oil exhibited more antifungal properties than antibacterial effects. The pure standards (thujone, cineole, camphor etc) showed no antimicrobial activity and it was concluded that the oil could exert antimicrobial properties be working in a synergistic way. *Artemisia afra* varies within and between natural populations and standardising of commercial products will be problematic without cloning of a favourable chemotype.

P. 146.**Synthetic odoriferous lactones**

Czesław Wawrzeni^a, Małgorzata Grabarczyk^a, Anna Nagielska^b, Robert Obara^c, Maia Szmigiel-Pieczewska^a, Antoni Szumny^a

C-Waw@OZI.AR.WROC.PL


^a Department of Chemistry, Agricultural University, Norwida 25, 50-375 Wrocław, Poland, e-mail: c-waw@ozi.ar.wroc.pl

^b Firmenich Sp. z o.o., Chrzanowska 10, 05-825 Grodzisk Mazowiecki, Poland

^c Institute of Chemistry, Świętokrzyska Academy, Chęcińska 5, 25-020 Kielce, Poland

Compounds with lactone moiety are widely spread in nature. They are components of many essential oils, usually responsible for their odoriferous properties (Ohlof, 1994). Lactones are also present in fruits, vegetables and other foods (Maga, 1976). We have also observed that some of terpenoid lactones synthesized by us possess very interesting odours. Recently we have published the synthesis and odour characteristics of some terpenoid lactones with limonene system (Paruch *et al*, 2000).

Now we present the synthesis of further mono-, bi- and tricyclic odoriferous terpenoid lactones. The structures of some of them are given below.



Lactones were obtained in four or five step syntheses from simple or natural starting materials *via* the iodolactonization of γ,δ -unsaturated acids. The saturated lactones were synthesized from iodolactones by their reductive (nBu₃SnH) dehalogenation. The unsaturated ones were obtained by dehydrohalogenation of iodolactones with DBU.

Maga J. A. (1976) *Crit. Rev. Food Sci. Nutr.*: 8, 1-56

Ohlof G. (1994) *Scent and Fragrances*, Springer Verlag, London, Berlin, 1-238

Paruch E., Ciunik Z., Nawrot J., Wawrzeni^c C. (2000) *J. Agric. Food. Chem.*: 48, 4973-4977

P. 147. Developments in SPME for fast analysis of herb components for quality control and variety recognition

Jörg Weber, Christine Bartzsch, Karl-Heinz Feller

joerg.weber@fh-jena.de

Department of Medical Engineering, University of Applied Science Carl-Zeiss-Promende 2, 07745 Jena, Germany

In quality control fast results of analysis are required. Traditional Methods are often time consuming. A lot of several activities has to be done. In some cases automation is hardly applicable. Therefore our main intention was to develop a fast, easy and representative method for analysing volatile compounds in natural sources. The use of headspace solid-phase microextraction (HS-SPME) coupled to gas chromatography offers a lot of advantages not only in the analysis of herbs or other essential oil containing materials (De la Calle Garcia *et al.*, 1997; Weber *et al.*, 1999)

For quantification of maceration extracts a useful method was developed. Influences of extraction process will be discussed on thyme and chamomille maceration extracts. These are examples for liquid samples analysed by HS-SPME.

A more interesting challenge is the analysis of solid essential oil containing samples. Differences to liquid samples in handling and extraction force the development of a new handling procedure. The results to be presented show the efficiency of the new developed method. Quantitative analyses were performed on solid thyme and chamomille samples as well as on flavoured candles. In addition, several varieties of thyme were investigated for variety recognition using the new method.

De la Calle Garcia D, M Reichenbächer, K Danzer, C Hurlbeck, C Bartzsch, K-H Feller (1997) *J. High Resol. Chromatogr.* 20: 665.

Weber J, M Beeg, C Bartzsch, K-H Feller, D De la Calle Garcia, M Reichenbächer, K Danzer (1999) *J. High Resol. Chromatogr.* 22: 322.

P. 148. Composition of *Achillea filipendulina* essential oil

T. Wenham, J. Birkby

tracey.wenham@botanix.co.uk

Botanix Ltd, Hop Pocket Lane, Paddock Wood, Tonbridge, Kent.TN12 6DQ. UK

Achillea fillipendulina is a deciduous perennial grown for its attractive summer and autumn flowers for the dried flower market. It is a member of the Yarrow family and is also known as Cloth of Gold. *Achillea fillipendulina* has flat heads of bright yellow flowers which last all summer, feathery foliage and grows to heights of 1.2m (4ft). The composition of *Achillea fillipendulina* has previously been reported in the literature (Maffei *et al*, 1994).

The flowering aerial parts of *Achillea fillipendulina* were collected from a farm in Kent. The aerial parts were extracted by hydro-distillation using a Clevenger – type apparatus to produce 0.07% oil yield.

The essential oil was analysed by combined GCMS using a DB-1 column (30m x 0.32mm x 1µm) and a BP20 wax column (25m x 0.32mm x 0.25µm). Temperature programmed retention indices of the compounds were determined relatively to the retention times of a series of *n*-alkanes. The constituents of the oil were identified by GCMS comparison with reference libraries (NIST98) and comparison of retention indices and GCMS of those reported in the literature (Adams, 1989; Adams, 1995; Jennings *et al.*, 1980). Quantification was determined by peak area normalisation without consideration of calibration factors.

Over thirty compounds were identified and the main constituent was 1,8-cineole (45.8%). The remaining oil was made up of a number of major and minor fractions. The major fractions included monoterpene alcohols (13.8%), monoterpene hydrocarbons (15.4%), monoterpene esters (10.2%) and sesquiterpenes (8.9%). The minor fractions included monoterpene ketones (1.7%), other oxygenated monoterpenes (0.4%) and sesquiterpene oxides (1.7%) and sesquiterpene alcohols (2.1%).

The composition of this oil was significantly different to that reported by Maffei *et al*, where the principal component was found to be artemisia acetate.

Adams, R. P. (1989) *Identification of essential oils by ion trap mass spectroscopy*, Academic press London.

Adams, R. P. (1995) *Identification of essential oil components by gas chromatography / mass spectroscopy*, Allured publishing corporation Illinois USA.

Jennings, W. , Shibamoto, T. (1980) *Qualitative analysis of flavor and fragrance volatiles by glass capillary gas chromatography*, Academic press London.

Maffei, M., Mucciarelli, M., Scannerini, S. (1994), *Biochem. Syst. Ecol.* **22**, No. 7, pp. 670-687

P. 149. Composition of methyl esters in European *Humulus lupulus* essential oils

T. Wenham, J. Birkby

tracey.wenham@botanix.co.uk

Botanix Ltd, Hop Pocket Lane, Paddock Wood, Tonbridge, Kent.TN12 6DQ. UK.

Methyl esters are extracted from the essential oil of *Humulus lupulus* (hop) and used as downstream aroma and flavour products in brewing and the flavour and fragrance industry. Methyl esters have principally been extracted from the Target hop variety. This research attempts to determine the methyl ester profile of twelve other European hop varieties by extracted ion GCMS of the whole essential oils.

Twelve varieties of *Humulus lupulus* and the Target variety were obtained from various parts of Europe and the essential oils were extracted by hydro-distillation using a Clevenger-type apparatus. The methyl esters were extracted from target hop oil using liquid-liquid extraction and column chromatography.

The essential oils were analysed by combined GCMS using a DB-1 column (30m x 0.25mm x 1µm). The two principal methyl ester ions 74 and 87 amu were extracted from the total ion chromatogram and the temperature programmed retention indices of the methyl esters were determined relatively to the retention times of a series of n-alkanes. The methyl esters were identified by GCMS comparison with reference libraries (NIST98) and comparison of retention indices of those reported in the literature (Adams, 1989; Adams, 1995; Jennings *et al.*, 1980). Quantification was determined by peak area normalisation without any consideration of calibration factors.

The extracted ion technique was verified by matching the extracted Target methyl ester profile to the extracted ion methyl ester profile from the target whole oil. The methyl ester profiles of the twelve hop oil varieties were grouped into four categories. The first category consists of the Target hop variety and eight of the twelve other hop oil varieties which all contain the major constituents methyl 4-decenoate and methyl 4,8-decadienoate. The second category consists of one variety, which contains the major constituent methyl heptanoate. The third category consists of one variety, which contains the major constituent methyl 9,12-octadecadienoate. The fourth category consists of two varieties which contain the major constituents methyl tetradecanoate, methyl hexadecanoate, methyl 9,12-octadecadienoate and methyl 13, 16-octadectrienoate.

Adams, R. P. (1989) *Identification of essential oils by ion trap mass spectroscopy*, Academic press London.

Adams, R. P. (1995) *Identification of essential oil components by gas chromatography / mass spectroscopy*, Allured publishing corporation Illinois USA.

Jennings, W. , Shibamoto, T. (1980) *Qualitative analysis of flavor and fragrance volatiles by glass capillary gas chromatography*, Academic press London.

P. 150. Optical purity of (R)-(-)-1-octen-3-ol in the aroma of various species of edible mushrooms

Renata Zawirska-Wojtasiak, Erwin Wąsowicz
renazaw@owl.au.poznan.pl

Institute of Food Technology of Plant Origin, Agricultural University of A. Cieszkowski
Wojska Polskiego 31, 60-624 Poznań, Poland

Mushrooms have long been used as a food or a food flavouring material because of their unique and subtle flavour. The characteristic flavour of certain mushrooms species have been well documented [Wąsowicz, 1974; Fischer & Grosch, 1987; Mau, 1992 and 1997; Venkateshwarlu *et al.*, 1999]. It is established that the main odorant of mushrooms is 1-octen-3-ol. It occurs in mushrooms in predominant minus optical form [Chambers *et al.*, 1998].

The characteristic enantiomeric ratio of 1-octen-3-ol in various species of edible mushrooms was the subject of the investigation. The microdistillation-extraction apparatus of Likens-Nickerson was used for aroma isolation. The volatiles were separated on usual GC capillary column HP-5 PHMS, and on chiral GC column RtβDex s.a. connected to HP-6890 gas chromatograph. The following species were analysed: *Agaricus bisporus* (two varieties), *Pleurotus ostreatus*, *Hericium erinaceum*, *Xerocomus badius* and *Lentinula edodes* known as *Shii-take*. The samples of *Agaricus bisporus*, *Pleurotus ostreatus* and *Hericium erinaceum* were grown on few kind of growing media.

Despite significant differences stated in the flavour profiles of investigated mushrooms, in all of the species 1-octen-3-ol was one of the most abundant compounds: at the highest concentration in *Agaricus bisporus* (about 60% of total volatiles). In all of the species the predominant form of this compound there was (R)-(-)-1-octen-3-ol. Its optical purity was the highest in *Agaricus bisporus* (over 98.9%) in both varieties, but also very high in the other mushrooms: *Pleurotus ostreatus* (over 97.2%), *Hericium erinaceum* (over 96.8%), *Xerocomus badius* (over 88.9%) and *Shii-take* (over 95.5%)

The data obtained allow to suggest high optical purity of (R)-(-)-1-octen-3-ol in the edible mushroom aroma independently from the species. It can serve as the criterion of aroma authenticity of mushroom flavouring substances on the market. Stereochemistry has to be considered when studying xenobiotics, agrochemicals, food additives, flavour and fragrances [Maier *et al.*, 2001]. Optical purity of some odorants can be useful in aroma authenticity control system [Mosandl, 1995]. Several mushroom-like flavouring substances were controlled in this work. In all of them the racemic ratio of 1-octen-3-ol was stated, thus mean not natural origin of flavour.

It was also found that SPME (solid phase microextraction) used instead of distillation can be beneficial in the control system because of its convenience and speed. The enantiomeric ratios measured by SPME – with polydimethylsiloxane fiber- were just the same as with distillation.

- Chambers E., A.C. Smith, L.M. Seitz, D.B. Sauer (1998) *Development in Food Science* 40: 173-180
Fischer K.H., W. Grosch (1987) *Lebensm.-Wiss. u.-Technol.* 20: 233-236
Mau J.L., R. Beelman, G. Ziegler (1992) *J.Food Sci* 3: 704-706
Mau J.L., Ch. Chyau, J. Li (1997) *J. Agric. Food Chem.* 45: 4726-4729
Maier N., P. Franco, W. Lindner (2001) *J. Chromatog. A* 906: 3-33
Mosandl A. (1995) *Food Rev. Int.* 11: 597-664
Venkateshwarlu C. (1999) *Flavour & Fragrance Journal* 14: 191-194
Wąsowicz E. (1974) *Bull. Pol. Sci.*, 22: 143-151

Index



<i>Species</i>	
<i>Abisidia glauca</i>	106
<i>Achillea filipendulina</i>	192
<i>Achillea pachycephala</i>	50
<i>Achyrocline saturooides</i>	120
<i>Acore calamus</i>	93
<i>Acorus calamus</i>	100
<i>Aeolanthus gamwelliae</i>	113
<i>Agaricus bisporus</i>	194
<i>Agastache foeniculum</i>	151
<i>Agathosma betulina</i>	118
<i>Agathosma crenulata</i>	118
<i>Agonis obtusissima</i>	115
<i>Aldrovanda vesiculosa</i>	96
<i>Aloysia sellowii</i>	171
<i>Aniba canelilla</i>	51, 121
<i>Aniba fragrans</i>	51
<i>Aniba rosaeodora</i>	51
<i>Anthemis altissima</i>	98
<i>Anthurium salvadorensis</i>	17
<i>Artemisia afra</i>	189
<i>Artemisia sp.</i>	76
<i>Artemisia vulgaris</i>	99
<i>Aspergillus niger</i>	34, 142
<i>Asterella blumeana</i>	139
<i>Bortyosphaeria dothidea</i>	34
<i>Botrytis cinerea</i>	78
<i>Brosimum gaudichaudii</i>	120
<i>Calea clematidea</i>	138
<i>Cananga odorata</i>	37
<i>Carapa guianensis</i>	51
<i>Carum carvi</i>	93
<i>Caryocar brasiliense</i>	120
<i>Cedrelopsis grevei</i>	180
<i>Chamomilla recutita</i>	126
<i>Chorella pyrenoidosa</i>	34
<i>Cistus albidus</i>	149
<i>Citrus aurantifolia</i>	53
<i>Citrus reticulata</i>	87
<i>Citrus sinensis</i>	53
<i>Copaifera spp.</i>	51
<i>Coriander sativum</i>	67
<i>Coryanthes sp.</i>	17
<i>Crithmum maritimum</i>	153
<i>Cryptocarya moschata</i>	66
<i>Cryptotermis brevis</i>	52, 53
<i>Cyclotrichium longiflorum</i>	148
<i>Cymbopogon flexuosus</i>	100
<i>Cymbopogon schoenanthus</i>	104
<i>Cymbopogon winterianus</i>	100
<i>Cymbopogon winterianus</i>	52
<i>Cyphomandra divaricata</i>	17
<i>Dionaea muscipula</i>	96
<i>Dipteryx odorata</i>	51
<i>Dittrichia viscosa</i>	161
<i>Drosera rotundifolia</i>	96
<i>Drosophyllum lusitanicum</i>	96
<i>Eleuterococcus senticosus</i>	107
<i>Eremaea pauciflora</i>	115
<i>Eremanthus erytropappus</i>	59
<i>Eryngium creticum</i>	49
<i>Eryngium vesiculosum</i>	150
<i>Eucalyptus globulus</i>	75
<i>Eucalyptus stoatei</i>	115
<i>Fagara zanthoxyloides</i>	41
<i>Foeniculum vulgare</i>	68, 132, 145, 173, 174
<i>Frullania sp.</i>	32
<i>Gennaria diphylla</i>	81
<i>Hedychium gardnerianum</i>	128
<i>Helietta longifoliata</i>	172
<i>Heracleum sphondylium</i>	95
<i>Hericium erinaceum</i>	194
<i>Hoya vs. sussuela</i>	17
<i>Humulus lupulus</i>	193
<i>Hymenophyton flabellatum</i>	147
<i>Hypericum androsaemum</i>	82, 141
<i>Hypericum calycinum</i>	141
<i>Hypericum elodes</i>	141
<i>Hypericum glandulosum</i>	47
<i>Hypericum hircinum</i>	141
<i>Hypericum humifusum</i>	141
<i>Hypericum joerstadii</i>	47
<i>Hypericum linarifolium</i>	141
<i>Hypericum montanum</i>	141
<i>Hypericum perforatum</i>	141
<i>Hypericum perforatum</i>	141
<i>Hypericum pubescens</i>	141
<i>Hypericum pulchrum</i>	141
<i>Hypericum sp.</i>	141
<i>Hypericum tomentosum</i>	141
<i>Hypericum undulatum</i>	141
<i>Johrenia ramosissima</i>	162
<i>Juniperus horizontalis</i>	101
<i>Juniperus oxycedrus</i>	94
<i>Kunzea ericoides</i>	118
<i>Kunzea pulchella</i>	115
<i>Laurus azorica</i>	70
<i>Laurus nobilis</i>	109, 114
<i>Lavandula latifolia</i>	152
<i>Lentinula edodes</i>	194
<i>Leptospermum scoparium</i>	118
<i>Levisticum officinale</i>	48, 105, 144, 165
<i>Licaria puchuri-minor</i>	51
<i>Lippia adoensis</i>	140
<i>Lippia affinis</i>	171
<i>Lippia alba</i>	63, 64, 134
<i>Lippia grandis</i>	46
<i>Lippia javanica</i>	186
<i>Lippia rugosa</i>	140
<i>Lonicera nummularifolia</i>	135
<i>Lycaste depepei</i>	17
<i>Lychnophora ericoides</i>	120
<i>Malleostemon tuberculatus</i>	115
<i>Marchantia polymorpha</i>	34
<i>Melaleuca alternifolia</i>	118
<i>Melaleuca quinquenervia</i>	182
<i>Melaleuca uncinata</i>	115
<i>Mentha pulegium</i>	56, 57
<i>Mentha x piperita</i>	19

<i>Micromeria juliana</i>	71	<i>Stachys pilifera</i>	125
<i>Myrtus communis</i>	57	<i>Stachys sylvatica</i>	122
<i>Nardostachys chinensis</i>	34	<i>Stellaria media</i>	133
<i>Nepeta haussknechtii</i>	91	<i>Syzygium cuminii</i>	41
<i>Ocimum americanum</i>	100	<i>Syzygium travancoricum</i>	41
<i>Ocimum basilicum</i>	52, 58	<i>Targionia lorbeeriana</i>	139
<i>Ocimum spp.</i>	124	<i>Teucrium lusitanicum</i>	65
<i>Origanum dictamnus</i>	93	<i>Teucrium polium</i>	72
<i>Origanum majorana</i>	33, 146	<i>Teucrium polium</i> subsp. <i>vicentinum</i>	65
<i>Origanum sp.</i>	116	<i>Thryptomene australis</i>	115
<i>Origanum virens</i>	159	<i>Thryptomene kochii</i>	115
<i>Origanum vulgare</i>	93, 154, 183	<i>Thymbra capitata</i>	131
<i>Osmitopsis asteriscoides</i>	187	<i>Thymbra spicata</i>	163, 175
<i>Osyris alba</i>	79	<i>Thymi herba et extracta</i>	55
<i>Palisota sp.</i>	17	<i>Thymus atticus</i>	185
<i>Papaver somniferum</i>	110	<i>Thymus caespititius</i>	130
<i>Passiflora chocoensis</i>	17	<i>Thymus camphoratus</i>	130, 131
<i>Pelargonium sp.</i>	118	<i>Thymus mastichina</i>	130, 131, 164
<i>Pentapleura subulifera</i>	54	<i>Thymus parnassicus</i>	185
<i>Peristylus cordatus</i>	81	<i>Thymus samius</i>	185
<i>Pimpinella anisum</i>	80, 86	<i>Thymus sp.</i>	116, 176
<i>Pinus sp.</i>	53	<i>Thymus vulgaris</i>	20, 55, 73, 97, 156, 164
<i>Pinus sylvestris</i>	108	<i>Thymus zygis</i>	130, 164
<i>Pittosporum undulatum</i>	74, 127	<i>Valeriana dioscoridis</i>	184
<i>Plagiochila fruticosa</i>	34	<i>Vanillosmopsis erythropappa</i>	59
<i>Plectranthus ecklonii</i>	84	<i>Vitex agnus-castus</i>	33
<i>Plectranthus laxiflorus</i>	85	<i>Xerocomus badius</i>	194
<i>Plectranthus sp.</i>	77	<i>Zanthoxylum rhoifolium</i>	89
<i>Plectranthus verticillatus</i>	84	<i>Zanthoxylum xanthoxyloides</i>	41
<i>Pleurotus ostreatus</i>	194	<i>Zanthoxylum zanthoxyloides</i>	41
<i>Pogostemon patchouli</i>	169	<i>Zingiber officinale</i>	170
<i>Pothomorphe peltata</i>	123		
<i>Pothomorphe umbellata</i>	123		
<i>Pterospartum tridentatum</i>	155		
<i>Rhodiola rosea</i>	160		
<i>Rosmarinus officinalis</i>	57, 116, 152		
<i>Salvia glutinosa</i>	103		
<i>Salvia nemorosa</i>	117		
<i>Salvia officinalis</i>	83, 90, 97, 116, 117, 152		
<i>Salvia pratensis</i>	117		
<i>Salvia sclarea</i>	117		
<i>Salvia triloba</i>	116		
<i>Sanguisorba officinalis</i>	112		
<i>Santalum insulare</i>	158		
<i>Santalum spicatum</i>	39		
<i>Santolina corsica</i>	181		
<i>Satureja fruticosa</i>	31		
<i>Satureja isophylla</i>	92		
<i>Satureja montana</i>	69, 157, 183		
<i>Satureja sp.</i>	116		
<i>Schusterella sp.</i>	32		
<i>Scrophulia ningpoensis</i>	137		
<i>Sideritis sp.</i>	35		
<i>Solidago microglossa</i>	179		
<i>Spiranthera odoratissima</i>	120		
<i>Spondias dulcis</i>	45		
<i>Spondias mombin</i>	45		
<i>Spondias purpurea</i>	45		
<i>Stachis acerosa</i>	125		
<i>Stachis benthamiana</i>	125		

Subject

Activities of essential oils	23
Adverse effects of essential oils	38
African wormwood	189
Amazon aromatic plants	51
Angelica root	118
Antidepressant	37
Antimicrobial activity	41, 59
Antioxidant activity	131
Attenuated total reflectance technology	29
Banana	60
Basil	52, 58, 111
Benzoin	38
Bergamot	38
Biodiversity	20
Biological activities	23, 40, 41, 58
Biosynthesis of essential oils	19
Biotransformation	34, 142, 166
Blashed thistle	116
Blood pressure	37
Breath-by-breath nosespace analysis	60
Calamus	38
Calcium channels	118
Camomile	59, 116, 126, 168
Chemical structures	21
Chiral odoriferous compounds	119
Citronella	52
Citrus	118
Clary sage	116
Coriander	67
Cosmetic applications	129
Cyclodextrins	30
Data Bank of the Amazon aromatic plants	36
Dill	118
Dragonhead	116
Ecological aspects	21
Emotional parameters	37
Enantiomeric ratio	194
Enantiomers	30, 136
Enantioselective cyclodextrin derivatives	30
Extraction methods	116
Fennel	118, 145, 173, 174
Fever tea	186
Feverfew	116
Fractional distillation and pervaporation	27
Fragrance notes	88
Frankincense	118
Genetic polymorphism	20
Geranium	118
Ginger	170
Glandular structures	47
Grape fruit	34
Guinea-pig	118
Hairy roots	80, 86, 165
Headspace sorptive extraction	18
Headspace SPME-GC analysis	168
Headspace trapping	17
Headspace-solid phase microextraction	18, 76, 79, 90, 110
Healing rites	40
Herbs and spice plants	28
High speed GC	18
Human liver microsomes	136
Human physiological parameters	37
Human physiology and subjective evaluation	39
Hyacinth	28
Hydroponics	28
Impedimetry	183
Intellectual property	22
Jasmin	28
Jasmine	118
Laurel	70, 114
Lavender	116, 118, 152
Lemon	28
Lemongrass	188
Lime	28
Liverworts	32, 139, 147
Lovage	105
Mandarin	87
Manuka	118
Manure	170
Marjoram	168
Mental parameters	37
Microbial activity	166
Microbial transformation	106
Microorganisms	34
Miglyol 812N	86
Mushrooms	194
Near-infrared spectroscopy	29
Ninde	113
NIR-FT-Raman spectroscopy	29
Nutmeg	118
Olfactometry	140
Olfactorimetry	110
Orange	53
Orange concentrated essences	27
Patchouli	169
Patents	22
Peppermint	118, 168
Percutaneous absorption	39
Perfumery materials	22
Peru balsam	38
Pervaporation	27
Pharmacological effect	118
Physiological activities	23
Phytopesticidal	100
Phytosanitary industry	93
Proton-Transfer-Reaction Mass-Spectrometry	60
Psychological effects	23
Pungent	118
Quantitative structure activity relationship	61
Rain forests	17
Repeated measurement analysis	33
Rose	28, 129
Rose root	160
Rosemary	62, 152
Sage	83, 152
Sample preparation and analysis	18
Sandalwood	39, 61

Sassafras	38
Sensitisers	38
Skeletal muscle	118
Smooth muscle	118
Solid phase extraction.....	17
Solid phase microextraction	62, 168, 191
Spasmogenic.....	118
Spasmolysis	118
SPME-GC/MS.....	167
Statistical procedure	33
Steel corrosion	102
Strawberry	28
Supercritical carbon dioxide extraction	31
Supercritical fluid extraction	31, 175, 188
Sympatholytic effect.....	37
Synthetic odoriferous lactones.....	190
Tea tree	118
Termite	52, 53
Thyme.....	118, 156, 168
Trichomes	19, 84, 85, 117, 122, 134
Uterus	118
Valerian	118
Vibrational spectroscopy methods.....	29
Volatile signals	21
Wintergreen	38
Wormseed.....	38
Ylang-ylang	37

Authors


Abdalla, A. E.	143
Acamovic, T.	73
Agnaniet, Huguette	129
Albert, H.	40
Alves, Sydney H.	138
Amorim, Lúcia R.	82
Andrade, Eloisa Helena A.	36, 45, 46, 121
András, Cs.	48
Antonelli, M.	122
Antunes, Teresa	47
Apáti, P.	48
Araújo, Carla C. de	66
Asakawa, Yoshinori	32, 34, 142, 147
Ascensão, Lia	84, 85
Atti-Santos, Ana C.	188
Atti-Serafini, Luciana	188
Ayoub, Nahla A.	112
Aytaç, Z.	54, 148
Balogh, B.	183
Bamasian, Shahrzad	50
Bandion, Franz	110
Barata, Lauro E. S.	51
Bardon, A.	32
Baron, Vincent	158
Barros, N. M.	52, 53
Barroso, José G.	31, 47, 69, 74, 80, 81, 84, 85, 86, 130, 131, 153, 154, 155, 165
Bartsch, Christine	167, 191
Başer, K. Hüsnü Can.	35, 40, 54, 76, 77, 78, 79, 95, 103, 106, 113, 148, 163, 175, 186, 187, 189
Bazylo, A.	55
Beirão-da-Costa, Maria Luísa	159
Ben Fadhel, N.	57
Ben Fadhel, Najeh	56
Ben Salah, A.	57
Bernardini, Antoine-François	72, 180
Bernáth, J.	145
Bettini, Mércia de Fátima M.	27
Bianchini, Jean-Pierre	158
Bicchi, Carlo	30, 141
Biljana Božin	58
Birbeck, Steve	39
Birkby, J.	192, 193
Blažević, Nikola	151, 152
Blüthner, Wolf-Dieter	146
Bouraïma-Madjebi, Saliou	182
Boussaid, M.	56, 57
Braggins, J.	32, 147
Brandão, M. G. L.	59
Brekalo, Marijana	152
Brevard, Hugues	60
Brkić, Dragomir	151
Brophy, Joseph J.	115, 150
Brunelli, Claudio	30
Buchbauer, Gerhard	23, 37, 39, 41, 61, 110, 140
Burillo, J.	31, 69
Butaud, Jean-François	158
Cabral, Lourdes Maria Corrêa	27
Camacho, C.	70
Campos, L. B.	128
Carreira, Léa Maria M.	36
Carvalho, L.	130, 131
Carvalho, M. A. Pinheiro	81
Casanova, Joseph	72, 180, 181
Cassel, Eduardo	62, 188
Castilho, P.	70
Castro, Dulce M.	63, 64, 134
Castro, Marília de M.	84
Cavaleiro, Carlos	65, 164, 176
Cavalheiro, Alberto J.	66
Cavalli, Jean François	180
Chartone-Souza, E.	59
Chatzopoulou, P. S.	67
Choudhary, M. Iqbal	78
Christensen, L. P.	97
Chung, H. G.	68, 105
Coelho, J. A.	31, 69
Constantinidis, Theophanis	185
Constantinidou, H-I. A.	177
Copeland, Lachlan M.	150
Cordero, Chiara	30
Costa, M.	161
Costa, M. Céu	70
Costa-Oliveira, S.	176
Couladis, Maria	71, 132, 133, 184
Cozzani, Stéphanie	72
Cross, D. E.	73
Croteau, Rodney	19
Cunha Luís, Tiago	74
Currais, António José M.	74
Dalcol, Ionara I.	89, 172
Dalfovo, V.	52, 53
Damjanovic, Biljana	75
Damjanovic, Jovanka	75
Davin, L. B.	127, 128
Davis, Edward M.	19
Deans, S. G.	165
Dellacassa, Eduardo	87
Demirci, Betül.	35, 40, 54, 77, 103, 113, 148, 186, 187, 189
Demirci, Fatih	76, 78, 79, 95, 106
Desjobert, Jean-Marie	72
Dessoy, Emilia C.	89, 138, 171
Dienes, Eszter	156
Discola, Karen F.	51
Distler, D.	168
Dragland, Steinar	160
Duarte, Carla S.	80
Duarte, Fernanda	141
Duman, Hayri	35
Dzurillay, Á.	48
Ehlert, Polyana A. D.	169
Ekici, M.	54
Elkamali, Hatil H.	104
Elkamali, Hizabr H.	104
Ernst, E.	187
Facanali, R.	173
Fadlalla, Babiker	104

Falcalani, Roselanine.....	66	Kamalinejad, M.	98
Falcato Simões, M.	130, 131	Kamiya, N. I.	32
Farooq, Afgan.....	78	Karamanolik, K.	177
Feller, Karl-Heinz.....	167, 191	Kardali, Mohamed.....	101
Fernandes, Francisco M.....	81	Katsiotis, S. T.	67
Fernandes, Priscila C.....	124	Kawahigashi, Tatsuo	102
Fernandes-Ferreira, Manuel.....	82, 83, 139	Kaya, Ayla.....	103
Ferrari, Bernard.....	181	Kéry, Á.	48, 116
Ferreira, Cláudia.....	84, 85	Khalid, H. S.	143
Ferreira, Nicolau J.	74	Khalid, Hassan E.	104
Ferri, A. F.	123, 173, 174	Kim, S. M.	68, 105
Figueiredo, A. Cristina.....	31, 47, 69, 74, 80, 81, 84, 85, 86, 130, 131, 153, 154, 155, 165	Kirimer, Neşe.....	35, 106
Flach, Adriana.....	138	Kolalite, M. R.	107, 108
Fortes, I. C. P.	59	Koteyeva, N. K.	108
Frąckowiak, Bożena.....	119	Kovačević, Nada.....	109
Francke, Wittko.....	21	Kovatcheva, Assia.....	61
Franz, Chlodwig M.....	33, 90, 146	Krist, Sabine.....	110
Frizzo, Caren D.	53, 87	Krüger, H.	111
Fronza, E.....	53	Kubeczka, Karl-Heinz	112
Furlani, Pedro R.....	124	Kujundžić, Sebastijan.....	132
Furusawa, Mai.....	34, 142	Kürkçüoğlu, Mine.....	95, 113, 163
Gearon, Valerie.....	39	Kuštrak, D.....	114
Giacomelli, Sandro R.	89	Lakusic, B.....	71
Gomes, Fabiana.....	62	Langbehn, Jan.....	146
Gomes, J. I.	123	Lassak, Erich V.....	115
Gomes, Paula B.	88	Leda, P. H. O.	179
Gonçalves, M. J.	164, 176	Lemberkovics, É.	48, 116, 117
Gonzaga, Wellington de A.	89, 172	Lewis, N. G.....	127, 128
Grabarczyk, Małgorzata	190	Leydet, Alain.....	129
Grassi, Paolo.....	90	Lima, Waterloo Napoleão.....	121
Guedes, Ana P.	82	Linares, Carlos E. B.....	138
Habibi, Zohreh.....	91, 92, 162	Lindinger, Werner.....	60
Hart, Stephen.....	118	Lindsey, K. L.....	40
Hashimoto, Toshihiro.....	34, 142	Lis-Balchin, Maria.....	38, 118
Hassan, Mohamed A.....	104	Lobo, Carlos.....	47
Hatch, S.	40	Lochyński, Stanisław.....	119
Hernandez Ochoa, Leon.....	93	Longhi, Luiz G.	188
Heuberger, Eva.....	39	Lopes, Lucia.....	120
Hillman, K.	73	López, Ginés.....	94
Hoenzel, Solange Cristina.....	171, 172	Lousã, Mário.....	141
Hongratanaworakit, Tapanee.....	37	Machado, Silvia R.	63, 64, 134
Ilhal, Vinicius.....	171	Maia, José Guilherme S.....	36, 45, 46, 121
Iñigo, Ana.....	94	Maistry, K.	77
İşcan, Gökalp.....	95	Maleci Bini, L.....	122
Ivanova, Alexandra N.....	96, 107	Marcelo Curto, M. J.....	141
Jafari, A.	135	Marczal, G.	117
Jakobsen, H. B.	97	Märk, Tilmann.....	60
Jamalian, Az.	98	Marques, Márcia O. M.....	63, 64, 66, 123, 124, 134, 169, 170, 173, 174
Jancic, R.	71	Martinez-de-Oliveira, J.....	176
Javidnia, K.	98, 135	Martins, Marcus Vinicius de Miranda.....	120
Jerković, I.	114	Masoudi, Shiva.....	50, 125
Jin, Baoping.....	28	Massoudi, Abdolnasser.....	126
Jirovetz, Leopold.....	41, 140	Mastelić, J.....	114
Jovanović, Marina.....	133	Mata, Vera.....	88
Junghanns, Wolfram.....	146	Matavulj, Milan.....	58
Kaiser, Roman.....	17	Mayr, Dagmar.....	60
Kakasy, A.	116	McDevitt, R.	73
Kalemba, Danuta.....	99	Medeiros, H.	127
Kalita, M. C.	100	Medeiros, J. R.	127, 128

Meireles de Sousa, Inês G.....	74	Pansera, Marcia R.....	188
Meireles, M. A. A.....	173, 174	Paolillo, A.....	122
Mendes Guimarães, Eliane.....	178	Partl, Anamarija.....	151, 152
Mendes, R. L.....	31, 69	Patel, Subha M.....	28
Mendonca, S. C.....	128	Pavkov, Ružica.....	58
Menut, Chantal.....	129	Pavlovic, Milica.....	184
Messaoud, C.....	57	Pedro, Luis G.....	47, 74, 80, 81, 84, 85, 86, 130, 131, 153, 154, 165
Miguel, M. Graça.....	65, 80, 86, 130, 131, 161	Pelele, T.....	189
Mihajlović, Bisererka.....	133	Peralba, Maria do Carmo R.....	62
Milos, Mladen.....	157	Pereira, A. P.....	31, 69
Mimica-Dukić, Neda.....	58, 132, 133	Pereira, Ana L.....	155
Ming, L. C.....	174	Pérez-Alonso, Maria José.....	94, 149, 150
Ming, Lin C.....	63, 64, 134, 169, 170, 173	Pérez-Alonso, María José.....	101
Miri, R.....	98, 135	Phukan, S.....	100
Miyazawa, Mitsuo.....	136, 137	Pina-Vaz, C.....	164, 176
Mkaddem, M.....	56	Pinto, E.....	164, 176
Mohácsi-Farkas, Cs.....	183	Pizzolato, Tânia M.....	62
Mohammed, Zeinab A.....	143	Pluhár, Zsuzsanna.....	144, 156
Moldão-Martins, Margarida.....	159	Porto, Carla.....	171, 172
Montero, Jean Louis.....	129	Proença da Cunha, A.....	65
Mookherjee, Braja D.....	28	Radonić, Ani.....	157
Moraes, M.....	123	Raharivelomanana, Phila.....	158
Morais, Rui.....	139	Rahman, Atta-ur-.....	78
Moreira, L. F.....	59	Rakvaag, Grete.....	160
Morel, Ademir F.....	89, 138, 171, 172	Ribeiro, Carlos.....	159
Mouloungui, Z.....	93	Ristić, Mihailo.....	109
Moura, Neusa F.....	172	Rodrigues, A.....	70, 164
Muravnik, L. E.....	96	Rodrigues, A. E.....	88
Muselli, Alain.....	72	Rodrigues, A. G.....	176
Nagielska, Anna.....	190	Rohloff, Jens.....	160
Nascimento, A. M. A.....	59	Romano, A.....	161
Nawwar, Mahmoud A. M.....	49	Rubiolo, Patrizia.....	18, 30, 141
Nemeth, E.....	68, 105	Rustaiyan, Abdolhossein.....	50, 125, 162
Neves, Marta.....	139	Sabet, R.....	135
Ngassoum, Martin Benoit.....	41, 140	Sahin, Serpil.....	163, 175
Nikusokhan, Maryam.....	162	Salgueiro, Lígia.....	65, 164, 176
Nishimatsu, N.....	34	Salvatera-Garcia, Maria.....	22
Nogueira, J. M. F.....	161	Santos, Pedro A. G.....	155, 165
Nogueira, Teresa.....	141	Santos-Gomes, Paula C.....	83
Noma, Yoshiaki.....	34, 106, 142	Sanz, J.....	149
Nour, Amal M.....	143	Sarkarzadeh, H.....	98
Novák, I.....	144, 145	Sbeghen, A. C.....	52, 53
Novak, Johannes.....	33, 90, 146	Schäfer, S.....	166
Nsanzimana, Jean Baptiste.....	99	Scheffer, J. J. C.....	80, 86, 165
Obara, Robert.....	190	Schoenefeld, Karina.....	167
Okuno, Yoshiharu.....	137	Schrader, J.....	166
Oliveira, G. B.....	59	Schulz, Hartwig.....	29, 168
Oliveira, J.....	164	Sedaghat, Sajjad.....	91, 92
Oliveira, M. M.....	165	Selas, Mafalda.....	84
Oliveira, S.....	164	Sell, D.....	166
Omatsu, I.....	147	Seong, N. S.....	105
Onishi, S.....	34	Serafini, Luciana A.....	52, 62
Özcan, Musa.....	153, 154	Sevinate-Pinto, Isabel.....	47
Özek, Temel.....	148, 175, 187	Shafī, Muhammed.....	41
Pal, Boza.....	133	Shahabi, Manochehr.....	140
Palá-Paúl, Jesús.....	149, 150	Shimada, Tsutomu.....	136
Palavra, A. M. F.....	31, 69	Shindo, Masaki.....	136
Pálfí, M.....	48	Shiroma, K.....	123
Palmeira, A.....	164, 176	Silva, Francisco F.....	86
Pank, Friedrich.....	146		

Silva, Magnólia A S.....	169, 170	von Konrat, M.	32
Silva, Milton Helio L.....	36, 46	Wąsowicz, Erwin.....	194
Silva, Ubiratan F. da.....	138, 171	Wawrzeńczyk, Czesław.....	190
Simándi, B.	48, 116	Weber, Andréia D.....	89
Simić, Mirjana.....	109	Weber, Jörg.....	167, 191
Simin, Nataša.....	58	Wenham, T.	192, 193
Simionatto, Euclésio.....	89, 138, 171, 172	Wilcock, Christopher C.	81
Slavkovska, Violeta.....	71, 109	Wolschann, Peter.....	61
Sokovic, Marina.....	184	Yeretzian, Chahan.....	60
Sonsuzer, Serap.....	163, 175	Yilmaz, Levent.....	163, 175
Stahl-Biskup, Elisabeth.....	20	Zaouali, Y.	57
Stefanini, Mirian B.	173, 174	Zawirska-Wojtasiak, Renata.....	194
Stefańska, J.	55	Zivkovic, Vladimir.....	75
Steinlesberger, Heidi.....	90	Zoghbi, Maria das Graças B.	36, 45, 46
Stilinović, Božidar.....	151, 152		
Strzelecka, H.....	55		
Subramoney, S.....	186		
Sumnu, S. Gulum.....	175		
Svoboda, K. P.	73		
Székely, Gabriella.....	144, 145		
Szmigiel-Pieczewska, Maia.....	190		
Szóke, É.....	116		
Szumny, Antoni.....	190		
Tabanca, N.....	54		
Tahara, Satoshi.....	78		
Tatsadjeu, Leopold.....	140		
Tavares, C.....	164, 176		
Tavares, Regina.....	141		
Taveira, Francisca Socorro N.	46, 121		
Teixeira, Generosa.....	155		
Teixeira, J. P. F.....	123, 124		
Telascrea, Marcelo.....	66		
Termentzik, K.....	177		
Then, M.	117		
Theodoro, Suzi Huff.....	178		
Tirillini, B.	122		
Tomassini, T. C. B.....	179		
Tomi, Félix.....	180, 181		
Tot, Andrea.....	133		
Toyota, M.	32, 147		
Trenkle, Robert W.....	28		
Trilles, Bénédicte.....	182		
Trivara, N.	177		
Tulok, M.	183		
Tzakou, Olga.....	71, 132, 184		
Unterweger, Heidrun.....	110		
Urieta, J. S.	31, 69		
van Niekerk, Léon.....	146		
van Staden, J.....	40		
van Vuuren, S. F.....	40, 77, 186, 187, 189		
van Zyl, R. L.....	40		
Vargas, Rubem M. F.....	188		
Vasconcellos, Vanessa R.....	123, 124		
Velasco-Negueruela, Arturo.....	94, 101, 149, 150		
Venâncio, F.....	70		
Vender, Carla.....	146		
Vicente, Ana.....	82		
Vilarem, G.	93		
Viljoen, A. M.....	40, 77, 186, 187, 189		
Vollmann, Carsten.....	112		

Participants List



Prof. Dr Aasen, Arne J.

University of Oslo
Dept of Pharmacy
P.O Box 1068
Oslo N-0316 Blindern
Norway
Phone :+4722854180 / Fax :+4722855947
ajaasen@farmasi.uio.no

Dr. Albert Lllana, Francisco José

Conselleria de Medi Ambient
Generalitat Valenciana
C/ Francesc Cubells, 7
Valencia 46011
Spain
Phone :+34963867692 / Fax :+34963863768
gabriel.ballester@cma.m400.gva.es

Dr. Andrade, Eloisia Helena A.

Museu Emílio Goeldi
Departamento de Botânica
CP 399, 66040-170 Belém,
Brazil
eloisa@museu-goeldi.br

Mrs. Antosik, Kamilla

Firmenich SA
Chrzanowska, 10
Grodzisk Mazowiecki 05-825
Poland
Phone : +48227241760 / Fax :+48227555972
KAMILLA.ANTOSIK@firmenich.com

Prof. Dr. Antunes, Teresa

Faculdade de Ciências
Dept de Biologia Vegetal
Edf C2 Campo Grande
Lisboa 1749-016
Portugal
Phone :+351217500000 / Fax :+351217500048
teresa.antunes@fc.ul.pt

Mr. Apati, Pal

Semmelweis University
Institute of Pharmacognosy
Ulloi Street 26
Budapest H-1085
Hungary
Phone :+3612660120/5306 / Fax :+3613172979
lembi@drog.sote.hu

Prof. Dr Asakawa, Yoshinori

Faculty of Pharmaceutical Sciences
Tokushima Bunri University
Yamashiro-cho, 770-8514
Tokushima 770-8514
Japan
Phone :+886229611 / Fax :+886558774
asakawa@ph.bunri-u.ac.jp

Dr. Awadalla, Amal

Medicinal & Aromatic Plants Research Institute
P.O Box 11496
Khartoum 11496
Sudan
Phone : +249011784987 / Fax : +249011773771
hsubki@hotmail.com

Dr. Ayoub, Nahla

Faculty of Pharmacy
Abbasia Square
Kairo
Egypt
Phone : +2024512550 / Fax :+2024512550
ayoub.n@link.com.eg

Dr. Bamasian, Shahrzad

Dept of Chemistry
Islamic Azad University
No. 45, 11th Street Asad Abady
Avenue Tehran 14336
Iran
Phone : +980218711960
abam@hamgam.com

Prof. Dr Barata, Lauro E.S

Instituto de Química
Universidade Estadual de Campinas
CP 6154
Campinas 13083-970
Brazil
Phone : +551932876822 / Fax :+551937883023
lbarata@iqm.unicamp.br

Mrs. Barreto, Rosária

Revista Activa
Largo da Lagoa, 15-C
Linda-a-Velha 2795-116
Portugal
IsabelV@acj.pt

Prof. Dr Barros, Neiva

Universidade de Caxias do Sul
R. Francisco Getúlio Vargas, 1130
Caxias do Sul 95070-560
Brazil
Phone : +55542182149 / Fax : +55542182149
n.barros@terra.com.br

Prof. Dr Barroso, José Manuel

Faculdade de Ciências de Lisboa
Dept de Biologia Vegetal
Piso 1, Bloco C2
Campo Grande
Lisboa 1749-016
Portugal
Phone : +351217500069 / Fax : +351217500048
jgbarroso@kernel.cc.fc.ul.pt

Prof. Dr Başer, K. Hüsni Can

Medicinal and Aromatic Plant and Drug Research
Centre
TBAM
Anadolu University
Eskisehir 26470
Turkey
Phone : +902223352952 / Fax : +902223350127
khebaser@anadolu.edu.tr

Mrs. Bauermann, Ulrike

IGV Institut für Getreideverarbeitung GmbH
Arthur-Scheunert-Allee, 40/41
Bergholz-Rehbrücke 14558
Germany
Phone : +493320089207 / Fax : +493320089251
u_bauermann@igv-gmbh.de

Mrs. Bazylko, Agnieszka

Dept of Pharmacognosy
Faculty of Pharmacy / University of Medicine
ul. Banacha, 1
Warsaw 02-097
Poland
oklyzab@farm.amwaw.edu.pl

Prof. Dr Ben Fadhel, Najeh

Institut National des Sciences Appliquées et de
Technologie
Centre Urbain Nord B.P n° 676
Tunis Cedex 1080
Tunisie
Phone : +21671703627 / Fax : +21671704329
najehbenfadhel@lycos.fr

Dr. Bettini, Mércia F. M.

Flavor Tec - Aromas de Frutas, Ltda
Av. Bela Vista, 971 Parque Industrial
Pindorama 15830-000
Brazil
Phone : +55175721000 / Fax : +55175211450
mercia@flavortec.com.br

Mr. Bianchi, Antonio

COE Traditional Medicine
Via Lazzaroni, 8
Milano 20124
Italy
Phone : +39035983017 / Fax : +390266714338
bantonio@globalnet.it

Prof. Dr Bicchi, Carlo

Dipartimento di Scienza e Tecnologia del Farmaco
University of Torino
Via Pietro Giuria, 9
Torino I-10125
Italy
Phone : +390116707662 / Fax : +390116707687
carlo.bicchi@unito.it

Dr. Bini, Claudio

Dept Biologia Vegetale
University of Florence
Via La Pira 4
Firenze 50121
Italy
Phone : +390552727396 / Fax : +390552757399
maleci@unifi.it

Mrs. Birkby, Jane

Botanix, Ltd
Hop Pocket Lane, Paddock Wood
Tonbridge, Kent
Tonbridge Kent TN9 6DQ
UK
Phone : +441842833415 / Fax : +441892836987
jane.birkby@botanix.co.uk

Prof. Dr Boussaid, Mohamed

Institut National des Sciences Appliquées et de
Technologie
Centre Urbain Nord
B.P n° 676
Tunis 1080
Tunisie
Phone : +21671703627 / Fax : +21671704329
mhdboussaid@rnu.insat.tn

Mrs. Božin, Biljana

Institute of Biology Faculty of Sciences
Trg Dositeja Obradovica, 3
Novi Sad 21000
Yugoslavia
Phone : +38121350122 / Fax : +3812154065
mimica@ih.ns.ac.yu

Mrs. Brandão, Maria das Graças Lins

Universidade Federal de Minas Gerais
Faculdade de Farmácia
Av. Olegário Maciel, 2360
Belo Horizonte 30180-112
Brazil
Phone : +553133397670 / Fax : +553133397663
branlins@dedalus.lcc.ufmg.br

Dr. Brevard, Hugues

Nestlé Research Center
P.O Box 44
Vers-chez-les-Blanc Lausanne 26
1000 Switzerland
Phone : +41217858409 / Fax : +41217858554
hugues.brevard@rdls.nestle.com

Dr. Brud, Władysław

Pollena Aroma, Ltd
ul. Klasyków, 10
Warszawa 03-115
Poland
Phone : +48228114270 / Fax : +48228119228
w.sbrud@pollenaaroma.com.pl

Prof. Dr Buchbauer, Gerhard

Institute of Pharmaceutical Chemistry
University of Vienna
Althanstrasse, 14
Vienna A-1090
Austria
Phone : +431427755150 / Fax : +43142779551
gerhard.buchbauer@univie.ac.at

Mr. Cadariu, Traian

" Bruder Unterweger " Oils
Thal-Aue, 13
Thal-Assling 9911
Austria
Phone : +43485582010 / Fax : +434855820122
labor.bu-oils@tirol.com

Dr. Cardoso, Beatriz

Faculdade de Ciências
Universidade de Lisboa
Parada do Alto S. João, nº 12 - 6º Frt
Lisboa 1900-052
Portugal Phone : +35121818368
bmac@fc.ul.pt

Dr. Carneiro, Ana Paula

Apartado 41007
Lisboa 1506-001
Portugal
Phone : +351219594719

Prof. Dr Cassel, Eduardo

Faculdade de Química - PUCRS
Av Ipiranga, 6681
Bairro Partenon
Porto Alegre 90619-900
Brazil
Phone : +555133203653 / Fax : +555133203625
cassel@puccrs.br

Dr. Castro, Dulce

Faculdade de Ciências Agronomicas de Botucatu
UNESP-Universidade Estadual Paulista
R.Alfredo di Lello, 10
São Manuel 18650-000
Brazil
Phone : +5501468411642
dulcem@fca.unesp.br

Prof. Dr Cavaleiro, Carlos

Faculty of Pharmacy
University of Coimbra
Rua do Norte
Coimbra 3004-534
Portugal
Phone : +351239859995 / Fax : +351239827126
cavaleir@ff.uc.pt

Dr. Cavaleiro, Alberto José

Instituto de Química
UNESP-Universidade Estadual Paulista
Rua Prof. Francisco Degni s/n
Araraquara-SP 14800-900
Brazil
Phone : +550162016667 / Fax : +550162227932
albjcava@iq.unesp.br

Dr. Chatzopoulou, Paschalina

National Agriculture Research Foundation
Dept of Aromatic and Medicinal Plants
P.O Box 60458
Thermi
Thessaloniki 57001
Greece
Phone : +30310471110 / Fax : +30310347418
xatzlin@yahoo.gr

Dr. Chung, Hae Gon

National Crop Experiment Station
209 Seodun-dong Kwansunku
Suweon 441-707
Republic of Korea
Phone : +82312906729 / Fax : +82312906787
haegon@rda.go.kr

Dr. Claussen, Susanne

Aromatechnics, Lda
Rua do Monte Grande
S. Bartolomeu do Sul
Castro Marim 8950-270
Portugal
Phone : +351281957560 / Fax : +351281957560
aromatec@oninet.pt

Prof. Dr Coelho, José

ISEL
R.Conselheiro Emidio Navarro, 1
Lisboa 1949-014
Portugal
Phone : +351218317066 / Fax : +351218317267
jcoelho@deq.isel.pt

Prof. Dr Costa, Maria do Céu

INETI
Estrada do Paço do Lumiar, Ed. F
Lisboa 1649-038
Portugal
Phone : +351217164211 / Fax : +351217168100
ceu.costa@mail.ineti.pt

Dr. Couladis, Maria

University of Athens
Dept of Pharmacognosy and Chemistry of Natural
Products / School of Pharmacy
Panepistimiopolis Zografou
Athens 15771
Greece
Phone : +30107274585 / Fax : +30107274591
kouladi@pharm.uoa.gr

Ms. Cozzani, Stéphanie

Université de Corse
Equipe Chimie des Produits Naturels UMR
CNRS 6134 - Quartier Grossetti - 20250 Corte
France

Ms. Cross, Deborah

Scottish Agricultural College
Avian Science Research Centre
Auchincruive Estate
Ayr KA6 5HW
Scotland UK
Phone : +4401292525117 / Fax : +4401292525098
D.Cross@au.sac.ac.uk

Mr. Cunha Luís, Tiago

Faculdade de Ciências de Lisboa
R. Augusto Costa(Costinha), nº 10 - 3º Dto.
Lisboa 1500-064
Portugal
Phone : +351217603916
tiago.c.luis@netcabo.pt

Mr. Currais, António José

Faculdade de Ciências de Lisboa
Praça Rainha Santa, nº 4 - 6º Dto.
Lisboa 1600-687
Portugal
Phone : +351217590076 / Fax : +351217590257
vickie@clix.pt

Prof. Dr Damjanovic, Biljana

Faculty of Metallurgy and Technology
Cetinjski bb.
Podgorica 81000
Montenegro
Yugoslavia
Phone : +38181245406 / Fax : +38181242301
bibana@cg.yu

Prof. Dr Davis, Edward

Institute of Biological Chemistry
Washington State University
P.O Box 646340
Pullman, WA 99164-6340
USA
Phone : +15093344117 / Fax : +15093357643
edd@mail.wsu.edu

Dr. Demirci, Betül

Anadolu University
TBAM
Eskisehir 26470
Turkey
Phone : +902223350580 / Fax : +902223350127
bdemirca@anadolu.edu.tr

Dr. Demirci, Fatih

Anadolu University
TBAM
Eskisehir 26470
Turkey
Phone : +902223350580 / Fax : +902223350127
fdemirci@anadolu.edu.tr

Ms. Duarte, Carla Sofia

Universidade do Algarve
Rua do Sobral, nº 2 Vale de Figueira
Santarém 2000-740
Portugal
Phone : +351964066120
cduarte@ureach.com

Dr. El Kamali, Hatil

Medicinal & Aromatic Plants Research Institute
P.O Box 11496
Khartoum 11496
Sudan
Phone : +249011784987 / Fax : +249011773771
htlkamali@yahoo.com

Dr. Fernandes, Francisco Manuel

Jardim Botânico da Madeira
Quinta do Bom Sucesso
Funchal 9050
Portugal
Phone : +351291211200 / Fax : +351291211206
francisco.fernandes.sra@gov-madeira.pt

Prof. Dr Fernandes Ferreira, Manuel

Universidade do Minho
Departamento de Biologia
Braga 4710-057
Portugal
Phone : +351253604310 / Fax : +351253678980
mfferreira@bio.uminho.pt

Ms. Ferreira, Cláudia

Faculdade de Ciências / Universidade de Lisboa
Dept de Biologia Vegetal
Piso 1, Bloco C2
Campo Grande
Lisboa 1749-016
Portugal
Phone : +351217500000 / Fax : +351217500048
c.ferreira@iol.pt

Ms. Ferreira, Léopoldine

5 Bis, Rue du Bas Chatron
Saint Germain de la Grange
78640 France
Phone : +33662697368
leopoldferreira@aol.com

Mr. Ferreira, Nicolau José
Faculdade de Ciências de Lisboa
R.da Paz, Lote 19-2º Esq.
S.Domingos de Rana 2785-500
Portugal
Phone : +351214455695
nicolau.ferreira@mail.pt

Prof. Dr Figueiredo, Ana Cristina
Faculdade de Ciências de Lisboa
Dept Biologia Vegetal
Piso 1, Bloco C2
Campo Grande
Lisboa 1749-016
Portugal
Phone : +351217500069 / Fax : +351217500048
acsf@fc.ul.pt

Ms. Francisco, Ana Margarida Costa
Faculdade de Ciências de Lisboa
Quinta de Santiago Cabeço de Mouro
Portalegre
7300-011 Portalegre
Phone : +351245207162
Portugal
anafrancisco80@hotmail.com

Prof. Dr Francke, Wittko
University of Hamburg
Institut für Organische Chemie
Martin-Luther-King-Platz, 6
Hamburg D-20146
Germany
Phone : +4940428382866 / Fax : +4940428383834
francke@chemie.uni-hamburg.de

Dr. Franz, Chlodwig
Institute for Applied Botany
Veterinarplatz, 1
Vienna 1210
Austria
Phone : +431250773101 / Fax : +431250773190
chlodwig.franz@vu-wien.ac.at

Dr. Frizzo, Caren
Aripê Citrus Ltda
RS124, Km 1.2
Montenegro 95780-000
Brazil
Phone : +55516321444 / Fax : +55516323174
aripe@zaz.com.br; caren.aripe@terra.com.br

Mrs. Fumagalli, Valeria
Moellhausen, S.P.A
Via A Ponchielli, 13
Cologno Monzese (MI) 20093
Italy
Phone : +390227301917 / fax : +390227302743
info@moellhausen.com

Ms. Furusawa, Mai
Tokushima Bunri University
Yamashiro-cho, 180
Tokushima 771-8514
Japan
Phone : +886229611 / Fax : +886223217
s010401@tokushima.bunri-u.ac.jp

Dr. Galvão, Helena
Universidade do Algarve
FCMA
Gambelas
Faro 8000-117
Portugal
Phone : +351289800900 / Fax : +351289818353
hgalvao@ualg.pt

Ms. Garbo, Emiliana
Martini-Bacardi
Piazza L. Rossi, 2
Pessione 10020
Italy
Phone : +390119419270 / Fax : +390119419324
egarbo@bacardi.com

Dr. Gehrman, Beatrice
Institute of Pharmacy
Humboldt-University of Berlin
Goethestrasse, 54
Berlin D-13086
Germany
Phone : +493096592321 / Fax : +49309248280
beatrice.gehrmann@rz.hu-berlin.de

Ms. Gomes, Paula
LSRE
Faculty of Engineering of University of Porto
R. Dr. Roberto Frias, Edif. E - Sala E403
Porto 4200-465
Portugal
Phone : +351225081669 / Fax : +351225081674
pgomes@fe.up.pt

Dr. Gonzaga, Wellington de Abreu
Universidade Federal de Santa Maria
Campus Universitário
Santa Maria 97105-900
Brazil
Phone : +5522082058869
welloeiras@bol.com.br

Dr. Grassi, Paolo
Institute for Applied Botany
University of Veterinary Medicine
Veterinarplatz, 1
Vienna 1200
Austria
Phone : +431250773101 / Fax : +431250773190
trucho3000@mixmail.com

Dr. Habibi, Zohreh

Dept of Chemistry
Shahid Beheshti University
Evin
Tehran
Iran
Phone : +98212720739 / Fax : 98212403041
zohre1340@hotmail.com

Dr. Harlalka, Ramakant

Nishant Aromas
424, Milan Ind. Estate, Off T.J Road
Cotton Green
Mumbai 400033
India
Phone : +91224711431 / Fax : +91224716502
nishantaromas@vsnl.com

Mr. Hernandez-Ochoa, Leon

ENSIACET (LCA-CATAR)
118, Route de Narbonne
Toulouse Cedex 4 31077
France +33562885724
Phone : +33562885730
leonzempoala@yahoo.com

Dr. Heuberger, Eva

Institute of Pharmaceutical Chemistry
University of Vienna
Althanstrasse, 14
Vienna A-1090
Austria
Phone : +431427755160 / Fax : +43142779551
Eva.Heuberger@univie.ac.at

Prof. Dr Hiltunen, Raimo

Dept of Pharmacy
University of Helsinki
P.O Box 56
Helsinki 00014
Finland
Phone : +358919159140 / Fax : +358919159138
raimo.hiltunen@helsinki.fi

Dr. Hongratanaworakit, Tapanee

Faculty of Pharmaceutical Sciences
Srinakharinwirot University
Rangsit-Ongkharak Road, 63
Nakorn-Nayok 26120
Thailand
Phone : +66373950945 / Fax : +6637395394
htapanee@hotmail.com

Mr. Hudewenz, Volker

Drom Fragrances International KG
Oberdiller Str. 18
Baierbrunn D-82065
Germany
Phone : +4989744250 / Fax : +49897934966
hud@drom.com

Ms. Iñigo, Ana

Facultad de Biología
Universidad Complutense Madrid
Av. Complutense s/n
Madrid 28040
Spain
Phone : +34913944433 / Fax : +34913945034
ainigo@terra.es

Ms. Ilori, Olukemi

University of Lagos
Department of Pharmacognosy / School of Pharmacy
P.M.B 12003
Idi-Araba, Lagos
Nigeria
Phone : + 23414803672 / Fax : + 23415851432
olukemiilori@yahoo.com

Mr. İşcan, Gökcalp

Medicinal and Aromatic Plant and Drug Research
Centre
Anadolu University
Eskisehir 26470
Turkey
Phone : +902223352952 / Fax : +902223350127
giscan@anadolu.edu.tr

Dr. Ivanova, Alexandra

Komarov Botanical Institute
Prof. Popov Str. 2
St.Petersburg 197376
Russia
Phone +78123464477 / Fax : +78122344512
Alyx@av-online.ru

Dr. Jakobsen, Henrik Byrial

Videvadgaard
Blaesborgvej, 9
Leyre 4320
Denmark
Phone : +4546480003
henrik.byrial@get2net.dk

Dr. Javidnia, Katayoun

Faculty of Pharmacy
Shiraz University of Medical Sciences
P.O Box 71345-1149
Shiraz 71345
Iran
Phone : +987112294849 / Fax : +987112290091
kjavidnia@albarta.com

Dr. Jirovetz, Leopold

Institute of Pharmaceutical Chemistry
University of Vienna Althanstrasse, 14
Vienna A-1090
Austria
Phone : +431427755091 / Fax : +43142779551
leopold.jirovetz@univie.ac.at

Dr. Joulain, Daniel

ROBERTET, S.A
37, Avenue Sidi-Brahim
B.P 52100
Grasse 06131
France
Phone : +33493403309 / Fax : +33493706809
daniel.joulain@robertet.fr

Prof. Dr Kaiser, Roman

Givaudan Dubendorf Ltd
Ueberlandstrasse, 138
Dubendorf 8600
Switzerland
Phone : +4118242355 / Fax : +4118242926
roman.kaiser@givaudan.com

Dr. Kalembe, Danuta

Technical University
Inst Food General Chemistry
ul. Stefanowskiego 4/10
Lodz 90-924
Poland
Phone : +48426313423 / Fax : +48426362860
dakal@snack.p.lodz.pl

Dr. Kalita, Mohan Chandra

Gauhati University
Dept of Biotechnology
Gauhati University
Guwahati 781014
Assam
India
Phone : +910361572408 / Fax +910361570133
mckalital@sancharnet.in

Dr. Kardali, Mohamed

Departamento de Biología Vegetal I
Facultad de Biología
Universidad Complutense Madrid
Madrid 28040
Spain
Phone : +34650336566 / Fax : +34913945034
kardali@bio.ucm.es

Prof. Dr Katsiotis, Stavros

National Agriculture Research Foundation
Dept of Aromatic and Medicinal Plants
P.O Box 60458
Thermi
Thessaloniki 57001
Greece
Phone : +3031471110 / Fax : +3031347418
stakat@auth.gr

Dr. Kawahigashi, Tatsuo

Dept of Applied Chemistry, Science & Technology
Kinki University
3-4-1, Kowakae
Higashi
Osaka 577-8502
Japan
Phone : +81667212332 / Fax : +81667305896
L1kawahi@cced.kindai.ac.jp

Dr. Kaya, Ayla

Medicinal and Aromatic Plant and Drug Research
Centre
Anadolu University
Eskisehir 26470
Turkey
Phone : +902223350581 / Fax : +902223350127
aykaya@anadolu.edu.tr

Dr. Khalid, Hassan S.

Medicinal and Aromatic Plants Research Institute
P.O Box 11496
Khartoum
Sudan
Phone : +24911784987 / Fax : +24911773771
hskhalidmo@yahoo.com

Dr. Kim, Seong Min

College of Industrial Science
Kongju National University
527 Yesangun
Chungnam 340-800
Republic of Korea
Phone : +824583322481 / Fax : +824583322485
smkim@knu.kongju.ac.kr

Prof. Dr Kırimer, Neşe

Medicinal and Aromatic Plant and Drug Research
Centre
Anadolu University
Eskisehir 26470
Turkey
Phone : +902223352952 / Fax : +902223350127
nkirimer@anadolu.edu.tr

Dr. Kolalite, Milana

Komarov Botanical Institute
Russian Academy of Sciences-Plant Resources Dept.
Prof. Popov Str. 2
Saint-Petersburg 197376
Russia
Phone : +78123464477 / Fax : +78122344512
milana_kolalite@mail.ru

Prof. Dr Kovacevic, Nada

Faculty of Pharmacy
Vojvode Stepe, 450
Belgrade 11000
Yugoslavia
Phone : +381113970379 / Fax : +381113972840
nadak@pharmacy.bg.ac.yu

Mrs. Krist, Sabine

Institute of Pharmaceutical Chemistry
Althanstrasse, 14
Vienna 1090
Austria
Phone : +43223629883

Dr. Krüger, Hans

Bundesanstalt fuer Zuechtungsforschung an
Kulturpflanzen
Neuer Weg, 22-23
Quedlinburg 06484
Germany
Phone : +490394647282 / Fax : +490394647234
h.krueger@bafz.de

Prof. Dr Kubeczka, Karl-Heinz

Untere Steigstrasse, 12b
Margetshochheim D-97276
Germany
Phone : +499314676210 / Fax : +499314676211
kubeczka@t-online.de

Dr. Kürkçüoğlu, Mine

Medicinal and Aromatic Plant and Drug Research
Centre
Anadolu University
Eskisehir 26470
Turkey
Phone : +902223352952 / Fax : +902223350127
mkurkcuo@anadolu.edu.tr

Prof. Dr Kustrak, Danica

Faculty of Pharmacy and Biochemistry
Ante Kovacica, 1
P.O Box 156
Zagreb 10000
Croatia
Phone : +38514818288 / Fax : +38514856201
dekanat@nana.pharma.hr

Dr. Lassak, Erich

Phytochemical Services
254, Quarter Sessions Road
Westleigh 2120
Australia
Phone : +61298751894 / Fax : +61298751791

Dr. Lawrence, Brian

Journal of Essential Oil Research
110 Staffordshire Court
Winston Salem NC-27104
USA
Phone : +13367412031 / Fax : +13367416343

Prof. Dr Lemberkovics, Éva

Semmelweis University
Institute of Pharmacognosy
Ulloi Street 26
Budapest H-1085
Hungary
Phone : +3612660120 / Fax : +3613172979
lembi@drog.sote.hu

Mr. Liddle, Peter

Bacardi Martini
B.P 50
Saint-Ouen Cedex 93401
France
Phone : +33149454873 / Fax : +33149454905
peliddle@bacardi.com

Dr. Liechti, Christoph

Givaudan Dubendorf Ltd
Ueberlandstrasse, 138
Dubendorf 8600
Switzerland
Phone : +4118242365 / Fax : +4118242976
christoph.liechti@givaudan.com

Dr. Linssen, Victoria

Soul of the Plant
5024 E. Marino Drive
Scottsdale, AZ 85254
USA
Phone : +16029713333 / Fax : +16029713333
vlinssen@hotmail.com

Dr. Lis-Balchin, Maria

South Bankl University
Borough Road
London SE1 OAA
UK
Phone : +442089428242 / Fax : +442089428242
lisbalmt@sbu.ac.uk

Dr. Lochynski, Stanislaw

Institute of Organic Chemistry, Biochemistry and
Biotechnology
Wroclaw University of Technology
Wybrzeze Wyspianskiego, 27
Wroclaw 50-370
Poland
Phone : +48713202400 / Fax : +48713284064
lochynski@kchf.ch.pwr.wroc.pl

Dr. Lopes, Lúcia

Centro de Desenvolvimento Sustentável
SAS Quadra 05 Bloco H - Sala 204
Brasilia-DF 70070-914
Brazil
Phone : +55613215001 / Fax : +55613228473
lulopes7@terra.com.br

Prof. Dr Maia, José Guilherme

Museu Emílio Goeldi
Av. Magalhães Barata, 376
Belém 66040-170
Brazil
Phone : +5591274425 / Fax : +55912744025
gmaia@museu-goeldi.br

Prof. Dr Maleci, Laura

Dept Biologia Vegetale
University of Florence
Via La Pira 4
Firenze 50121
Italy
Phone : +390552727396 / Fax : +390552757399
maleci@unifi.it

Dr. Marques, Márcia Ortiz Mayo

Instituto Agronomico de Campinas
Av. Barão de Itapura, 1481
CP 28
Campinas 13001-970
Brazil
Phone : +551932415188 / Fax : +551932415188
mortiz@cec.iac.br

Dr. Masoudi, Abdolnasser

KAF Joint Stock Company (DARUGAR)
3, West Armaghan Str, Opposite Park Mellat
Valiasr Ave.
Tehran 19678
Iran
Phone : +98212058784 / Fax : +98212058785
nasser_massoudi@yahoo.com

Prof. Dr Masoudi, Shiva

Depart of Chemistry
Islamic Azad University / Central Tehran Branch
8, 10 meter Golestan Street
Shahid Mahmoud Abadi Ave
Nabard Ave, Pirouzi Ave 1766637551
Tehran
Iran
Phone : +9821727995
shmasoudi@yahoo.com

Dr. Mateus, Eduardo

UNL-FCT/DCEA
Quinta da Torre
Costa da Caparica 2825-114
Portugal +351212948300
Phone : +351212948554
abr@mail.fct.unl.pt

Prof. Dr Medeiros, Jorge

Universidade dos Açores
Rua Mãe de Deus
Ponta Delgada
Açores / Portugal
Phone : +351296650175 / Fax : +351296650171
jrmediros@notes.uac.pt

Ms. Meireles de Sousa, Inês

Faculdade de Ciências de Lisboa
R. Capitães de Abril, nº 9 - 5º Esq.
Amadora 2700-048
Portugal
Phone : +351214743775
Ineq@netcabo.pt

Prof. Dr Menut, Chantal

Laboratoire de Chimie Biomoléculaire
ENSCM
8, Rue de l'École Normale
Montpellier 34296
France
Phone : +33467144340 / Fax : +33467144340
cmenut@univ-montp2.fr

Prof. Dr Miguel, Graça

Universidade do Algarve
Campus de Gambelas
Faro 8000-117
Portugal
Phone : +351289800957 / Fax : +351289818419
mgmiguel@ualg.pt

Mrs. Mihajlovic, Biserka

Institute of Public Health
Medical Faculty
Futoshi put, 103
Novi Sad 21000
Yugoslavia
mimica@ih.ns.ac.yu

Prof. Dr Mimica-Dukic, Neda

Institute of Chemistry
Faculty of Sciences
Trg Dositeja Obradovica, 3
Novi Sad 21000
Yugoslavia
Phone : +38121350122 / Fax : +3812154065
dukaned@eunet.yu

Mr. Ming Chau, Lin

Faculdade de Ciências Agronômicas de Botucatu
FCA/Unesp
Fazenda Experimental Lageado s/n
Caixa Postal 237
Botucatu/SP 18603-970
Brazil
Phone : +551468411642
dulcem@fca.unesp.br

Dr. Miri, Ramin

Faculty of Pharmacy
Shiraz University of Medical Sciences
P.O Box 71345-1149
Shiraz 71345
Iran
Phone : +987112294849 / Fax : +987112290091
mirir@sums.ac.ir

Prof. Dr Miyazawa, Mitsuo

Dept of Applied Chemistry, Science & Technology,
Kinki University
3-4-1, Kowakae
Higashi
Osaka 577-8502
Japan
Phone : +81667212332 / Fax : +81667274301
miyazawa@apch.kindai.ac.jp

Mr. Moellhausen, Luca

Moellhausen, S.P.A
Via A Ponchielli, 13
Cologno Monzese (MI) 20093
Italy
Phone : +390227301917 / Fax : +390227302743
info@moellhausen.com

Dr. Moldão Martins, Margarida

Instituto Superior de Agronomia
DAIAT/SCTA
Tapada da Ajuda
Lisboa 1349-017
Portugal
Phone : +351213653547 / Fax : +351213632000
mmoldao@isa.utl.pt

Dr. Mookherjee, Braja D.

International Flavors & Fragrances, Inc.
1515 US Highway 36
Union Beach 07735
USA
Phone : +17323354500 / Fax : +17323352591
braja.mookherjee@iff.com

Prof. Dr Morel, Ademir Farias

Universidade Federal de Santa Maria
Campus Universitário
Santa Maria 97105-900
Brazil
Phone : +5522082058869
afmorel@hanoi.base.ufsm.br

Ms. Mota, Luísa

Faculdade de Ciências / Universidade de Lisboa
Dept de Biologia Vegetal
Piso 1, Bloco C2
Campo Grande
Lisboa 1749-016
Portugal
Phone : +351217500000 / Fax : +351217500048
luisa_branquinho@hotmail.com

Dr. Najda, Ali

Consultor S.A
7, Rue du Mont-Blanc
P.O Box 1042
Geneva 1211
Switzerland
Phone : +41227383440 / Fax : +41227383472
ali.najda@consultor.ch

Dr. Nendel, Maja

Frey & Lau GmbH
Immenhacken, 12
Henstedt-Ulzburg 22548
Germany
Phone : +4941939953 / Fax : +494193995580
mnendel@freylau.de

Mr Neves, António Jorge Franca

Portugal

Prof. Dr Neves, Marta Lopes

ESB-UCP
R. Mestre Mateus Fernandes
Caldas da Rainha 2500-237
Portugal
Phone : +351262839338 / Fax : +351262839339
marta@esb.ucp.pt

Eng^a Neves, Susana

Escola Superior Agrária de Ponte de Lima
Trav. Antero de Quental, 6 r/c Esq.
Amadora 2700-061
Portugal
Phone : +351214942075
susananeves@esapl.pt

Prof. Dr Ngassoum, Martin Benoit

University of Ngaoundere
ENSAI-IUT, Dept of Applied Chemistry
B.P 455
Ngaoundere
Cameroon
Phone : +237252751 / Fax : +237252751
ngassoum@caramail.com

Mrs. Niesluchowska, Alicja

Firmenich SA
Chrzanowska, 10
Grodzisk Mazowiecki 05-825
Poland
Phone : +48227241760 / Fax : +48227555972
alicja.niesluchowska@firmenich.com

Mrs. Niestroj, Eva

Givaudan Deutschland GmbH
Giselherstrasse, 11
Dortmund 44319
Germany
Phone : +492312186420 / Fax : +4902312186266
eva.niestroj@givaudan.com

Dr. Nogueira, Maria Teresa

INETI
Estrada do Paço do Lumiar, Ed. F
Lisboa 1649-038
Portugal
Phone : +351217165141 / Fax : +351217168100
teresa.nogueira@mail.ineti.pt

Prof. Dr Noma, Yoshiaki

Tokushima Bunri University
Yamashiro-cho, 180
Tokushima 771-8514
Japan
Phone : +886229611 / Fax : +886223217
ynoma@tokushima.bunri-u.ac.jp

Mr. Nour, Amal Mukhat M

Faculty of Pharmacy
P.O Box 10777
Khartoum
Sudan
Phone : +24912214050
amal_mukhtar@hotmail.com

Dr. Novák, Ildikó

Szent István University
Dept of Medicinal and Aromatic Plants
29, Villány Str.
Budapest H-1118
Hungary
Phone : +3613726250 / Fax : +3613726330
inovak@omega.kee.hu

Dr. Novak, Johannes

Institute for Applied Botany
University of Veterinary Medicine
Veterinarplatz, 1
Vienna A-1210
Austria
Phone : +431250773104 / Fa x : +431250773190
Johannes.Novak@vu-wien.ac.at

Dr. Odukoya, Olukemi Abiodun

University of Lagos Pharmacognosy
Dept / School of Pharmacy / College of Medicine
Lagos PMB 12003
Idiaraba
Nigeria
Phone : +23408023144152 / Fax : +23415851432
olukemiodukoya@yahoo.com

Ms. Omatsu, Ikuko

Faculty of Pharmaceutical Sciences
Tokushima Bunri University
Yamashiro-cho, 180
Tokushima 770-8514
Japan
Phone : +886229611 / Fax : +886558174
phiku_usi@ph.bunri-u.ac.jp

Mr. Ormancey, Xavier

SCP
BP 11 - 855 Av. Maurice Donat
Mougins Cedex F-06252
France
Phone : +33492282050 / Fax : +33492282070
xavier.ormancey@biolandes.com

Dr. Ozek, Temel

Medicinal and Aromatic Plant and Drug Research
Centre
Anadolu University
Eskisehir 26470
Turkey
Phone : +902223352952 / Fax : +902223350127
tozek@anadolu.edu.tr

Mr. Palá-Paúl, Jesús

Depto Biología Vegetal I Botánica / Facultad de
Biología
Universidad Complutense Madrid
Madrid 28040
Spain
Phone : +34913944433 / Fax : +343945034
quibey@bio.ucm.es

Prof. Dr Palavra, António

Instituto Superior Técnico
Av. Rovisco Pais
Lisboa 1049-001
Portugal
Phone : +351218419387 / Fax : +351218464455
amgpalavra@popsrv.ist.utl.pt

Dr. Panero, Ombretta

Tradall Production Sarl (Bacardi-Martini Group)
267, Route de Meyrin
Meyrin 1217
Switzerland
Phone : +410227193475 / Fax : +410227193498
opanero@bacardi.com

Dr. Partl, Anamarija

Hercegovačka, 27
Zagreb HR-10000
Croatia
Phone : +38514683051 / Fax : +38514622895
apartl@croatica.botanic.hr

Mrs. Patel, Subha

International Flavors & Fragrances, Inc.
1515 US Highway 36
Union Beach 07735
USA
Phone : +17323354500 / Fax : +17323352591
subha.patel@iff.com

Prof. Dr Pedro, Luís Manuel

Faculdade de Ciências de Lisboa
Dept de Biologia Vegetal
Piso 1, Bloco C2
Campo Grande
Lisboa 1749-016
Portugal
Phone : +351217500069 / Fax : +351217500048
luis.pedro@fc.ul.pt

Ms. Pereira, Ana Luísa

Departamento de Biologia Vegetal - FCUL
Ed. C2 - Piso 1
Campo Grande
Lisboa 1749-016
Portugal
Phone : +351217500000 / Fax : +351217500048
ana.pereira@fc.ul.pt

Ms. Pereira, Ana Patricia

Instituto Superior Técnico
Av. Rovisco Pais
Lisboa 1049-001
Portugal
Phone : +351218419387 / Fax : +351218464455
apapereira@clix.pt

Dr. Piggott, John

University of Strathclyde
Dept of Bioscience
204 George Street
Glasgow G1 1XW
UK
Phone : +441415482150 / Fax : +441415534124
j.r.piggott@strath.ac.uk

Dr. Pluhár, Zsuzsanna

Dept of Medicinal and Aromatic Plants
Szent István University
29 Villányi Str.
Budapest H-1118
Hungary
Phone : +3613726250 / Fax : +3613726330
zpluhar@omega.kee.hu

Mr Prata, Pedro Miguel Cardoso

Portugal

Prof. Dr Proença da Cunha, António

Faculdade de Farmácia
Laboratório de Farmacognosia
Rua do Norte
Coimbra 3000
Portugal
Phone : +351239859995
pdacunha@ci.uc.pt

Mr. Radoias, Georges

" Bruder Unterweger " Oils
Thal-Aue, 13
Thal-Assling 9911
Austria
Phone : +43485582010 / Fax : +434855820122
labor.bu-oils@tirol.com

Dr. Radonic, Ani

Faculty of Chemical Technology
N. Tesle 10/V
Split 21000
Croatia
Phone : +38521385633 / Fax : +38521384964
radonic@ktf-split.hr

Dr. Raharivelomanana, Phila

Université de la Polynésie Française
BP 6570 FAAA
FAAA / Tahiti 98702
French Polynesia
Phone : +689803822 / Fax : +689439390
raharive@upf.pf

Dr. Rebelo de Almeida, M^a Margarida

Farmácia RioMouro
R.Óscar Monteiro Torres, 6
Rio de Mouro 2635-383
Portugal
Phone : +351219169200 / Fax : +351219171857

Mr. Regetzki, Soren

European Frutarom Corp.
Belle Alliance Str., 58
Hamburg D-20259
Germany
Phone : +494043257627 / Fax : +494043257650
soerenre@hotmail.com

Mr. Ribeiro, Carlos

Instituto Superior de Agronomia
DAIAT/SCTA
Tapada da Ajuda
Lisboa 1349-017
Portugal
Phone : +351914546100
c2mr@mail.pt

Ms. Richter, Rita

Universitaet Hamburg
Organische Chemie
Martin-Luther-King-Platz, 6
Hamburg D-20246
Germany
Phone : +49408511714 / Fax : +494088302257
ririchter@gmx.de

Dr. Rohloff, Jens

The Plant Biocentre Dept of Botany
Dragvoll
Trondheim N-7491
Norway
Phone : +4773590174 / Fax : +4773590177
jens.rohloff@chembio.ntnu.no

Dr. Romano, Anabela

Universidade do Algarve (FERN)
Campus de Gambelas
Faro 8000-117
Portugal
Phone : +351289800910 / Fax : +351289818419
aromano@ualg.pt

Dr. Roque, Odete

Faculdade de Farmácia
Laboratório de Farmacognosia
Rua do Norte
Coimbra 3000
Portugal
Phone : +351239859995
pdacunha@ci.uc.pt

Prof. Dr Rubiolo, Patrizia

Dipartimento di Scienza e Tecnologia del Farmaco
University of Turin
Via Pietro Giuria, 9
Turin 10125
Italy
Phone : +39116707661 / Fax : + 39116707687
patrizia.rubiolo@unito.it

Prof. Dr Rustaiyan, Abdolhossein

Dept of Chemistry, Science and Research
Campus I.A University
P.O Box 14515-775
Tehran
Iran
Phone : +98212716370
rustaiyan@excite.com

Dr. Sahin, Serpil

Middle East Technical University
Food Engineering Dept.
Ankara 06531
Turkey
Phone : +903122105627 / Fax : +903122101270
serp@metu.edu.tr

Prof. Dr Salgueiro, Lígia

Universidade de Coimbra - Faculdade de Farmácia
Laboratório de Farmacognosia
Rua do Norte
Coimbra 3000
Portugal
Phone : +351239859995 / Fax : +351239827126
ligia@ff.uc.pt

Prof. Dr Salvaterra-Garcia, Maria

Firmenich SA
Patent & Trademark Dept
P.O Box 239
Geneva 8 1211
Switzerland
Phone : +41227803341 / Fax : +41227803338
maria.garcia@firmenich.com

Mr. Santos, Pedro

ISA
R. António de Abreu, 126
Cascais 2750
Portugal
Phone : +351214841758
pedro.santos-926@clix.pt

Dr. Santos, Pedro A.

Centro de Biotecnologia Vegetal
Faculdade de Ciências / Universidade de Lisboa
Piso 1, Bloco C2
Campo Grande
Lisboa 1749-016
Portugal
Phone : +351217500000 / Fax : +351217500048
psantos@fc.ul.pt

Mr. Sarmento, David

IBET
Av. João XXI, 62-1º Dto.
Lisboa 1000
Portugal
Phone : +351917852602
David_sarmento@netcabo.pt

Mr. Scazzola, Franco

Moellhausen, S.P.A
Via A Ponchielli, 13
Cologno Monzese (MI) 20093
Italy
Phone : +390227301917 / Fax : +390227302743
info@moellhausen.com

Dr. Schäfer, Silvia

Karl-Winnacker-Institute, Dechema e.V.
Theodor-Heuss-Allee, 25
Frankfurt am Main 60486
Germany
Phone : +49697564347 / Fax : +49697564388
silvia.schaefer@dechema.de

Prof. Dr Scheffer, Johannes

Div. Pharmacognosy, LACDR
Leiden University
Gorlaeus Labs.
P.O Box 9502
Leiden 2300 RA
The Netherlands
Phone : +31715274474 / Fax : +31715274511
scheffer@chem.leidenuniv.nl

Dr Scheffer-Heeringa, Ella

Div. Pharmacognosy, LACDR
Leiden University
Gorlaeus Labs.
P.O Box 9502
Leiden 2300 RA
The Netherlands
Phone : +31715274474 / Fax : +31715274511
scheffer@chem.leidenuniv.nl

Dr. Schmidt, Erich

Kurt Kitzing GmbH
Hinterm Alten Schloss, 21
Wallerstein 86757
Germany
Phone : +499081275080 / Fax : +49908179200
info@artandfragrance.de

Ms. Schoenefeld, Karina

University of Applied Science Jena
Dept of Medical Engineering
Carl-Zeiss-Promenade, 2
Jena 07745
Germany
Phone : +493641205623 / Fax : +493641205601
Karina_Schoenefeld@web.de

Prof. Dr Schulz, Hartwig

BAZ, Institute for Plant Analysis
Neuer Weg, 22-23
Quedlinburg D-06484
Germany
h.schulz@bafz.de

Mrs. Sedaghat, Soheila

Islamic Azad University
North Tehran Branch
P.O Box 15815-3575
Tehran 14147-53313
Iran
Phone : +89212222667 / Fax : +89212949650
soheila_67@hotmail.com

Mr. Smith, Shamusideen A

University of Lagos
Department of Pharmacognosy / School of Pharmacy
P.M.B 12003
Lagos
Nigeria
Phone : + 2348023390877
sagboye@yahoo.com

Dr. Sevinate-Pinto, Isabel

Faculdade de Ciências
Dept de Biologia Vegetal
Edifício C2
Campo Grande
Lisboa 1749-016
Portugal
Phone : +351217500000 / Fax : +351217500048
teresa.antunes@fc.ul.pt

Eng^a Agr^a Silva, Magnólia

Faculdade de Ciências Agronômicas
Universidade Estadual Paulista
Fazenda Experimental Lageado s/n
Caixa Postal 237
Botucatu/SP 8603-970
Brazil
Phone : +551468234741 / Fax : +551468213438
magnolia@fca.unesp.br, magsilvas@bol.com.br

Mrs. Simin, Natasa

Institute of Chemistry
Faculty of Sciences
Trg Dositeja Obradovica, 3
Novi Sad 21000
Yugoslavia
Phone : +38121350122 / Fax : +3812154065
mimica@ih.ns.ac.yu

Dr. Simionatto, Euclésio

Universidade Federal de Santa Maria
Campus Universitário
Santa Maria 97105-900
Brazil
Phone : +5522082058869
a2060469@alunop.ufsm.br

Prof. Dr Stahl-Biskup, Elisabeth

University of Hamburg
Dept of Pharmaceutical Biology and Microbiology
Bundesstrasse, 45
Hamburg D-20146
Germany
Phone : +4940428383896 / Fax : +4940428383895
elisabeth.stahl-biskup@uni-hamburg.de

Dr. Stefanini, Mirian Baptista

Faculdade de Ciências Agronômicas
UNESP-Universidade Estadual Paulista
Fazenda Lageado, s/n
Campus de Botucatu
Botucatu/SP 18603-970
Brazil
Phone : +551468027172 / Fax : +551468213438
mstefanini@fca.unesp.br

Dr. Storesund, Hans Johan

Department of Pharmacy
University of Oslo
P.O Box 1068
Oslo 0316
Norway
Phone : +4722855002 / Fax : +4722855947
h.j.storesund@farmasi.uio.no

Dr. Sumnu, Servet Gulum

Middle East Technical University
Food Engineering Dept.
Ankara 06531
Turkey
Phone : +903122105628 / Fax : +903122101270
gulum@metu.edu.tr

Prof. Dr Tapiéro, Claude

Université Montpellier II
CC006
Place Eugène Bataillon
Montpellier 34085
France
Phone : +33467143210 / Fax : +33467042029
tapiero@univ-montp2.fr

Mrs. Tavares, Christina Bianca

Faculdade de Medicina - Universidade do Porto
Serviço de Microbiologia
Hospital S. João
Av. Hernani Monteiro
Porto
Portugal
Phone : +351917304036
cbrj@hotmail.com

Ms. Termentzik, K.

Laboratory of Agricultural Chemistry
School of Agriculture
Aristotle University of Thessaloniki
Thessaloniki 54124, Greece
aterment@agro.auth.gr

Dr. Theodoro, Suzi Huff

Universidade de Brasília UNB
Centro de Desenvolvimento Sustentável
SAS Quadra 05 Bloco H - Sala 205
Brasília 70070-914
Brazil
Phone : +55613215001 / Fax : +55613228473
suzitheodoro@cds.unb.br

Mr. Tigges, Wolfgang

Givaudan Deutschland GmbH
Giselherstrasse, 11
Dortmund 44319
Germany
Phone : +492312186410 / Fax : +492312186266
wolfgang.tigges@givaudan.com

Dr. Tinoco, Maria Teresa

Universidade de Évora
R. Romão Ramalho, 59
Évora 7000-671
Portugal
Phone : +351266745300 / Fax : +351266744971
mtft@uevora.pt

Dr. Tomassini, Therezinha

Fundação Oswaldo Cruz
Far-Manguinhos
R. Sizenando Nabuco, 100
Rio de Janeiro 21041-250
Brazil
Phone : +5502139772490 / Fax : +5502122703912
tecoebarto@hotmail.com

Prof. Dr Tomi, Félix

Université de Corse
Route des Sanguinaires
Ajaccio 20000
France
Phone : +33495524122 / Fax : +33495524142
tomi@vignola.univ-corse.fr

Dr. Trilles, Bénédicte

Université de la Nouvelle-Calédonie
Unité de Production d'Espèces Végétales
P.O Box 4477
Nouméa 98847
New Caledonia
Phone : +687836160 / Fax : +687432415
coustenoblefh@canl.nc

Dr. Tulok, Mária Heltmanné

Szent István University
Dept of Medicinal and Aromatic Plants
29, Villány Str.
Budapest H-1118
Hungary
Phone : +3613726250 / Fax : +3613726330
mtulok@omega.kee.hu

Dr. Tzakou-Tzini, Olga

University of Athens
Dept of Pharmacognosy and Chemistry of Natural
Products / School of Pharmacy
Panepistimiopolis Zografou
Athens 15771
Greece
Phone : +30107274102 / Fax : +30107274591
tzakou@pharm.uoa.gr

Mrs. van Vuuren, Sandy

University of the Witwatersrand
Dept of Pharmacy & Pharmacology
7, York Road
Park Town 2193
South Africa
Phone : +27117172157 / Fax : +27116424355
vanvuurens@therapy.wits.ac.za

Dr. Vargas, Rubem

Faculdade de Engenharia - PUCRS
Av.Ipiranga, 6681
Porto Alegre 90619-900
Brazil
Phone : +555133203653 / Fax : +555133203653
rvargas@pucrs.br

Mrs. Vidal, Isabel

Revista Activa
Largo da Lagoa, 15-C
Linda-a-Velha 2795-116
Portugal
IsabelV@acj.pt

Dr. Viljoen, Álvaro

University of the Witwatersrand
Dept of Pharmacy & Pharmacology
7, York Road
Park Town 2193
South Africa
Phone : +27117172169 / Fax : +27116424355
viljoenam@therapy.wits.ac.za

Mr. Waraschitz, Wolfgang

Obere Hauptstrasse, 28
Lasse A-2291
Austria
Phone : +4322132254 / Fax : +4322133138
wolfgang.waraschitz@A1.net

Prof. Dr Wawrzencyk, Czeslaw

Agricultural University
Dept of Chemistry
Norwida, 25
Wroclaw 50-375
Poland
Phone : +480713205257 / Fax : +480713284124
c-waw@ozi.ar.wroc.pl

Dr. Weber, Joerg

University of Applied Science Jena
Carl-Zeiss-Promenade, 2
Jena 07745
Germany
Phone : +493641205623 / Fax : +493641205601
joerg.weber@fh-jena.de

Mrs. Wenham, Tracey

Botanix, Ltd
Hop Pocket Lane, Paddock Wood
Tonbridge Kent TN12 6DQ
UK
Phone : +441892833415 / Fax : +441892836987
tracey.wenham@botanix.co.uk

Dr. Zawirska-Wojtasiak, Renata

Institute of Food Technology
Agricultural University of A Cieszkowski
Wojska Polskiego, 31
Poznan 60-624
Poland
Phone : +48618487275 / Fax : +48618487314
renazaw@owl.au.poznan.pl