





43<sup>rd</sup>  
ISEO

# International Symposium on Essential Oils

5 to 8 September 2012  
Lisbon – Portugal



[iseo2012.fc.ul.pt](http://iseo2012.fc.ul.pt)



**43<sup>rd</sup>**  
**ISEO**

**International  
Symposium  
on Essential Oils**

5 to 8 September 2012  
Lisboa • Portugal

## **Program, Book of Abstracts and Participants List**

### **Organized by**

Prof. Dr Ana Cristina Figueiredo  
Prof. Dr José Gonçalves Barroso  
Prof. Dr Luis Gaspar Pedro

# ***43rd International Symposium on Essential Oils (ISEO2012)***

**Faculty of Sciences of Lisbon, 5 to 8 September 2012, Lisbon, Portugal**

**Program,  
Book of Abstracts  
and Participants List**

**Editors**

Ana Cristina Figueiredo  
José Gonçalves Barroso  
Luis Gaspar Pedro

**Publisher**

Centro de Biotecnologia Vegetal - Fundação da Faculdade de Ciências de Lisboa

**Printer**

Europress

**Cover**

In9

**Legal deposit:** 347181/12

**ISBN:** 978-989-20-3188-0

Additional copies can be obtained from the Editors at Faculdade de Ciências de Lisboa, C2, Piso 1,  
Campo Grande, 1749-016 Lisboa, Portugal

The content of the Abstracts is the authors responsibility

## Table of Contents

<b>Welcome Addresses</b> .....	<b>1</b>
<b>Acknowledgements</b> .....	<b>3</b>
<b>Young Students Fellowships</b> .....	<b>4</b>
<b>Organization</b> .....	<b>5</b>
<b>General Information</b> .....	<b>7</b>
<b>Scientific Program</b> .....	<b>9</b>
<b>Social Program</b> .....	<b>12</b>
<b>Plenary Lectures Abstracts</b> .....	<b>13</b>
PL 1. Umami and Kokumi: latest developments in taste perception .....	15
PL 2. The "invisible hand" of floral volatiles in pollination ecology: dosage, context and signal evolution .....	16
PL 3. Ecological role of essential oils and volatiles in plants .....	17
PL 4. Aroma compounds in tomato flavour .....	18
PL 5. The molecular mechanisms of essential oil composition .....	19
PL 6. Chemical ecology of essential oils: analytical methods and biological activity .....	20
PL 7. Are essential oils natural products? .....	21
PL 8. Propolis volatiles: chemical diversity and biological activity .....	22
<b>Oral Communications Abstracts</b> .....	<b>23</b>
OC 1. Essential oils as allelochemicals – the mode of action and their potential use as bio-herbicides .....	25
OC 2. Qualitative and quantitative analysis of <i>Vetiver</i> oils by comprehensive two-dimensional Gas-Chromatography .....	26
OC 3. Dealing with phthalic-acid ester (PAE) residues in raw materials and compounded flavourings .....	27
OC 4. Kinetics of isolation of essential oils from plants .....	28
OC 5. Structural factors in the odor of $\alpha$ -santalol derivatives .....	29
OC 6. Cryptic speciation in liverworts: <i>Conocephalum conicum</i> , <i>Reboulia hemisphaerica</i> and <i>Marchantia polymorpha</i> .....	30
OC 7. Synergistic biological activity and chemodiversity of Himalayan medicinal plants from Nepal .....	31
OC 8. Screening of essential oils for inhibitory effects on the activity of BACE-1, the $\beta$ -secretase involved in Alzheimer's disease .....	32
OC 9. Portuguese Lavenders as a source of anti-inflammatory drugs .....	33
<b>Young Scientists (YS) Oral Communications Abstracts</b> .....	<b>35</b>
YSOC 1. Effects of PSBS gene silencing by RNAi on volatile compound biosynthesis and accumulation in tomato ( <i>Solanum lycopersicum</i> ) .....	37
YSOC 2. Coriander oil antimicrobial activity: A flow cytometric study .....	38
YSOC 3. Application of vibrational spectroscopy and chemometrics as a rapid and low cost alternative in the quality assessment of indigenous South African essential oils .....	39
YSOC 4. Effect of hydroxypropyl- $\beta$ -cyclodextrins on the antioxidant activity of <i>Lavandula</i> essential oils .....	40
YSOC 5. Classification of Lavender essential oils for the evaluation of sedative effects on Human .....	41
YSOC 6. Essential oils do not inhibit <i>Escherichia coli</i> peptidyl-tRNA hydrolase .....	42
<b>Posters Abstracts</b> .....	<b>43</b>
<b>Indexes</b> .....	<b>266</b>
<b>Species</b> .....	<b>268</b>
<b>Subjects</b> .....	<b>271</b>
<b>Authors</b> .....	<b>276</b>
<b>Participants List</b> .....	<b>283</b>
<b>Sponsors</b> .....	<b>293</b>



## Welcome Addresses

Welcome

On behalf of the *Faculdade de Ciências da Universidade de Lisboa* I am quite pleased to welcome you to our University and to the 43<sup>rd</sup> *International Symposium on Essential Oils*.

This is the second time that such an important meeting takes place in Portugal at our facilities. I believe that is a clear indication that happy memories remain from the first occasion and we are very honoured for hosting you again. Also, I want to congratulate the organizers of the Symposium, specially my colleagues Ana Cristina Figueiredo, José Gonçalves Barroso and Luis Gaspar Pedro who made a great effort for putting things together, timely and efficiently.

Looking at your scientific programme, I realize that you planned an intense activity with several plenary lectures and working sessions for oral communications and posters presentations. Hence, I anticipate four days of fruitful scientific exchange but plenty of social contact among all the participants. In particular, I hope that you still have some spare time to visit our beautiful city and to enjoy the almost mythical light of Lisbon.

I wish you all a very pleasant stay in Portugal.

Prof. Dr. José Manuel Pinto Paixão  
Dean of the *Faculdade de Ciências da Universidade de Lisboa*

Welcome

As Director of the *Laboratório Associado Institute for Biotechnology and Bioengineering (IBB)*, a network of research units in Portugal, in which the Plant Biotechnology Research Group (Centro de Biotecnologia Vegetal, CBV), that organizes this event, is integrated, I am delighted to welcome you all to Lisbon to contribute and attend the *43rd International Symposium on Essential Oils (ISEO2012)*.

The organization of the ISEO2012, is not only timely and relevant, but also of utmost importance for both the CBV and to IBB in view of the increasing importance of this research field in Portugal and abroad. The scientific program of this meeting promises compelling science in the field of essential oils as well as in related fields, namely ecology, odour perception, biological activities, among others. The outstanding level of the invited speakers as well as of many oral and poster contributions warrants the success of the Symposium.

I would like to express my hearty greetings and thanks to scientists and researchers, coming from all around the world, who brought to this venue their know-how and the result of years of studies, to share with the scientific community, the general public, and the media.

I am certain that the ISEO2012 is the right forum for further consolidation and development of the medicinal and aromatic plants field, in general, and essential oils and volatile compounds in particular.

Despite the tight schedule of the scientific program, I hope you can find out some time to enjoy the social and cultural wealth and allurements of our beautiful city of Lisbon.

I therefore renew my welcome, and wish you all a pleasant and fruitful stay.

Prof. Dr Joaquim M. Sampaio Cabral  
The Director of the *Laboratório Associado Institute for Biotechnology and Bioengineering (IBB)*

## Acknowledgements

The 43rd International Symposium on Essential Oils was generously sponsored by:

- ❖ Faculdade de Ciências da Universidade de Lisboa
- ❖ Centro de Biotecnologia Vegetal (CBV), Institute for Biotechnology and Bioengineering (IBB)
- ❖ International Federation of Essential Oils and Aroma Trades (IFEAT)
- ❖ Fundação Calouste Gulbenkian
- ❖ Estética Viva
- ❖ John Wiley & Sons Limited
- ❖ Milestone
- ❖ Taylor & Francis group
- ❖ ILC-Instrumentos de Laboratório e Científicos, Lda
- ❖ Fundação para a Ciência e a Tecnologia (FCT) under CBV / IBB plurianual funding and research contract PTDC/AGR-CFL/117026/2010

The generous obligingness of the following entities is gratefully acknowledged:

- ❖ Bacardi Martini Portugal, Lda
- ❖ Ach. Brito & C.<sup>a</sup>, S.A.
- ❖ Quinta Picos do Couto Lda
- ❖ Gorreana Chás
- ❖ Universidade de Lisboa
- ❖ Câmara Municipal de Lisboa
- ❖ Planalto Dourado
- ❖ Associação Portuguesa de Horticultura

Gratitude is given to all those that contributed to make this event public:

- ❖ Centro de Informática da Faculdade de Ciências da Universidade de Lisboa
- ❖ Gabinete de Comunicação, Imagem e Cultura da Faculdade de Ciências da Universidade de Lisboa
- ❖ Câmara Municipal de Lisboa
- ❖ Estética Viva
- ❖ Associação Portuguesa de Horticultura
- ❖ John Wiley & Sons Limited

ISEO2012 Organizing Committee deeply acknowledges the Young Scientists Fellowships Selection Committee for the excellent and invaluable involvement in the evaluation of Young Scientists CVs and abstracts contributions. Gratitude is also given to the ISEO permanent Scientific Committee and ISEO2012 Scientific Committee for their commitment.



## Young Students Fellowships

Thanks to the support of the *International Federation of Essential Oils and Aroma Trades* (IFEAT) (14) together with the ISEO2012 Organizing Committee (6), a total of twenty *Registration Fellowships* to assist young researchers to attend ISEO2012, could be available.

The Young Scientists ISEO2012 *Registration Fellowship* Selection Committee has accepted 37 out of the 40 application proposals. Candidates from seventeen countries (Algeria, Brazil, China, Czech Republic, Germany, Iran, Italy, Japan, Poland, Portugal, Romania, Serbia, South Africa, Spain, The Netherlands, USA and Yemen) submitted an appropriate application.

In total 20 high quality abstracts and applicants CVs were recommended by the Selection Committee as *Registration Fellowship* award winners. Due to the tight symposium schedule not all works could be presented as oral communications. Thus, of the selected 20 contributions, 14 were set as poster presentations. The remaining 6 were chosen for oral presentations, in a special session devoted to Young Scientists ISEO2012.

The *Registration Fellowship* award consists of the student registration fee reimbursement (€265). The awardees will also be entitled to a certificate stating their prize and the awardees names will be listed in the ISEO2012 book of abstracts.

During the conference, the Fellowships Selection Committee will announce and reimburse the awardees with the pecuniary prize (€265) and prize certificate. The first author of the selected 20 contributions will receive the award during the special session devoted to Young Scientists ISEO2012. Six awardees will additionally be offered the Symposium dinner, supported by *John Wiley & Sons, Milestone, Taylor & Francis and Estética Viva*.

- ❖ The International Federation of Essential Oils and Aroma Trades (IFEAT) and the Organizing Committee supported the Registration fees of the following students:

Andrea Occhipinti - University of Turin, Italy.

Filomena Silva - Universidade da Beira Interior, Portugal.

Maxleene Sandasi - Tshwane University of Technology, South Africa.

Patrícia Costa - Universidade do Algarve, Portugal.

Makiko Kitao - Kinki University, Japan.

Rachel Thompson - University of Alabama in Huntsville, USA.

Jakub Smid - Czech University of Life Sciences, Czech Republic.

Jorge Faria - Faculdade de Ciências da Universidade de Lisboa, Portugal.

Nikola Stojanovic - University of Niš, Serbia.

Adela Frankova - Czech University of Life Sciences, Czech Republic.

Amel Bouzabata - University of Corsica, France.

Satoshi Asada - Meijo University, Japan.

Inés Méndez Tovar - Agricultural Technical Institute of Castilla y León, Spain.

Jennifer Bufalo - Universidade Estadual Paulista, Brazil.

Michelle Cristiane Bufalo - Universidade Estadual Paulista, Brazil.

Beata Zawitowska - Nicolaus Copernicus University, Poland.

Agustina Nurcahyanti - Heidelberg University, Germany.

Miljana Đorđević - University of Niš, Serbia.

Prabodh Satyal - University of Alabama in Huntsville, USA.

Ana Miltojevic - University of Niš, Serbia.

## Organization

### ISEO2012 Organizing Committee


Prof. Dr Ana Cristina Figueiredo  
 Prof. Dr José Gonçalves Barroso  
 Prof. Dr Luis Gaspar Pedro

### ISEO2012 Scientific Committee

Prof. Dr Ana Cristina Figueiredo  
 Prof. Dr Carlos Cavaleiro  
 Prof. Dr Hans Scheffer  
 Prof. Dr Helena Trindade  
 Prof. Dr José Gonçalves Barroso  
 Prof. Dr Lígia Salgueiro  
 Prof. Dr Luis Gaspar Pedro

### Symposium Venue

The Faculty of Sciences of Lisbon hosts the *43rd International Symposium on Essential Oils*.



**WWW Information**

<http://iseo2012.fc.ul.pt/>

**International Symposium on Essential Oils (ISEO) Permanent Scientific Committee**

Dr Agnieszka Ludwiczuk  
Dr Alain Chaintreau  
Prof. Dr Alvaro Viljoen  
Prof. Dr Ana Cristina Figueiredo  
Prof. Dr Carlo Bicchi  
Prof. Dr Chlodwig Franz  
Prof. Dr Fatih Demirci  
Prof. Dr Gerhard Buchbauer  
Prof. Dr Hans Scheffer  
Prof. Dr Hüsnü Can Başer  
Dr Jan Demyttenaere  
Prof. Dr Jan Karlsen  
Prof. Dr Johannes Novak  
Prof. Dr Karl-Heinz Kubeczka  
Prof. Dr Luigi Mondello  
Prof. Dr Nicolas Baldovini  
Prof. Dr Patrizia Rubiolo  
Prof. Dr. Stanislaw Lochynski  
Prof. Dr Yoshinori Asakawa  
Prof. Dr Zamborine Eva Nemeth

**Organizing Committee Contact**

Faculdade de Ciências de Lisboa  
Centro de Biotecnologia Vegetal  
Departamento de Biologia Vegetal  
C2, Piso 1, Campo Grande 1749-016 Lisboa, Portugal  
Tel.: +351217500257 | FAX: +351217500048  
email: [iseo2012@fc.ul.pt](mailto:iseo2012@fc.ul.pt)

**Secretariat Address**

Cidália Pacheco / Cristina Vicente  
Rua Augusto Macedo, 12-D Escritório 2  
1600-503 Lisboa – Portugal  
Tels.: +351217120778 / 79 / 80 | Fax: +351217120204  
[cidaliampacheco@veranatura.pt](mailto:cidaliampacheco@veranatura.pt)

## General Information

### Symposium Office

The Symposium Office will be located at the Foyer of the Lecture Halls in the building C3 of the Faculty of Sciences of Lisbon.

Registration and Opening Hours:

Wednesday, September 5:	14:00 – 19:00
Thursday, September 6 and Friday, September 7:	09:00 – 18:00
Saturday, September 8:	09:00 – 13:00

### Symposium Language

English will be the official language of the ISEO2012. There will be no simultaneous translation.

### Badges

Participants are requested to wear their Symposium badges for identification and admittance to Scientific and Social venues. Symposium badges are required at the lunch's canteen.

### Plenary Lectures and Oral Communications

The Plenary Lectures are limited to 50min, Oral Communications to 10min, all with plus 5min for discussion. A room for presentation preview will be available. MS-Power Point presentation facilities are available. Presentation files should be handed at the Symposium Office, in the Foyer of the Lecture Halls, with the name of the presenting author at least 1h prior to the beginning of the respective session.

### Posters

All posters will remain exhibited throughout the symposium and the authors are recommended to be present at their posters during posters sessions.

Two poster sessions will be scheduled, odd numbers on Thursday and even numbers on Friday.

Poster boards measuring 90cm wide and 120cm height will be available for poster presentation with portrait orientation and A0 size (84.1 cm X 118.9cm). The Registration Office provides fixation facilities.

**Meals**

Two lunches, of Thursday, September 6, and Friday, September 7, 2012 are included in the registration fee of regular and student participants.

The lunches will be served at the C6 building Canteen (information on location provided at *Symposium Venue* section).


The Canteen serves meals from 12:00 to 14:00. Personal badges are required at the Canteen. Accompanying persons lunch vouchers can be purchased at the Symposium Office. Neither the Canteen nor the Campus coffee shops accept foreign currency.

There are also several restaurants, snack bars and coffee shops in the area around the University Campus.

**Currency**

All payments should be done in Euro.

**Means of conveyance to the Symposium venue**

 796 - 778 - 777 - 768 - 767 - 764 - 755 - 747 - 738 - 735 - 701 - 207 - 108 - 36

 Cidade Universitária - Campo Grande

**Insurance / Liability**

The Organizers are in no respect responsible for any accommodation problem, nor for any accident, injury, theft, loss and property damage to any person during the Symposium.

## Scientific Program

### Wednesday, September 5, 2012

All lectures, the poster presentations and the symposium office will be located at the building C3 of the Faculty of Sciences of Lisbon (information on location provided at *Symposium Venue* section).

The Get-together party will be held at the building C6 of the Faculty of Sciences of Lisbon (information on location provided at *Symposium Venue* section).

14.00 - 19.00	<b>Registration at the Symposium Office</b>	Foyer of the Lecture Halls, C3
17.00 - 17.45	<b>Opening Ceremony</b> <b>Welcome Addresses</b> Performance of the <i>Vicentuna</i>	Main Lecture Hall
18.00 - 19.00	<b>Plenary Lecture</b> Chair: Prof. Dr Karl-Heinz Kubeczka <b>Umami and Kokumi: latest developments in taste perception</b> <i>Dr Chris Winkel, Givaudan, Flavour Ingredients Research, the Netherlands</i>	Main Lecture Hall
19.00 - 21.00	<b>Get-together Party</b>	C6 building canteen

### Thursday, September 6, 2012

09.00 – 10.00	<b>Plenary Lecture</b> Chairs: Prof. Dr Johannes Scheffer & Prof. Dr Lígia Salgueiro <b>The “invisible hand” of floral volatiles in pollination ecology: dosage, context and signal evolution</b> <i>Prof. Dr Robert Raguso, Cornell University, USA</i>	Main Lecture Hall
10.00 – 10.15	<b>Oral Communications</b> <b>Essential oils as allelochemicals – the mode of action and their potential use as bio-herbicides</b> <i>Dr Nativ Dudai, Division of Aromatic and Medicinal Plants, Israel</i>	Main Lecture Hall
10.15 – 11.00	<b>Coffee Break</b>	Foyer of the Lecture Halls
11.00 – 12.00	<b>Plenary Lecture</b> Chair: Prof. Dr Agnieszka Ludwiczuk <b>Ecological role of essential oils and volatiles in plants</b> <i>Prof. Dr Anna-Karin Borg-Karlson, KTH, Stockholm</i>	Main Lecture Hall
12.00 – 14.00	<b>Lunch</b>	C6 Canteen/Local Restaurants
14.00 – 15.00	<b>Plenary Lecture</b> Chair: Prof. Dr Chlodwig Franz <b>Aroma compounds in tomato flavour</b> <i>Prof. Dr Denise Teman, University of Florida, USA</i>	Main Lecture Hall
15.00 – 16.00	<b>Oral Communications</b> Chairs: Prof. Dr Patrizia Rubiolo & Prof. Dr Stanislaw Lochynski	Main Lecture Hall
15.00 – 15.15	<b>Qualitative and quantitative analysis of Vetiver oils by comprehensive two-dimensional Gas-Chromatography</b> <i>Dr Jean-Jacques Filippi, Université de Nice-Sophia Antipolis, France</i>	
15.15 – 15.30	<b>Dealing with phthalic-acid ester (PAE) residues in raw materials and compounded flavourings</b> <i>Dr Thierry Cachet, International Organization of the Flavor Industry (IOFI), Belgium</i>	
15.30 – 15.45	<b>Kinetics of isolation of essential oils from plants</b> <i>Dr Helena Sovova, ASCR, Czech Republic</i>	
16.00 – 17.00	<b>Coffee Break</b>	Foyer of the Lecture Halls
16.00 – 19.00	<b>Poster Session (Odd Numbers)</b>	Foyers of the Lecture Halls

**Friday, September 7, 2012**

09.00 – 10.00	<b>Plenary Lecture</b>	Main Lecture Hall
	Chairs: Prof. Dr Helena Trindade & Prof. Dr Eva Nemeth <b>The molecular mechanisms of essential oil composition</b> Prof. Dr Jörg Degenhardt, Martin Luther University Halle-Wittenberg, Germany	
10.00 – 10.15	<b>Oral Communications</b>	Main Lecture Hall
	<b>Structural factors in the odor of <math>\alpha</math>-santalol derivatives</b> Dr Toshio Hasegawa, Saitama University, Japan	
10.15 – 11.00	<b>Coffee Break</b>	Foyer of the Lecture Halls
11.00 – 12.00	<b>Plenary Lecture</b>	Main Lecture Hall
	Chair: Prof. Dr Carlo Bicchi <b>Chemical ecology of essential oils: analytical methods and biological activity</b> Prof. Dr Alés Svatós, Max Planck Institute, Germany	
12.00 – 14.00	<b>Lunch</b>	C6 Canteen/Local Restaurants
14.00 – 15.00	<b>Plenary Lecture</b>	Main Lecture Hall
	Chair: Prof. Dr Jan Karlsen <b>Are essential oils natural products?</b> Dr Daniel Joulain, SCBZ, France	
15.00 – 16.30	<b>Young Scientists Symposium</b>	Main Lecture Hall
	Chairs: Prof. Dr Alvaro Viljoen and Dr Jan Demyttenaere	
15.00 – 15.10	<b>Effects of PSBS gene silencing by RNAi on volatile compound biosynthesis and accumulation in tomato (<i>Solanum lycopersicum</i>)</b> Dr Andrea Occhipinti, University of Turin, Italy	
15.15 – 15.25	<b>Coriander oil antimicrobial activity: A flow cytometric study</b> Dr Filomena Silva, CICS-UBI, Portugal	
15.30 – 15.40	<b>Application of vibrational spectroscopy and chemometrics as a rapid and low cost alternative in the quality assessment of indigenous South African essential oils</b> Dr Maxleene Sandasi, Tshwane University of Technology, South Africa	
15.45 – 15.55	<b>Effect of hydroxypropyl-<math>\beta</math>-cyclodextrins on the antioxidant activity of Lavandula essential oils</b> Dr Patricia Costa, University of Algarve, Portugal	
16.00 – 16.10	<b>Classification of Lavender essential oils for the evaluation of sedative effects on Human</b> Dr Makiko Kitao, Kinki University, Japan	
16.15 – 16.25	<b>Essential oils do not inhibit Escherichia coli peptidyl-tRNA hydrolase</b> Dr Rachel Thompson, University of Alabama in Huntsville, USA	
16.30 – 17.30	<b>Coffee Break</b>	Foyer of the Lecture Halls
17.00 - 19.00	<b>Poster Session (Even Numbers)</b>	Foyers of the Lecture Halls
19.30 – 23.00	<b>Symposium Dinner</b>	Casa do Leão, at Castelo de S. Jorge

**Saturday, September 8, 2012**

09.00 – 10.00	<b>Plenary Lecture</b>	Main Lecture Hall
	Chairs: Prof. Dr Gerhard Buchbauer & Prof. Dr Paola Dugo <b>Propolis volatiles: chemical diversity and biological activity</b> Prof. Dr Vassya Bankova, Institute of Organic Chemistry, Bulgaria	
10.00 – 10.15	<b>Oral Communications</b>	Main Lecture Hall
	<b>Cryptic speciation in liverworts: <i>Conocephalum conicum</i>, <i>Reboulia hemisphaerica</i> and <i>Marchantia polymorpha</i></b> Prof. Dr Agnieszka Ludwiczuk, Medical University of Lublin, Poland	
10.15 – 11.00	<b>Coffee Break</b>	Foyer of the Lecture Halls

---

11.00 – 12.00	<b>Oral Communications</b>	Main Lecture Hall
	Chairs: Prof. Dr Yoshinori Asakawa & Prof. Dr Carlos Cavaleiro	
11.00 – 11.15	<b><i>Synergistic biological activity and chemodiversity of Himalayan medicinal plants from Nepal</i></b> <i>Dr Prabodh Satyal, University of Alabama in Huntsville, USA</i>	
10.15 – 11.30	<b><i>Screening of essential oils for inhibitory effects on the activity of BACE-1, the <math>\beta</math>-secretase involved in Alzheimer's disease</i></b> <i>Dr Rita Videira, University of Coimbra, Portugal</i>	
10.30 – 11.45	<b><i>Portuguese Lavenders as a source of anti-inflammatory drugs</i></b> <i>Dr Mónica Zuzarte, University of Coimbra, Portugal</i>	
12.00 – 13.00	<b>Closing Ceremony</b>	Main Lecture Hall
	<b>Performance of the Vicentina</b>	

---



## Social Program

### Get-together Party

*Wednesday – 5 September*

The Get-together Party is included in the registration fee. The Get-together party will be held at the building C6 of the Faculty of Sciences of Lisbon (information on location provided at *Symposium Venue* section).

### Symposium Dinner

*Friday – 7 September*

The Symposium dinner will take place at *Casa do Leão*, located at *Castelo de S. Jorge*, downtown Lisbon.

The Symposium dinner is not included in the registration fee, and participants wishing to join it may register to take part in it at the Symposium Office.

### Tours

The Secretariat makes available different tours programs. For Tours booking, contact the ISEO2012 Secretariat.

## Plenary Lectures Abstracts





**PL 1. Umami and Kokumi: latest developments in taste perception**

Winkel C

*Givaudan, Flavour Ingredients Research, Huizerstraatweg 28, 1411 GP Naarden, the Netherlands*

*Chris.winkel@givaudan.com*

Keywords: Umami, Kokumi, Taste

Since health has become one of the main drivers in nutrition, the scientific interest in taste perception is high. Academia as well as food and flavour companies aim to make food more healthy but not less appetizing. A good example is salt (NaCl) reduction. Salt has a pleasant taste but too much sodium is considered unhealthy. Salt reduction per se will therefore lead to less attractive food. Currently there is no good replacer for sodium chloride. To compensate for the loss in salty taste, umami taste and kokumi taste can help.

This review will give a general overview on savoury taste with a focus on umami taste, kokumi taste and syneristic effects of odour and taste molecules.

Umami is now generally accepted to be a primary taste and the umami receptor is known. Many new molecules that have umami taste or umami enhancing properties have been discovered.

Kokumi is a much less defined taste property. Kokumi has been linked to mouthfeelness, complexity and authenticity. Only recently the first papers dealing with this effect have been published

Synergy of odour and taste is known for some time for sweet tastants. Recently this effect has been studied for savoury tastants as well.

## PL 2. The “invisible hand” of floral volatiles in pollination ecology: dosage, context and signal evolution

Raguso RA<sup>1</sup>, Moré M<sup>2</sup>; Svensson G<sup>3</sup>

<sup>1</sup>Cornell University, USA;

<sup>2</sup>Universidad Nacional de Córdoba, Argentina;

<sup>3</sup>Lund University, Sweden

rar229@cornell.edu

Keywords: Bioassays, Deception, Floral scent, Foraging, Mutualism

Floral volatiles have long been appreciated as pollinator attractants in specialized tropical plants or those engaged in obligate mutualisms (e.g. figs with fig wasps). In contrast, generalized plant-pollinator interactions more commonly studied in ecological or agricultural contexts typically are considered visually-mediated phenomena, with scant reference to floral chemistry. Here I discuss the evidence challenging this perception, focusing on how context determines the ecological functions of plant volatiles. I will use common floral volatiles – 2-phenylethanol (2PE), dimethyl disulfide (DMDS) and (E)-4,8-dimethyl-1,3,7-nonatriene (DMNT) – to illustrate my thesis.

Volatiles were analyzed using static (SPME) or dynamic headspace collection coupled with GC-MS, and behavioural bioassays monitoring pollinator responses to flowers whose odours were manipulated using extracts, emitters or traps.

2PE is common in diurnal wildflowers pollinated by bees. As a low dosage pollen odour, 2PE attracts andrenid bees to *Fragaria* flowers (1), but at higher dosage in floral nectar, it repels ants and limits bumblebee visits to *Polemonium* flowers, reducing nectar thievery and optimizing selection on floral shape (2). DMDS is a universal component of deceptive flowers that mimic carrion. When present in diurnal, temperate flowers or boreal mosses, it attracts flies that disperse the plants' pollen or spores (3, 4). However, when DMDS is emitted by nocturnal, nectar rich flowers of Neotropical vines, it attracts bats as pollinators (5). Finally, DMNT is a common herbivore-induced vegetative volatile in maize, cotton and other crop plants, attracting parasitic wasps as an indirect defense. However, DMNT is the primary floral attractant in two specialized systems, a South American bee-pollinated orchid (6) and an obligate North American yucca-yucca moth interaction (7). Concentration, timing and community context determine the functional roles of these three common floral volatiles.

1. TL Ashman et al. (2005) Ecology 86: 2099-2105.

2. C Galen et al. (2011) Am. Nat. 177: 258-272.

3. M Moré et al. (2012) Ann. Bot. (in review).

4. P Marino et al. (2009) Symbiosis 47: 61-76.

5. von Helversen et al. (2000) J. Comp. Physiol. (A) 186: 143-153.

6. A Wiemer et al. (2009) Plant Biology 11: 506-514.

7. Svensson et al. (2011) Oikos 120: 1577-1583.

### PL 3. Ecological role of essential oils and volatiles in plants

Borg-Karlson A-K

*KTH, School of Chemical Science and Engineering, Department of Chemistry, Ecological Chemistry group, 100 44 Stockholm*

*akbk@kth.se*

Keywords: Volatiles, Terpenoids, Chirality, Behavior

Essential oils are important and sustainable sources for bioactive compounds, especially for those with complex molecular structures that are difficult or expensive to synthesize. Plants compounds are often used for control of insects and microorganisms as they function as repellents, antifeedants, inhibitors, bactericides, and fungicides (1). Some of these compounds have more than one biological function and can in specific cases attract and/ or repel various insects. Attacking insects or microorganisms induce the biosynthesis of the plant which then starts to produce more of specific compounds or synthesize new which function as physical and biological barriers towards the infestations (2). A large number of the plant compounds are chiral and the importance of knowing the different function of the enantiomers will be exemplified (4,5,6). Recent findings and ideas of how to utilize essential oils for controlling insects and other organisms will be discussed.

1. A. Sofrata, et al. (2011) Plos One 11
2. T. Zhao, et al. (2011) Oecologia: Volume 167, Issue 3, Page 691-699
3. K. Pålsson. Et al. (2008) J Med Entom. 45,1,88-93(6)
4. A.-K.-Borg-Karlson, et al. (2003) J.Chem. Ecol. 29,1,1-14.
5. M. Strandén, et al. (2002) Chemical Senses. 27:2,142-152
6. A. Wibe, et al. (1998) J. Chem Ecol. 24, 2

#### PL 4. Aroma compounds in tomato flavour

Tieman DM, Klee HJ

*University of Florida Horticultural Sciences Department, Gainesville, FL 32611, USA*

*dtieman@ufl.edu*

Keywords: Tomato, *Solanum lycopersicum*, Solanaceae, Aroma, Volatiles

Flavour of modern tomato varieties is generally considered to need improvement. Older heirloom varieties are considered to have better flavour, but do not have the agronomic properties for commercial production. Tomato flavour is a result of interactions between sugars, acids and aroma compounds. Analysis of heirloom tomato aroma volatiles indicated that these varieties have surprising diversity in aroma compound levels. This diversity allows us to determine the aroma volatiles important for good tasting tomatoes. Traditional methods have focused on aroma compounds that are above the threshold of detection by the human olfactory system; however, this method does not account for interactions between compounds in the mouth and nose. We have analyzed the biochemical components of flavour in over 60 heirloom varieties. Correlations between aroma compound levels and ratings from consumer taste panels have allowed us to define the components of a good tasting tomato. Some aroma volatiles that were identified as important for tomato flavour using the odour threshold method are shown to not contribute to tomato flavour using these methods. In addition, we have identified volatiles that add to the perceived sweetness of tomato fruit. Using this approach we can identify targets for breeding better tasting tomatoes.

## PL 5. The molecular mechanisms of essential oil composition

Degenhardt J, Schimmel J, Krause S

Martin Luther University Halle-Wittenberg, Institute for Pharmacy, Hoher Weg 8, 06120 Halle/Saale, Germany

[joerg.degenhardt@pharmazie.uni-halle.de](mailto:joerg.degenhardt@pharmazie.uni-halle.de)

Keywords: Thyme, Chemotype regulation, Terpene biosynthesis

Essential oils are usually complex mixtures of monoterpenes, sesquiterpenes and simple phenylpropanoids. The economic value of an essential oil depends on its composition, especially the presence of compounds that are pharmaceutically active, antimicrobial, or impart a desired scent or flavour. On the other hand, compounds with toxic and irritant properties or undesired flavours need to be absent from the oil. For many centuries, selection and classical breeding of essential oil plants have been used to optimize oil production. However, some components of the oils are difficult to separate from each other and the essential oil yield of a plant is hard to raise. In recent years, the biosynthesis of essential oils has been studied with molecular and biochemical methods. These efforts provide a better understanding of the biosynthesis of the essential oils and can reveal new strategies for the breeding of high-value plants.

Common thyme (*Thymus vulgaris*) is an excellent system to study the biosynthesis of essential oils and its regulation. Thyme forms several chemotypes which are morphologically identical but differ in the terpene content of their essential oils. Each chemotype is characterized by its dominant monoterpene alcohol, which can be geraniol, alpha-terpineol, sabinene hydrate, linalool, carvacrol, or thymol. Two of these monoterpenes, the phenolic monoterpene alcohols thymol and carvacrol, are especially valuable due to their antibacterial, antiseptic and spasmolytic effects. In all chemotypes, we characterized the key enzymes of terpene biosynthesis, the class of terpene synthases. Since most of these terpene synthases form multiple products, relatively few enzymes are sufficient to generate the complex terpene blends of each of the chemotypes. While each chemotype appears to have a complete set of terpene synthase genes coded in its genome, only the genes necessary for the chemotype-specific terpenes are activated. The terpene synthase TPS1 is expressed in the carvacrol and thymol types. The major product of TPS1 is  $\gamma$ -terpinene, which can be converted to carvacrol and thymol. We identified a family of cytochrome P450 monooxygenases that might catalyse this conversion *in planta*. Elucidation of the biosynthetic pathways and the enzymes involved in the biosynthesis of essential oil components allows developing molecular markers for directed breeding efforts to optimize oil production.



## PL 6. Chemical ecology of essential oils: analytical methods and biological activity

Svatoš A

*Mass spectrometry/Proteomics research Group, Max Planck Institute for Chemical Ecology, Hans-Knöll-Straße 8, D-07745 Jena, Germany*

*svatos@ice.mpg.de*

Keywords: Structure elucidation, Mass spectrometry, Volatiles, Imaging

Essential oils (EO) are mixtures of individual compounds which are typically found in terpenoid skeletons; EOs have widespread industrial application. However, in nature, plants produce terpenoids for their own protection, survival, and reproduction [1]. To understand the biochemical processes behind EO production, first the chemical structures of these mixtures need to be identified. Due to its high sensitivity, modern mass spectrometry (MS) is an important tool for the structural elucidation of EOs. Additionally, nuclear magnetic resonance (NMR) is used to refine stereochemistry of the compounds.

In this lecture I will describe theoretical and practical aspects of the MS of terpenoids, placing specific emphasis on their biosynthesis and biological activity. Typically, EO mixtures are separated by gas-phase liquid chromatography, and the eluted peaks are analyzed by electron-ionization mass spectrometry (EI-MS) using databases of MS spectra [2]. However, the mass spectra of EO components are highly similar and more detailed analysis is typically needed to identify the compound. Additional chemical reactions (hydrogenation, ozonization, methylthiolation) should be performed to obtain additional structural information [3,4]. Very recently, a new computer algorithm ("fragmentation trees" [5]) was developed to assist researchers in high-throughput structure elucidation.

Dramatic developments in ambient MS technology now allow for the rapid sampling of compounds emitted from plant surfaces. Our current interest is to follow emission of compounds from individual cells in plants' organs. This method, called MS imaging, has already increased our understanding of the spatial distribution of non-volatile compounds [6,7] and of the function of these molecules in the chemical ecology of plants.

Acknowledgments: Financial and technical support from Max Planck Society is acknowledged with pleasure. Work of current and previous members of our group and of co-operating partners' is highly recognized and appreciated.

1. J Gershenzon, N Dudareva (2007) *Nature Chem. Biol.* 3: 408-414.
2. RP Adams (2007) *Identification of Essential Oil Components By Gas Chromatography/Mass Spectrometry*. 4th, Allured Pub Corp. 804 pp.
3. A Svatoš et al. (2004). *Rapid Commun. Mass Spectrom.* 18: 816-821.
4. T G Köllner et al. (2008) *J. Biol. Chem.* 283: 20779-20788.
5. F Rasche et al. (2011) *Anal. Chem.* 83: 1243-1251.
6. A Svatoš (2010) *Trends Biotech.* 28: 425-434.
7. J Kroiß (2010) *Nature Chem. Biol.* 6: 261-263.

## PL 7. Are essential oils natural products?

Joulain D

SCBZ Conseil, 99 avenue Sidi Brahim, F-06130 Grasse

dajoulain@wanadoo.fr

Keywords: Essential oils, Naturalness

During the last two decades, mainly by the vector of the communication on the Internet, we observed a considerable tendency to promote essential oils as beneficial natural products in many aspects of well-being, and in various domains such as cosmetic ingredients, aromatherapy, "aromachology", various more or less defined therapeutic applications, and foods, to cite only a few.

In considering on the one hand the strict definition of essential oils according to the ISO standard, and on the other hand the accepted and/or enforced definitions of natural products in various countries, one can observe a disparity of judgments, and often obvious inaccuracies, which account for public ignorance, and even of some professional or official organizations.

Regardless, the relevance of these more or less supported claims in relation with their alleged benefits, one can wonder whether essential oils as complex chemicals, are entirely constituted of the same substances which exist naturally in the plant raw material from which they are derived. As a result, one may question the relevance of their labelling as "natural" or "100% natural" or "100% pure and natural" and so on. In contrast, products obtained by other techniques often recover more faithfully the bulk of constituents originally present in the plant material, but are often deprived of the "natural" label by certain governmental or private organizations.

Examples will be presented to illustrate these considerations, with the aim of correcting inaccurate declarations or exaggerated claims.

## PL 8. Propolis volatiles: chemical diversity and biological activity

Bankova V

*Institute of Organic Chemistry with Centre of Phytochemistry, BAS, Sofia, Bulgaria*

*bankova@orgchm.bas.bg*

Keywords: Propolis, Essential oils, Biological activity, Plant origin

Propolis (bee glue) is a sticky material collected by bees from plants, and serves dual purposes in the hive: as building material and as defensive substance. It has been used as a remedy in Europe since ancient times for treatment of wounds and burns, sore throat, etc. Nowadays, propolis use in over-the-counter preparations, “bio”-cosmetics and functional foods and for other purposes continues to increase.

Modern science has confirmed the antimicrobial action of bee glue and has discovered many other beneficial pharmacological properties: immunostimulating, antitumor, anti-inflammatory, antiulcer, local anaesthetic, hepatoprotective, antioxidant. Volatile compounds are found in low concentrations in propolis, but their aroma and significant biological activity make them of importance for the characterisation of propolis. As bee glue is a plant derived product, its chemical composition is highly variable and depends on the local flora at the side of collection and on the choice of propolis plant sources made by bees. For this reason it offers a significant chemical diversity and the study of propolis from different locations often reveals new constituents and new bioactive compounds. The available data about chemical composition of propolis essential oils from different geographic regions is reviewed. The role of propolis volatiles in identification of its plant origin is also discussed. The contribution of essential oils and their constituents to the diverse biological activities of propolis is considered. Propolis essential oils have demonstrated antibacterial, antifungal, antiprotozoan activity, they were found to inhibit the proliferation of cancer cells by inducing apoptosis and to have immunostimulating effect in humans, etc. Future perspectives in research on propolis essential oils are outlined.

## Oral Communications Abstracts





## OC 1. Essential oils as allelochemicals – the mode of action and their potential use as bio-herbicides

Dudai N, Chaimovitsh D

*Division of Aromatic and Medicinal Plants, Neve Ya'ar Research Center, Agricultural Research Organization, Israel*

*Natvdud@gmail.com*

Keywords: Essential oils, Monoterpenes, Allelopathy, Allelochemicals

Allelopathic activity of aromatic plants has been described frequently. Essential oils inhibit germination and growth of various plants. Residues of leaves of aromatic plants in soil inhibit the growth and the emergence of seedlings. We have developed a good quantitative assay to test for the inhibitory activity of volatiles on seed germination and have screened dozens of essential oils and their components. The major active allelochemicals were found to be aldehydic and ketonic monoterpenes, having a  $\alpha$ - $\beta$  unsaturated double bond, such as citral, (geranial and neral) and pulegone. Exposure of seeds for 4h to the compounds was enough to cause inhibition of the rate of germination and of seedling development in wheat, *Amaranthus palmeri* and *Brassica nigra*. Using histochemistry, we observed that citral is absorbed by wheat seed through the abscission layer, and that it reaches highest concentration in the embryo, where it accumulated in the aleurone, scutellum and parts of the endosperm. After exposure of wheat seeds to pure components of essential oils, GC-MS analysis of extracts of the seeds showed the presence of new products in the endosperm and embryo. These were probably formed by the metabolic detoxification of the inhibitors by the seed. Detoxification occurs in both embryo and endosperm during the first day of imbibition. In further work it was found that microtubules of Arabidopsis or wheat seedlings were disrupted within minutes after exposure to very small concentrations of some monoterpenes in the gaseous phase, whereas actin filaments remained intact. In field studies, in which known amounts of residues of aromatic plants were mixed with soil in which wheat seeds sown, some components of the essential oil and their derivatives were found in the seed embryo and endosperm. The agro-technology aspects will be discussed and further work of formulation and application of essential oils as bio-herbicides will be present.

## OC 2. Qualitative and quantitative analysis of Vetiver oils by comprehensive two-dimensional Gas-Chromatography

Filippi J-J<sup>1</sup>, Baldovini N<sup>1</sup>, Belhassen E<sup>1</sup>, Meierhenrich U<sup>1</sup>, Brevard H<sup>2</sup>

<sup>1</sup>LCMBA UMR 6001, Université de Nice-Sophia Antipolis, CNRS, parc Valrose, 06108 Nice Cedex 02, France.

<sup>2</sup>Robertet S.A., 37 avenue Sidi Brahim, 06130 Grasse, France.

Jean-Jacques.Filippi@unice.fr

Keywords: Vetiver; GCxGC; quantitative analysis

Vetiver oil is among the main basic ingredients of every modern perfumer [1]. However, vetiver oil has been lately incorporated in western fragrances, although being known for centuries in India. In the 1960's, many fragrance manufacturers started to include vetiver in perfume compositions and from that time, the worldwide production has increased so as to reach ~300-400t/year nowadays [2]. Since more than 200 constituents have already been described in the literature, mostly consisting of sesquiterpenoid derivatives, vetiver oil is actually considered as one of the most complex essential oils [3]. Indeed, such a complexity has always been a barrier to the fast and direct quantitative analysis. We thus aimed at using GCxGC (with both FID and MS detection) as the main tool for the identification and quantitation of vetiver oils constituents.

The correct understanding of 1D- and 2D-chromatographic data is the first step of an analytical process leading to the identification of odour impact compounds in vetiver extracts.

[1] K. Bauer, et al. (2001) Common fragrance and flavor materials, Wiley-VCH, Weinheim.

[2] M. Maffei (2002) Vetiveria, The Genus Vetiveria., CRC Press.

[3] P. Weyerstahl, et al. (2000) Flavour Fragr. J. 15: 395-412.

### **OC 3. Dealing with phthalic-acid ester (PAE) residues in raw materials and compounded flavourings**

Cachet T, Liddle P

*International Organization of the Flavor Industry (IOFI), Avenue des Arts 6, B- 1210 Brussels, Belgium*

*tcachet@iofiorg.org*

Keywords: Phthalic-acid esters, Phthalates, DEHP, Flavourings, Contaminants

Phthalic-acid esters (PAE, often referred to simply as “phthalates”) are a group of diesters of ortho-phthalic acid.

PAE have become ubiquitous environmental contaminants due to leaching from their widespread applications as plasticisers to soften polyvinyl chloride (PVC) and other widely-used polymeric materials. However, except for the use of diethyl phthalate which is accepted as a carrier for fragrances for use in cosmetics and personal care products, there is NO authorized use of PAE in flavour and fragrance raw materials and compounded flavourings. The illegal use of di-(2-ethylhexyl)phthalate (DEHP) in specific food flavourings in Taiwan uncovered early 2011 has triggered a major food supply chain issue in Asia with massive food recalls, particularly in the P.R. China.

As a consequence of the adulteration issue the China Health Ministry set limits on the presence of DEHP, di-isononyl phthalate (DINP) and di-n-butyl phthalate (DBP) of 1.5 mg/kg (DEHP), 9.0 mg/kg (DINP) and 0.3 mg/kg (DBP) in food and food additives, with no distinction between final food and food ingredients, including flavourings. These limits have subsequently been qualified by another rule specifying that food flavourings, in which the total sum of phthalates does not exceed 60 mg/kg, are permitted for use in production, for sale and applications.

IOFI reacted to this issue by initiating several projects. One project focused on reviewing the presence of PAE, and in particular of DEHP, in food from the use of flavourings. The objective of this effort was to be able to present evidence that the contribution of DEHP to the final food that may be contributed by the flavourings is negligible compared to overall human exposure. The results of this review will be presented.

In another project the IOFI Working Group on Methods of Analysis (WGMA) reviewed the extensive literature on the analysis of PAE in food and food components. From this review the WGMA has developed a set of recommendations for the analysis of PAE in flavourings. These recommendations were used for initiating additional method evaluation work to detect PAE in spiked and unspiked essential oils and extracts with the support of a large number of flavour industry analytical laboratories. Initial results of the method evaluation will be provided and discussed along with recommendations for the reduction of PAE residues in raw materials from natural and synthetic sources and in compounded flavourings.



## OC 4. Kinetics of isolation of essential oils from plants

Sovova H

*Institute of Chemical Process Fundamentals of the ASCR, Czech Republic*

*sovova@icpf.cas.cz*

Keywords: Essential oils, Distillation, Extraction, Kinetics, Plant microstructure

The rate of isolation of essential oils from plants depends on the method used, on operation conditions and on the plant pre-treatment. From chemical engineering point of view, the rate of the process is determined by phase equilibrium, mass transfer resistances, and flow pattern, whatever method is used – steam distillation, hydro-distillation [1], solvent extraction, supercritical fluid extraction [2] or the extraction with subcritical water.

The aim of this contribution was to compare these processes on the basis of their kinetic parameters and with respect to microstructure of plant materials, as significant progress was made recently in modelling the extraction of essential oils from different types of glands visualised by scanning electron microscopy [3–6].

Kinetic data and mathematical models are taken from the literature on different types of distillation and extraction techniques. It is often necessary to distinguish between the easily accessible oil and the oil captured in a low-permeable tissue or to take into account loss of volatile substances during the plant pre-treatment in order to explain different yields of essential oil obtained using different techniques.

The results show how the extraction or distillation rate depends on equilibrium and mass transfer parameters, being controlled first by equilibrium and then by mass transfer resistance. It is also shown that the parameters of geometry and microstructure of a given plant material determined for one of these separation techniques are applicable in the models for other separation techniques so that a prediction of extraction or distillation rate is possible.

Acknowledgements This work was supported by Technology Agency of the Czech Republic (project TA01010578).

1. H Sovova, SA Aleksovski (2006) *Flav. Fragr. J.* 21: 881–889.
2. M Bocevska, H Sovova (2007) *J. Supercrit. Fluids* 40: 360–367.
3. I Zizovic et al. (2005) *Chem. Eng. Sci.* 60: 6747–6756.
4. I Zizovic et al. (2005) *J. Supercrit. Fluids* 39: 338–346.
5. I Zizovic et al. (2005) *J. Supercrit. Fluids* 43: 249–258.
6. M Stamenic et al. (2008) *J. Supercrit. Fluids* 46: 285–292.

## OC 5. Structural factors in the odor of $\alpha$ -santalol derivatives

Hasegawa T<sup>1</sup>, Izumi H<sup>1</sup>, Yamada H<sup>2</sup>

<sup>1</sup>Department of Chemistry, Graduate School of Science and Engineering, Saitama University

<sup>2</sup>Yamada-matsu Co., Ltd.

toshihas@mail.saitama-u.ac.jp

Keywords:  $\alpha$ -Santalol, Structure–odour relation, *E/Z*-isomers, Derivatives

$\alpha$ -Santalol and  $\beta$ -santalol are the main constituents that give sandalwood (*Santalum album* L.) its distinctive woody odour. Other santalol derivatives with a formyl group have been identified as minor components that notably contribute to the sandalwood fragrance [1][2]. Here, we examine how structural factors affect the odour of  $\alpha$ -santalol derivatives by comparing the odour of their *E/Z*-isomers and saturated analogues.  $\alpha$ -Santalol is a sesquiterpene with a polycyclic moiety and a hydroxyl group. These parts were replaced with other structures, and we compared the odours of the *E/Z*-isomers and their saturated analogue for the prepared derivatives. We have previously studied the structure–odour relations of  $\alpha$ -santalols bearing various functional groups (hydroxyl, formyl, formyloxy, and acetoxy), and found a similarity in odour between the *Z*-isomer and its saturated analogue [3]. Next we selected  $\alpha$ -santalols with a benzyl group in place of a hydroxyl group. Many benzyl compounds have strong characteristic odours, suggesting that a benzyl group should strongly affect the odor of these compounds. We found similar odours between the *E*-isomer and its saturated analogue, in contrast to the case of an  $\alpha$ -santalol derivative with a hydroxyl, formyl, formyloxy, or acetoxy group. In addition, we investigated the influence of the bulky polycyclic moiety on the odor of the derivatives. The odour was evaluated when the polycyclic moiety was replaced with a linear alkyl chain. The polycyclic moiety was found to be the most important structural factor in the characteristic sandalwood odour. Through the synthesis of derivatives and the evaluation of their odour, we could identify key structural factors in  $\alpha$ -santalol odour.

1. T Hasegawa et al. (2011) *Flavour Fragr. J.* 26, 98–100.
2. T Hasegawa (2011) *In: S Zereshki (Ed.) Separation of Odour Constituents by Microscale Fractional Bulb-to-Bulb Distillation, Distillation-Advances from Modelling to Application, InTech, Croatia, Chapter 9.*
3. T Hasegawa et al. (2012) *Molecules*, 17, 2259–2270.

## OC 6. Cryptic speciation in liverworts: *Conocephalum conicum*, *Reboulia hemisphaerica* and *Marchantia polymorpha*

Ludwiczuk A<sup>1</sup>, Glowniak K<sup>1</sup>, Asakawa Y<sup>2</sup>

<sup>1</sup>Department of Pharmacognosy with Medicinal Plant Unit, Medical University of Lublin, Poland

<sup>2</sup>Faculty of Pharmaceutical Sciences, Tokushima Bunri University, Japan

aludwiczuk@pharmacognosy.org

Keywords: Volatile components, Liverworts, Cryptic speciation, Chemotypes, GC/MS technique

Cryptic speciation is a concept that has been adopted along with the use of molecular methods in studies of intraspecific phylogenies. It refers to cases where two species or populations are clearly genetically differentiated, but show little or no morphological differences. Cryptic species have been also identified in several liverworts, and these may be either allopatric or sympatric [1,2].

The aim of this work was to apply GC/MS technique as a simple method in identification of cryptic species in liverworts. The volatile components of different collections of three liverwort species; *Conocephalum conicum*, *Reboulia hemisphaerica*, and *Marchantia polymorpha* were analyzed. On the basis of GC/MS data, all analyzed *C. conicum* samples can be divided into 5 chemical groups, which correspond to 5 cryptic species found with use of isozymes and psbA sequencing [3]. The formerly single species, *Reboulia hemisphaerica*, recently was subdivided into several subspecies [4]. The analysis of the volatiles of the Japanese collections of this liverwort shows the occurrence of two very distinct chemotypes. One chemotype is characterized by the presence of cyclomylytayne and chamigrane sesquiterpenoids, and the second one produces mainly gymnomitranes. Studies of enzyme polymorphism of the European colonies of *Marchantia polymorpha* indicated the presence of three subspecies, which can be distinguished mainly on the basis of isozyme and ecological preferences [5]. On the basis of the chemical composition of Tahitian *M. polymorpha*, this liverwort can be divided into 4 chemotypes. Each of the chemotype is characterized by the presence of their own peculiar components, but all of them are identical from morphological point of view.

In conclusion, the analyzed liverwort species are characterized by high diversity of lipophilic terpenoids, especially sesquiterpenoids, and these secondary metabolites can assist in taxonomic differentiation and identification of cryptic species within one liverwort species.

Acknowledgments: The authors gratefully acknowledge Prof. I.J. Odrzykoski (Poland) for *C. conicum* samples and Prof. P. Raharivelomanana and Prof. J.-P. Bianchini (French Polynesia) for *M. polymorpha* collections.

1. AJ Shaw (2001) *J. Biogeogr.* 28: 253-261.
2. H Miwa et al. (2009) *Gene* 441: 169-175.
3. IJ Odrzykoski J Szweykowski (1991) *Pl. Syst. Evol.* 178: 135-151.
4. MC Boisselier-Dubayle et al. (1998) *The Bryologist* 101: 61-69.
5. MC Boisselier-Dubayle et al. (1995) *Taxon* 44: 363-376.

## OC 7. Synergistic biological activity and chemodiversity of Himalayan medicinal plants from Nepal

Satyal P, Paudel P; Setzer WN

*Department of Chemistry, University of Alabama in Huntsville, AL-35899, USA*

*ps0013@uah.edu*

Keywords: Essential oil, Himalayan range, Biological activity, Chemotypes, natural medicines, Volatile chemicals

Nepal is a small country, located between two emerging giants of the Asian continent: India and China. However, like the Himalayan range, Nepal's biodiversity, a mosaic of species, is an unmatched giant in its own right. In coordination with a botanist and a chemist from Tribhuvan University in Nepal, and the Natural Products Group at University of Alabama in Huntsville, around 100 essential oil samples were collected and further analyzed using GC-MS for chemical composition as well as tested for biological activity including cytotoxicity, antimicrobial, brine shrimp lethality, allelopathy, larvicidal, and insecticidal activity. In addition, separate testing of essential oil components was conducted to determine if any single component was responsible for the biological activity recorded in our samples. The results could not attribute any one compound responsible for the observed activities in the samples and we therefore conclude the activities to be a result of synergistic effect amongst the components in the samples.

Aside from biological testing, the result of our systematic and detailed analyses about plant chemical composition has revealed new chemotaxonomic divisions for some of the different species. Plant chemotypes are not only dependent on genetic difference between species but vary among geographical and climatic conditions. This was studied in some cases through comparison of oil compositions of the same species which were collected from different locations.

In conclusion, through international collaboration this project was undertaken with the sole goal of identifying potential biologically active essential oils in Nepalese plants. However our efforts have also translated into a wider understanding of chemodiversity and species differentiation. Through our research we believe that the Himalayan range has the potential to be one of the most important sources for alternate natural medicine.

Acknowledgments: We would also like to express our gratitude to Tilak Gautam for identification of plants and to Dr. Bernhard Vogler for technical assistance.

1. P Satyal et al.(2012) J. Chem. Pharmaceut. Res. 4:437-439.
2. P Satyal et al. (2012) J. Essen. Oil Bearing Plants 15: (Accepted).
3. WN Setzer (2009) Nat. Prod. Commun. 4:1305-1316.
4. P Satyal et al. (2012) Phamacog. Res. 4:(Accepted).

## OC 8. Screening of essential oils for inhibitory effects on the activity of BACE-1, the $\beta$ -secretase involved in Alzheimer's disease

Videira R<sup>1</sup>, Castanheira P<sup>2</sup>; Grãos M<sup>2</sup>; Salgueiro L<sup>3</sup>; Faro C<sup>4</sup>; Cavaleiro C<sup>3</sup>

<sup>1</sup>Center of Pharmaceutical Studies, Faculty of Pharmacy, University of Coimbra, Pólo das Ciências da Saúde, Azinhaga de Santa Comba, 3000-548 Coimbra, Portugal / Biocant - Center of Innovation in Biotechnology, Parque Tecnológico de Cantanhede, Núcleo 04, L;

<sup>2</sup>Biocant - Center of Innovation in Biotechnology, Parque Tecnológico de Cantanhede, Núcleo 04, Lote 2, 3060-197 Cantanhede, Portugal;

<sup>3</sup>Center of Pharmaceutical Studies, Faculty of Pharmacy, University of Coimbra, Pólo das Ciências da Saúde, Azinhaga de Santa Comba, 3000-548 Coimbra, Portugal;

<sup>4</sup>Biocant - Center of Innovation in Biotechnology, Parque Tecnológico de Cantanhede, Núcleo 04, Lote 2, 3060-197 Cantanhede, Portugal / Center for Neuroscience and Cell Biology, University of Coimbra, Coimbra, Portugal

cavaleir@ff.uc.pt

Keywords: BACE-1; cell-permeable inhibitors; *Lavandula luisieri*; essential oil; Alzheimer's disease

One of the most important features in Alzheimer's disease is the generation and deposition of neurotoxic  $\beta$ -amyloid peptide (A $\beta$ ) outside and around brain neurons. The inhibition of BACE-1, a key enzyme in A $\beta$  formation, is pointed as a promising therapeutic alternative for this disease. A crucial attribute for the *in vivo* effectiveness of BACE-1 inhibitors is the ability to crossover blood-brain barrier to reach brain targets. Pharmacokinetic theory points hydrophobicity and low molecular weight as advantageous attributes of drug candidates considering cellular membranes and barriers crossover. Recognizing such physicochemical feature in the ordinary components of essential oils we undertook a screening of essential oils for their ability to inhibit recombinant proBACE-1.

Among dozens of essential oils tested, the oil from *Lavandula luisieri* was signalized as inhibiting the enzyme (IC<sub>50</sub>  $\approx$  120  $\mu$ g.ml<sup>-1</sup>), showing an effect comparable to those of standard inhibitors used for research purposes. Moreover, the oil of *L. luisieri* was tested on the endogenous BACE-1 in cultured CHO cells, being responsible for a reduction in A $\beta$  production (EC<sub>50</sub>  $\approx$  86  $\mu$ g.ml<sup>-1</sup>) with no significant toxicity. The composition of the oil was established by combination of GC and GC-MS data, with the identification of approximately 100 constituents, representing 88.4% of the composition. It is mainly composed of oxygen-containing monoterpenes (62.3%), including 33.0% of necrodane skeleton compounds.

Results prove the effectiveness of the essential oil of *L. luisieri* as inhibitor of BACE-1 activity, both in enzymatic and cellular assays, and supported the bioguided fractioning of the oil towards the isolation of a new BACE-1 inhibitor.

Acknowledgements: Authors are grateful to Fundação para a Ciência e Tecnologia (FCT), POCI 2010/FEDER for financial support. R. Videira was founded through the PhD grant SFRH / BD / 19199 / 2004.

## OC 9. Portuguese Lavenders as a source of anti-inflammatory drugs

Zuzarte M<sup>1</sup>, Gonçalves MJ<sup>1</sup>, Cruz MT<sup>2</sup>, Cavaleiro C<sup>1</sup>, Canhoto J<sup>3</sup>, Salgueiro L<sup>1</sup>

<sup>1</sup>Center of Pharmaceutical Studies, Faculty of Pharmacy, Health Science Campus, University of Coimbra, Azinhaga de S. Comba 3000-354, Coimbra, Portugal, <sup>2</sup>Faculty of Pharmacy and Center for Neuroscience and Cell Biology, University of Coimbra, 3000-295 Coimbra, Portugal, <sup>3</sup>Center for Functional Ecology, Department of Life Sciences, University of Coimbra, Ap. 3046, 3001-401 Coimbra, Portugal

mzuzarte@hotmail.com

Keywords: *Lavandula* spp., Essential oils, Anti-inflammatory, Cytotoxicity, iNOS, COX-2

Inflammatory diseases are a major burden on humanity, despite recent successes with biopharmaceuticals. The lack of responsiveness and drug resistance, delivery problems and the cost of biopharmaceuticals justify the search for effective and less toxic anti-inflammatory agents. As part of our ongoing work on the valorization of Portuguese lavenders [1, 2, 3, 4] we now report their anti-inflammatory potential.

The essential oils were isolated by hydrodistillation and analysed by GC and GC/MS using fused silica capillary columns with two different stationary phases. The main constituents of the oils were carvacrol (42.8%) and *cis*- $\beta$ -ocimene (27.4%) for *L. multifida*; 1,8-cineole (33.9%), fenchone (18.2%), camphor (2.2%) and rare necrodane derivatives for *L. luisieri*; fenchone (42.1%), camphor (17.9%), and 1,8-cineol (10.4%) for *Lavandula pedunculata*; and 1,8-cineol (34.5%), camphor (13.4%),  $\alpha$ -pinene (9.0%) and linalool (7.9%) for *Lavandula viridis*.

To evaluate the anti-inflammatory potential of the oils, the *in vitro* model of lipopolysaccharide (LPS)-stimulated macrophages was used. The following inflammatory parameters were evaluated: nitrite oxide (NO) production by Griess reaction and inducible NO synthase (iNOS) and cyclooxygenase-2 (COX-2) expression by western blot analysis using specific antibodies.

All the oils (0.64  $\mu$ L/mL), excluding *L. multifida* were able to significantly ( $p < 0.001$ ) inhibit NO production without affecting cell viability. *L. luisieri* and *L. viridis* were the most effective, showing an inhibitory activity around 100%. Regarding the inhibition of iNOS and COX-2 expression, *L. viridis* was the most effective, inhibiting completely iNOS expression and COX-2 expression around 50%. *L. luisieri* and *L. pedunculata* were also effective on the inhibition of iNOS expression.

These promising results highlight the potential of Portuguese lavenders as anti-inflammatory drugs, thus justifying further *in vivo* assays to confirm the effectiveness of the *in vitro* results.

Acknowledgments: This work was funded through national funds from FCT – Fundação para a Ciência e a Tecnologia under the project CEF/POCI2010/FEDER and a PhD fellowship to Mónica R. Zuzarte (SFRH/BD/40218/2007).

1. M Zuzarte et al. (2009) Chem. Biodivers. 6: 1283-1292.
2. M Zuzarte et al. (2010) Ind. Crop. Prod. 32: 580-587.
3. M Zuzarte et al. (2010) J. Med. Microbiol. 60: 612-618.
4. M Zuzarte et al. (2011) Eur. J. Clin. Microbiol. Infect. Dis. DOI 10.1007/s10096-011-1450-4.



## Young Scientists (YS) Oral Communications Abstracts







## YSOC 1. Effects of PSBS gene silencing by RNAi on volatile compound biosynthesis and accumulation in tomato (*Solanum lycopersicum*)

Occhipinti A<sup>1</sup>, Ballottari M<sup>2</sup>, Termolino P<sup>3</sup>, Di Dato F<sup>3</sup>, Cammareri M<sup>3</sup>, Bossi S<sup>1</sup>, Maffei M<sup>1</sup>, Bassi R<sup>2</sup>

<sup>1</sup>Department of Life Sciences and Systems Biology, University of Turin, 10135 Italy

<sup>2</sup>Department of Biotechnologies, University of Verona, 37134 Italy; <sup>3</sup>Institute of Plant Genetics – CNR, Research Division Portici, 80055 Italy

andrea.occhipinti@unito.it

Keywords: Gene silencing, Solanaceae, Nonphotochemical quenching, GC-MS, qPCR, Terpene synthases

Photosynthetic organisms convert light energy into chemical energy, however, high light irradiation can lead to the production of Reactive Oxygen Species (ROS) if the light energy is not dissipated safely. ROS burst can damage cell membranes (photoinhibition). In order to reduce photoinhibition, nonphotochemical quenching (NPQ) is the main regulated mechanism used by plants to dissipate the excess of light energy. NPQ in higher plants is strictly dependent on the presence of a Lhc (Light Harvesting Complex)-like protein, PsbS which does not bind pigments, unlike the other Lhc proteins, and stimulates NPQ interacting with Photosystem II antenna proteins. This mechanism is very efficient, inducing thermal dissipation up to 80% of absorbed light even at moderate irradiance, suggesting that evolution lead higher plants to fully protect themselves at the expense of partial a loss of light energy conversion efficiency. In this work, we produced different transformed lines of *Solanum lycopersicum* with RNA interference (RNAi) construct in order to silence *PsbS* gene. Here we report the effects on biosynthesis and accumulation of volatile compounds (VOCs) in mutant plants. Leaf and fruit tissue extracts were analyzed by GC-MS, whereas quantitative gene expression analysis was carried out through qPCR for genes involved in terpene biosynthesis.

The metabolic and biochemical analyses revealed an increased accumulation in *psbs* mutants of some monoterpenes. The results suggest that deletion of this photoprotective mechanism might have an industrial application for plants cultivated in control conditions as in greenhouses, where the light can be artificially modulated without impairing the growth and fruit production, but increased accumulation of volatiles compounds that should influence fruit flavour and increase the resistance to herbivores.

## YSOC 2. Coriander oil antimicrobial activity: A flow cytometric study

Silva E, Ferreira S, Duarte A, Queiroz JA, Domingues FC

CICS-UBI – Health Sciences Research Centre, University of Beira Interior, Portugal

*filomena@fcsaude.ubi.pt*

Keywords: Coriander oil, Antibacterial activity, Antifungal activity, Flow cytometry, Germ tube, Synergism

The aim of this work was to study the antimicrobial effect and mechanism of action of coriander essential oil against Gram-positive, Gram-negative bacteria and *Candida* yeast cells.

Antibacterial and antifungal susceptibilities were evaluated using classical microbiological techniques with MIC and MLC determination. The effect of coriander oil on respiratory activity, membrane polarization, efflux activity, cell cycle progression and membrane permeability was evaluated by flow cytometry. Furthermore, in *Candida* cells, the coriander oil effect on germ tube formation and the potential synergism of coriander oil with amphotericin B were also studied.

Our results showed that coriander oil has an effective antibacterial activity against all bacteria tested with a potent bactericidal activity against almost all bacteria tested, with the exception of *B. cereus* and *E. faecalis* [1]. For fungal cells, coriander oil also revealed a fungicidal activity against the *Candida* strains tested with MLC values equal to the MIC value [2]. Propidium iodide (PI) incorporation and concomitant loss of all other cellular functions such as efflux activity, respiratory activity and membrane potential seem to suggest that the primary mechanism of action of coriander oil on both Gram-positive and Gram-negative bacteria as well as in *Candida* cells is membrane damage, which leads to cell death [1,2]. Also, concentrations below the MIC value caused a marked reduction in the percentage of germ tube formation for *C. albicans* strains with a synergistic effect between coriander oil and amphotericin B also observed for *C. albicans* strains [2].

These results, showing a potent antibacterial and antifungal activity due to membrane permeability, are noteworthy and justify the use of this plant not only as a food flavouring agent, but also as a food preservative and even in medical devices in order to prevent fungal infections.

Acknowledgments: This study was supported by grants SFRH/BD/41521/2007 and SFRH/BD/66857/2009 from the Portuguese Foundation of Science and Technology.

1. F Silva et al. (2011) J. Med. Microbiol. 60: 1479-86.
2. F. Silva et al. (2011) Phytomed. 19: 42-7.

### **YSOC 3. Application of vibrational spectroscopy and chemometrics as a rapid and low cost alternative in the quality assessment of indigenous South African essential oils**

Sandasi M<sup>1</sup>, Viljoen AM<sup>1</sup>, Baranska M<sup>2</sup>

<sup>1</sup>*Department of Pharmaceutical Sciences, Tshwane University of Technology, Private Bag X680, Pretoria, South Africa*

<sup>2</sup>*Faculty of Chemistry, Jagiellonian University, 3 Ingardena 30-060, Krakow, Poland*

[sandasim@tut.ac.za](mailto:sandasim@tut.ac.za)

Keywords: Vibrational spectroscopy, Chemometrics, Buchu, Sage, Lippia, Cape chamomile

As the market for herbal products continues to grow, concerns over the safety of some products has been raised. Adulteration especially in essential oils has become a point of great concern which compromises the quality of the product and ultimately the safety of the consumer. Current analytical methods are based on the identification and quantification of marker constituents which in some cases may not be biologically active. Due to the complexity and variation in chemical composition of these mixtures, it is important to profile the whole metabolome. Spectroscopic techniques that include mid- and near- infrared spectroscopy (MIR and NIR) have been identified as potential quality control alternatives.

Four commercially important essential oils from South Africa (Buchu, Cape chamomile, Sage and *Lippia* oils) were used to demonstrate the application of vibrational spectroscopy and chemometrics in multivariate classification and class predictions. Additionally calibration models were developed for marker constituents that would be used to predict these constituents in future samples.

Plant material was obtained from cultivation sites and the wild. The essential oils were isolated by hydrodistillation and reference analysis performed using GC-MS. The oils were also scanned on MIR and NIR spectrometers for spectral acquisition. For identification and classification, orthogonal projections to latent structures (OPLS-DA) were employed and the models built were used to further predict the botanical origin of commercial oils. Partial least squares regression (PLSR) analysis was used to develop multivariate calibration models that were also used to predict the quantities of marker constituents in future samples.

The results showed that OPLS-DA is a useful tool in the differentiation of essential oils from different species. Class prediction of unknown oil samples was possible with high predictive power. The PLS calibration models developed showed the best model performance with very high correlation coefficients of > 90% and low errors of prediction for marker constituents. The feasibility of using vibrational spectroscopy as an alternative quality assessment technique has been demonstrated.

Acknowledgement: The authors would like to thank the National research Foundation (NRF) of South Africa and Tshwane University of Technology for financial support.

#### YSOC 4. Effect of hydroxypropyl- $\beta$ -cyclodextrins on the antioxidant activity of *Lavandula* essential oils

Costa P, Gonçalves S; Romano A

*Institute for Biotechnology and Bioengineering, Centre of Genomics and Biotechnology (IBB-CGB),  
Faculty of Sciences and Technology, University of Algarve, Campus de Gambelas, Ed. 8, 8005-  
139 Faro, Portugal*

a26166@ualg.pt

Keywords: Antioxidant activity, Cyclodextrins, Essential oils, ORAC assay

Excess free radicals are produced under stress and in response to pathogens, often resulting in oxidative damage to lipids, proteins and nucleic acids. In humans, this damage is implicated in ageing and in numerous diseases [1]. The ability of natural antioxidants to protect cells from oxidative stress has been previously demonstrated [2] and several methods have been proposed to evaluate the antioxidant activity of natural products [3]. However, many assays are conducted in an aqueous system and, therefore, must be adapted to measure lipophilic natural products, such as essential oils. Cyclodextrins are a group of naturally occurring cyclic oligosaccharides with a lipophilic central cavity and a hydrophilic outer surface [4], which have been used as solubilizing agents for hydrophobic volatile oils [5]. In this work, the antioxidant activities of the essential oils from *Lavandula viridis* L'Hér and *Lavandula pedunculata* subsp. *lusitanica* (Chaytor) Franco, endemic species from Portugal (Algarve region), were assessed in the absence and presence of hydroxypropyl- $\beta$ -cyclodextrins (HP- $\beta$ -CD) by means of the oxygen radical absorbance capacity (ORAC) assay [6]. The results showed that the peroxy radical scavenging activity increased in the presence of HP- $\beta$ -CD in both species, however the *L. viridis* essential oil exhibited the highest ORAC values. The results obtained demonstrated that the addition of HP- $\beta$ -CD improves the solubility of *Lavandula* essential oils and, consequently, their chain-breaking antioxidant activities.

Acknowledgements: C. Patrícia, S. Gonçalves acknowledge a grant from Portuguese Science and Technology Foundation (FCT, SFRH/BD/63505/2009 and SFRH/BPD/31534/2006).

- [1] M Valko et al. (2007). *Int. J. Biochem. Cell Biol.* 39: 44-84
- [2] CD Stalikas (2007) *J. Sep. Sci.* 30: 3268-3295
- [3] RL Prior et al. (2005). *J. Agric. Food Chem.* 53: 4290-4302
- [4] T Loftsson et al. (2005). *Expert Opin. Drug Deliv.* 2: 335-351
- [5] HMC Marques (2010). *Flavour Fragr. J.* 25: 313-326
- [6] KM Gillespie et al. (2007). *Nat. Protoc.* 2: 867-870.

## YSOC 5. Classification of Lavender essential oils for the evaluation of sedative effects on Human

Kitao M<sup>1</sup>, Tomi K<sup>1</sup>; Murakami H<sup>2</sup>; Matsumura Y<sup>2</sup>; Hayashi T<sup>1</sup>

<sup>1</sup>Faculty of Agriculture, Kinki University

<sup>2</sup>Graduate School of Agriculture, Kyoto University

1133650004n@nara.kindai.ac.jp

Keywords: Lavender, Principal Component Analysis, Aroma component composition, Enantioselective GC-MS, Optical isomer

Lavender essential oils are used for aromatherapy. Linalool is the main aroma component and thought to have a sedative effect on human [1]. The other components must affect the autonomic nervous system and then the whole effect of an essential oil may be determined by the component composition. Two steps are important for aromatherapy: the categorization of essential oils and the proper use based on scientific evaluation. In this study, lavender essential oils were categorized by principal component analysis of aroma component compositions.

Twenty-seven types of lavender essential oils, which include several *Lavandula* species, were purchased in markets in Japan. These essential oils were analyzed by GC-MS. Furthermore, optical isomer compositions of some aroma components were also determined using enantioselective GC-MS. Principal component analysis was conducted based on the data obtained by GC-MS.

High amount of linalool was commonly found in the lavender essential oils (20-50%) except *L. stoechas*. Linalyl acetate was also predominantly contained but not in *L. stoechas* and *L. spica*. *L. angustifolia* oils showed lower camphor content. Optical resolution analysis revealed the (R) form optical isomers are predominant in most of the essential oils, although (S) formula of borneol was abundant in some essential oils. The essential oils tested in this experiment could be divided into several groups by principal component analysis according to the PC1 and PC2. Loading factors indicated that the components which mainly contributed to the PC1 were camphor, alpha-pinene and eucalyptol.

From the chemical profile analysis of lavender essential oils, some oils including *L. stoechas* were selected according to their unique compositions. Principal component analysis results might make it possible to reveal the relationships between the compositions and sedative effects on human.

1. K Kuroda et al. (2005) Eur. J. Appl. Physiol. 95:107-114

**YSOC 6. Essential oils do not inhibit *Escherichia coli* peptidyl-tRNA hydrolase**

Thompson RM, McFeeters H, Setzer WN, McFeeters RL

Department of Chemistry, University of Alabama in Huntsville, USA

rmt0002@uah.edu

Keywords: Peptidyl-tRNA hydrolase, Essential oil inhibition, *Escherichia coli* Pth

The purpose of the study was to purify peptidyl-tRNA hydrolase (Pth) and peptidyl-tRNA from bacterial cells in order to determine the inhibitory activity of essential oils from aromatic medicinal plants located in Nigeria, South Africa, and Tajikistan. *E. coli* Pth was expressed and purified using metal chelation chromatography and gel shift assays were used to determine the inhibitory factor of the essential oils against Pth. Throughout all assays, the peptide was cleaved off from peptidyl-tRNA. Thus, enzymatic activity was observed from Pth when incubated with each essential oil and peptidyl-tRNAs in solution. As a result, essential oils show no sign of inhibitory activity against Pth. These findings are beneficial for future antibacterial research because they prevent further time and money exhausted and allows for the focus on other plant extracts that display inhibitory activity against Pth. The plant extracts that show inhibitory activity were not essential oils. Future studies include analysis by gas chromatography and mass spectrophotometry in order to determine what compound is responsible for inhibition. Plant extracts show great promise for the future development of antibacterials.

Acknowledgments: I greatly appreciate the McFeeters lab group in the Biochemistry department at The University of Alabama in Huntsville. Also, thanks to the Research Experience for Undergraduates Program at the University of Alabama Huntsville as well as Alabama Space Grant Consortium, President's office of the University of Alabama in Huntsville, and Funds from the Department of Chemistry Patent account.

## Posters Abstracts







## P 1. Chemical composition, antifungal activity and cytotoxicity of *Otanthus maritimus* (L.) Hoffmanns. & link from Portugal

Cabral C<sup>1</sup>; Cavaleiro C<sup>1</sup>; Gonçalves MJ<sup>1</sup>; Cruz MT<sup>2</sup>; Lopes MC<sup>2</sup>; Salgueiro L<sup>1</sup>

<sup>1</sup>Grupo de Bromatologia, Farmacognosia e Ciências Analíticas, Faculdade de Farmácia/Centre of Pharmaceutical Studies, Pólo das Ciências da Saúde, Azinhaga de Santa Comba, Universidade de Coimbra, 3000-548 Coimbra, Portugal

<sup>2</sup>Centro de Neurociências e Biologia Celular e Faculdade de Farmácia, Universidade de Coimbra, 3000-295 Coimbra, Portugal

cmdscabral@gmail.com

Keywords: *Otanthus maritimus*; essential oil; GC-MS; antifungal activity; cytotoxicity

*Otanthus maritimus* (L.) Hoffmanns. & Link belongs to the Asteraceae family that includes well-known aromatic and medicinal species such as chamomile, yarrow, wormwood and feverfew [1]. The use of *O. maritimus* has been reported in traditional medicine for various purposes including the treatment of toothache, asthmatic bronchitis, dysentery and inflammation of the urinary bladder [2-4]. The present work aimed to characterize the chemical composition of the essential oils obtained from the aerial parts of *O. maritimus* collected in Portugal, as well as to evaluate the antifungal activity of the oil at concentrations without skin cytotoxicity.

The oils were isolated by hydrodistillation and analyzed by gas chromatography (GC) and gas chromatography-mass spectrometry (GC-MS). Broth macrodilution methods based on the Clinical and Laboratory Standards Institute (CLSI) reference protocols were used to determine minimum inhibitory concentrations (MICs) and minimum lethal concentrations (MLCs) of the essential oils. Cytotoxicity was tested in HaCaT keratinocytes through the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay.

The main compounds of the oils are chrysanthenone (40.4-57.2%), filifolone (12.2-15.5%), *cis*-chrysanthenyl acetate (10.1-12.2%) and  $\alpha$ -pinene (6.7-7.2%). Results concerning the antifungal activity demonstrated the potential of *Otanthus* oil against dermatophytes. Furthermore, evaluation of cell viability showed no significant cytotoxicity in HaCaT keratinocytes at concentrations between 0.16 and 0.32  $\mu$ L/mL.

These findings add significant information to the knowledge on the pharmacological activity of *O. maritimus* essential oils, specifically due to its antifungal properties. This evidence of the therapeutic effects of *O. maritimus* essential oil, suggest that doses with antifungal activity with very low detrimental effect on keratinocytes combined with its pleasant odor could be of great value to cosmetic and pharmaceutical industries.

Acknowledgements. Our thanks to CEF/POCI2010/FEDER, for financial support.

1. TG Tutin (1976) In: TG Tutin *et al.* (Eds.) *Flora Europaea* 4. Cambridge University Press, Cambridge, UK, pp. 168.
2. L Reutter (1923) *Traité de matière médicale et de chimie végétale*. Paris Librairie J. B. Baillière et Fils, Paris, France.
3. CA Thanos, K Georghiou, DJ Douma, CJ Marangaki (1991) *Anal. Bot.* 68: 469-475.
4. M Tsoukatou, C Vagias, C Harvala, V Roussis (2000) *J. Essent. Oil Res.* 12: 360-364.

## P 2. Physical and chemical characterization of inclusion complexes: essential oil from *Ocimum basilicum* and $\beta$ -cyclodextrin

Serafini MR<sup>1</sup>; Menezes PP<sup>2</sup>; Nascimento SS<sup>1</sup>; Quintans-Junior LJ<sup>1</sup>; Almeida EDP<sup>1</sup>; da Silva GF<sup>1</sup>; Grangeiro-Junior S<sup>3</sup>; Lima AAN<sup>4</sup>; Araujo AAS<sup>1</sup>

<sup>1</sup>Federal University of Sergipe- Brazil

<sup>2</sup>Universidade Federal de Sergipe- Brazil

<sup>3</sup>Pharmaceutical Laboratory of Pernambuco- Brazil

<sup>4</sup>Federal University of Amazônia- Brazil

maiserafini@hotmail.com

Keywords: inclusion complexes,  $\beta$ -cyclodextrin, *Ocimum basilicum*

Several *Ocimum* species (Lamiaceae) are used to treat central nervous system (CNS) disorders in various parts of the world and its anticonvulsant activity is frequently reported [1,2]. *Ocimum basilicum* L. essential oil (OBEO) is rich in monoterpenes, such as  $\delta$ -cadinol (10.2%), estragole (22.6%), and linalool (47.3%) [3]. Thus, essential oils have shown potential biology activity, but its insolubility makes it difficult to use in therapeutic systems. Accordingly, this study aimed to characterize the physical-chemical properties and assess the thermal properties of inclusion complexes of *Ocimum basilicum* with  $\beta$ -cyclodextrin ( $\beta$ -CD). Inclusion complexes were prepared by three different procedures. A physical mixture (PM) was prepared by the addition of OBEO to a powdered  $\beta$ -CD under manual agitation. Paste complexation (PC) was carried out by homogenization of  $\beta$ -CD with water (1:1 molar, 10 mL) under constant manual agitation. Then, the material was dried at room temperature (in a desiccator) till the formation of a glass film, which was removed by manual trituration and stored in airtight glass containers. Co-evaporated (CE) was carried out by the addition of water to a beaker containing  $\beta$ -CD (1:1 molar, 20 mL) and stirred for 36 h by a magnetic stirring device operating at 400 rpm. DSC curves can be observed that the preparations PM, PC and CE showed three events endothermic (25-400°C), followed by elimination of the carbonaceous material. The thermoanalytical curve of PM was a superposition of the guest and host curves, which indicates a lower evidence of inclusion and significant interaction between the host and guest molecules. On the other hand, the thermal curves of PC and CE showed in the interval from 130 to 275°C a gradual mass loss attributed to oil release. The FTIR spectra also suggest the complexation, since the bands for the oil were present also in three different preparations. Further studies should be performed to confirm these results.

1. LJ Quintans-Júnior, JRGS Almeida, JT Lima, et al. (2008). Plants with anticonvulsant properties - a review. Rev Bras Farmacogn 18: 798-819.
2. JS Oliveira, LA Porto, CS Estevam, et al. (2009). Phytochemical screening and anticonvulsant property of *Ocimum basilicum* leaf essential oil. BLACPMA 8: 195-202.
3. M Mazutti, B Beledelli, AJ Mossi, et al. (2006). Caracterização química de extratos de *Ocimum basilicum* L. obtidos através de extração com CO<sub>2</sub> a altas pressões. Quim Nova 29: 1198-1202.

### P 3. Antimicrobial and antioxidant activities of volatile oils of three species of Piperaceae grown in Cuba

Saucedo Y<sup>1</sup>; Jorge E<sup>1</sup>; Ramos GR<sup>2</sup>; van der Heyden Y<sup>3</sup>; Monteagudo U<sup>1</sup>; Simó EF<sup>2</sup>; Lerma MJ<sup>4</sup>; Bravo LR<sup>1</sup>; Medinilla M<sup>1</sup>; Herrero JM<sup>2</sup>

<sup>1</sup>Pharmacy Department, Faculty of Chemistry, Central University "Marta Abreu" of Las Villas, C-54830 Santa Clara, Cuba

<sup>2</sup>Department of Analytical Chemistry, Faculty of Chemistry, University of Valencia, E-46100 Burjassot (Valencia), Spain

<sup>3</sup>Department of Analytical Chemistry and Pharmaceutical Technology, Faculty of Medicine and Pharmacy, Vrije Universiteit Brussel-VUB, Belgium

<sup>4</sup>Departamento de Química Analítica y Tecnología de los Alimentos, Universidad de Castilla – La Mancha, Avda. Camilo José Cela 10, E-13004, Ciudad Real, Spain

ysaucedo@uclv.edu.cu

Keywords: *Piper aduncum*, *Piper auritum*, *Piper angustifolium*; essential oil; antioxidant activity; antimicrobial activity.

The Piperaceae family is recognized in the traditional medicine in all America for the use of its oils as astringent, digestive, diuretic and antimicrobial [1-4]. The present study has as purpose evaluate the antimicrobial and antioxidant activities by different *in vitro* techniques of volatile oils obtained by hydrodistillation of dried leaves of three species of Piperaceas grown in Cuba.

The antioxidant activity was evaluated by several assays such as free radical DPPH activity, reducing power and complementary antioxidant activity in linoleic acid system (FTC method). The antioxidant activity of these essential oils was also evaluated against unrefined crude *Cucurbita* seed oil by peroxide, thiobarbituric acid and *p*-anisidine methods.

The essential oil *Piper auritum* showed antioxidant activity similar to natural antioxidants in some cases and better results than *Piper angustifolium* and *Piper aduncum* oils. The *Piper auritum* showed a stronger antioxidant activity than BHA and BHT but lower than that of propyl gallate. Moreover, this antioxidant activity was supported by the complementary antioxidant assay in linoleic acid system and 2, 2'-diphenyl-1-picrylhydrazyl. In other hand the essential oils showed good to moderate inhibitory effects against three microorganisms evaluated, *Klebsiella pneumoniae* ATCC 13833, *Bacillus subtilis* ATCC 6633 and *Candida albicans* ATCC 18231 but these didn't had activity with *Staphylococcus aureus* ATCC 25932 and *Pseudomonas aeruginosa* ATCC 27853.

This study gives a wide view on the composition of Piperaceas essential oils produced during the entire productive season, useful to identify quality parameters for the analytical characterization of this product.

1. J Cicció and C Ballester. (1997). International Journal of Tropical Biology and Conservation. 15 (2): 783-790.
2. Y Sanchez et al. (2002). Rev Cubana Invest Biomed. 21: 214- 216.
3. E. Andrade et al (2008). Chem Biodivers, 5: 97 - 208.
4. L.G Ranilla et al. (2010). Bioresour Technol. 101(12): 4676-89.

#### P 4. Antimicrobial activity of essential oils from *Lavandula luisieri* and cineol against *Botrytis cinerea*

Graça SFN<sup>1</sup>, Delgado FMG<sup>2</sup>, Silva LA<sup>3</sup>, Ferreira MASS<sup>4</sup>, Pintado CMBS<sup>2</sup>

<sup>1</sup>Escola Superior Agrária do Instituto Politécnico de Castelo Branco, Quinta da Senhora de Mércules, Apartado 119, 6001-909 Castelo Branco, Portugal

<sup>2</sup>CERNAS/IPCB, Centro de Estudos de Recursos Naturais, Ambiente e Sociedade/Instituto Politécnico de Castelo Branco, Quinta da Senhora de Mércules, Apartado 119, 6001-909 Castelo Branco, Portugal

<sup>3</sup>Universidade da Beira Interior, Rua Marquês d'Ávila e Bolama, 6201-001 Covilhã, Portugal

<sup>4</sup>CBAA/ Instituto Superior de Agronomia, Universidade Técnica de Lisboa, Edifício Ferreira Lapa, Tapada da Ajuda, 1349-017 Lisboa, Portugal

*fdelgado@ipcb.pt*

Keywords: Essential oils; *Botrytis cinerea*; food spoilage; *Lavandula luisieri*; antimicrobial activity;

The microbiological contamination of strawberry, caused by filamentous fungi, leads to high fruit stock losses during his transport and storage period. The use of edible films with essential oils from *Lavandula luisieri* may be a powerful solution. So, the main goal of this work was to test the antifungal activity of essential oils (EO) of *L. luisieri* and cineol against *Botrytis cinerea*, a common spoilage mold of strawberry.

In order to test the antifungal activity of essential oils against *Botrytis cinerea* the minimum inhibitory concentration (MIC) and the minimum fungicidal concentration (MFC) were determined, respectively, by the broth microdilution method [1] and by the drop plate method onto solid medium [2]. The suspensions,  $0.4 \times 10^4$ - $5.0 \times 10^4$  spores/ml, were performed according NCCLS [3] and Alizadeh-Salteh [4].

The MIC values of essential oils against *Botrytis cinerea* were between 0,38% (v/v) and 1,5% (v/v) and the MFC values were between 1,25% (v/v) and 3% (v/v). Cineol was less effective in mold inhibition than the essential oils of *L. luisieri*.

The results showed an effective antifungal activity of essential oils of *L. luisieri*.

Acknowledgements: CERNAS/IPC is supported by FCT, *Fundação para a Ciência e a Tecnologia* (Science and Technology National Foundation)

1. JH Jorgensen, MJ Ferraro (2009) Medical Microbiology. 49: 1749-1754.

2. V Tullio *et al.* (2007) Journal of Applied Microbiology. 102: 1544-1550.

3. NCCLS (2002) Document M38-A. Wayne, Pennsylvania, USA.

4. S Alizadeh-Salteh *et al* (2010). Europ. J. Hort. Sci. 75(6): 278-282.

## P 5. Morphological and essential oil variability of Portuguese fennel (*Foeniculum vulgare* Mill.)

Lopes VR<sup>1</sup>; Barata AM<sup>1</sup>; Bettencourt E<sup>2</sup>; Pedro LG<sup>3</sup>; Barroso JG<sup>3</sup>; Figueiredo AC<sup>3</sup>

<sup>1</sup>Institutions: <sup>1</sup>Banco Português de Germoplasma Vegetal, UIRGEMP, INRB, I.P., Quinta de S. José, S. Pedro de Merelim, 4700 859, Braga, Portugal

<sup>2</sup>Genetic Resources, Ecophysiology and Plant Breeding Unit, INRB, I.P. (currently on leave of absence)

<sup>3</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portugal

violeta.lopes@inrb.pt

Keywords: *Foeniculum vulgare*; Apiaceae; Portugal; descriptors; essential oils; morphological characterization

Previous studies on Portuguese fennels (*Foeniculum vulgare* Mill.) essential oils [1-3], showed the existence of chemotypes (anethole, anethole / fenchone, anethole / methyl chavicol) that diverge from international standards [4,5]. In the present work, twenty accessions of *F. vulgare* consisting of wild material and fruits' progeny were assessed for diversity through a combined morphochemical evaluation.

The accessions, collected in 2005 and 2006, were morphologically characterized using 34 qualitative and quantitative variables scored on 20 randomly selected plants per population [3]. The essential oils were isolated by hydrodistillation from the fruits (seeds) progeny and analysed by Gas Chromatography (GC) and Gas Chromatography-Mass Spectrometry (GC-MS). The percentage composition of the essential oils was used to determine the relationship between the different oil samples by cluster analysis as in [3].

Essential oils analysis defined two main clusters: cluster I, composed of 9 samples, included *trans*-anethole (43-70%) and fenchone (18-33%) rich oils, and cluster II, with 11 samples, included methyl chavicol (= estragole) (15-44%), *trans*-anethole (14-37%) and fenchone (19-39%) rich oils.

Combined morphological and essential oil assessment, showed that the *trans*-anethole dominant cluster I included plants with larger umbels, larger leaves and higher number of umbels per plant. Accessions whose essential oils gathered in cluster II were morphologically characterized by being taller plants, with higher stems number, larger petiole length and distance between 1st and 2nd leaf segment pairs, small umbels with lower number of rays/umbel and short leaflets.

Two subspecies of *F. vulgare* Mill. (= *F. officinale* All.) occur in Portugal, *F. vulgare* Mill. subsp. *capillaceum* (= *F. vulgare* Mill. subsp. *vulgare*) and *F. vulgare* Mill. subsp. *piperitum* [references in 3]. The latter subspecies is differentiated from subsp. *capillaceum* by smaller umbels with lower number of rays/umbel. The chemical analysis allowed a taxonomical clarification between the two subspecies, despite their morphological similarity.

1. OR Roque, A Proença da Cunha (1989). Bol. Fac. Far. Coimbra 13: 45-52.

2. CMF Cavaleiro et al. (1993) J. Essent. Oil Res. 5: 223-225.

3. VR Lopes et al. (2010) Acta Horticulturae 860: 33-50.

4. Council of Europe (COE) (2008) *European Pharmacopoeia* 6th Edition. Strasbourg.

5. ISO 17412:2007 (2007)

**P 6. Anti-hypernociceptive activity of citronellol, a monoterpene presents in lemongrass essential oil, involves periaqueductal gray and piriform cortex**

Quintans-Júnior LJ<sup>1</sup>; Brito RG<sup>2</sup>; Prado DS<sup>2</sup>; Santos PL<sup>2</sup>; Feitosa VLC<sup>2</sup>; Sousa DP<sup>2</sup>; Araujo AAS<sup>2</sup>; Santos MRV<sup>2</sup>; Quintans JSS<sup>2</sup>; Lucca-Júnior W<sup>2</sup>

<sup>1</sup>Federal University of Sergipe – Brazil

<sup>2</sup>Federal University of Sergipe

lucindojr@gmail.com

Keywords: Monoterpenes, citronellol, hypernociception, pain

Citronellol (CT) is a monoterpene present in lemongrass essential oil. We investigated the anti-hypernociceptive and central nervous system (CNS) effects of CT on rodents.

Male Swiss mice (n=6/group) were treated with CT (25, 50 or 100 mg/kg; i.p.) or saline, 0.5h after the plantar injection of carrageenan (CG; 300 µg/paw), TNF-α (100 pg/paw), prostaglandin (PGE<sub>2</sub>; 100 ng/paw) or dopamine (DA; 30 µg/paw). The hypernociception was evaluated at times of 0.5, 1, 2, and 3h after the injection of agents, using the digital analgesymeter (von Frey). To evaluate the central effects, the animals were perfused 90 min after the treatment, the brains were removed, frozen, cut (20 µm) and subjected to immunofluorescence for Fos. The protocols were approved by Animal Ethics Committee at the UFS (CEPA/UFS: 72/11). The data were analyzed by ANOVA (one-way) followed by Tukey's test (p<0.05).

CT significantly (p<0.001) reversed the hypernociception induced by CG at 100 mg/kg (0.5h: 5.9±0.6, 1h: 6.3±0.6, 2h: 6.4±0.6, 3h: 6.2±0.4) comparing with vehicle group (0.5h: 1.5±0.1, 1h: 2.1±0.2, 2h: 1.9±0.2, 3h: 1.9±0.2). The hypernociception induced by TNF-α was significantly (p<0.001) reversed at 25 (0.5h: 5.0±0.2, 1h: 5.0±0.3, 2h: 4.8±0.4, 3h: 5.9±0.2), 50 (0.5h: 4.6±0.5, 1h: 5.0±0.4, 2h: 5.1±0.5, 3h: 5.6±0.5) and 100 (0.5h: 4.7±0.2, 1h: 4.9±0.3, 2h: 4.9±0.2, 3h: 4.7±0.3) comparing with vehicle (0.5h: 1.9±0.4, 1h: 2.0±0.4, 2h: 1.8±0.3, 3h: 2.1±0.3). The hypernociception induced by PGE<sub>2</sub> was also significantly (p<0.001) reversed at 25 (0.5h: 3.7±0.3, 1h: 3.8±0.3, 2h: 4.6±0.4, 3h: 4.1±0.2), 50 (0.5h: 4.2±0.3, 1h: 4.8±0.1, 2h: 4.7±0.3, 3h: 4.6±0.4) and 100 (0.5h: 4.4±0.3, 1h: 4.8±0.2, 2h: 5.1±0.1, 3h: 5.2±0.3) when compared with vehicle (0.5h: 1.9±0.2, 1h: 1.8±0.05, 2h: 1.9±0.2, 3h: 1.7±0.2). The hypernociception induced by DA was significantly (p<0.001) reversed at 25 (2h: 5.4±0.4, 3h: 5.0±0.2) and 100 (3h: 5.4±0.1) comparing with vehicle (0.5h: 3.4±0.2, 1h: 3.4±0.2, 2h: 2.6±0.3, 3h: 3.2±0.1). The immunofluorescence assay demonstrated a significant increase in the number of Fos positive cells in the piriform cortex (25: 8±1.4, 50: 23.7±4.3, 100: 24±4.8, vehicle: 1.2±0.2) and in the periaqueductal gray (25: 12.2±1.2, 50: 25.7±2.8, 100: 26±3.4, vehicle: 1.5±0.8).

Our results suggest an anti-hypernociceptive effect of CT, probably by the activation of the piriform cortex and periaqueductal gray CNS regions.

Acknowledgments: CAPES, CNPq, FAPITEC-SE.

## P 7. Citronellol, a monoterpene alcohol, reduces the orofacial nociceptive behavior in mice

Quintans-Júnior LJ<sup>1</sup>; Brito RG<sup>1</sup>; Prado DS<sup>1</sup>; Santana MT<sup>1</sup>; Lucca-Júnior W<sup>1</sup>; Quintans JSS<sup>1</sup>; Bonjardim LR<sup>1</sup>; Almeida JRGS<sup>2</sup>; Barreto RSS<sup>1</sup>; Santos MRV<sup>1</sup>

<sup>1</sup>Federal University of Sergipe – Brazil

<sup>2</sup>Federal University of São Francisco Valley - Brazil

lucindojr@gmail.com

Keywords: Monoterpenes, citronellol, orofacial pain, analgesic

Citronellol (CT) is a monoterpenoid alcohol present in the essential oil of many plants, as the *Cymbopogon citratus*. We evaluated the effects of CT on orofacial nociceptive behavior in mice.

Male Swiss mice (n=6/group) were treated intraperitoneally with CT (25, 50 or 100 mg/kg) or saline (control). Orofacial nociception was induced by injection (s.c.) of 20 ul of 2% formalin, 20 ul (2.5 ug) of capsaicin or 40 ul of glutamate (25 uM), 0.5 h before the treatment. The quantification of nociception was performed by measuring the time (seconds) that the animal spent face-rubbing in the injected area with its fore or hindpaws. In the formalin test, the animals were observed at periods of 0 to 5 minutes (first phase) and 15 to 40 minutes (second phase) after formalin injection, in the capsaicin test, at periods of 10 to 20 minutes after capsaicin injection, and in the glutamate test, at for 15 minutes following the glutamate injection. The protocols were approved by Animal Ethics Committee at the UFS (CEPA/UFS: 72/11). The data were analyzed by ANOVA (one-way) followed by Tukey's test (p<0.05).

CT significantly (p < 0.001) reduced, in both phases, the orofacial pain in the formalin test. In the first phase at doses of 25 (42.8 ± 6.7), 50 (42.5 ± 7.2) and 100 mg/kg (26.3 ± 2.0) when compared with control (93.5 ± 6.3), and in the second phase at 25 (47.3 ± 18.8), 50 (21.6 ± 10.2) and 100 (2.0 ± 1.3) when compared with control (168.7 ± 33.0). The orofacial pain induced by capsaicin was, also, significantly (p < 0.001) reversed at 25 (67.1 ± 14.4), 50 (64.3 ± 15.2) and 100 mg/kg (42.3 ± 17.4) comparing with control (193.7 ± 16.2). In the glutamate-induced nociception, the CT significantly (p < 0.01 or p < 0.001) reduced the orofacial pain at doses of 25 (48.3 ± 8.1), 50 (37.3 ± 7.4) and 100 mg/kg (24.1 ± 7.0) comparing with control (86.1 ± 8.7).

Our results suggest that CT reduces the orofacial nociceptive behavior in mice.

Acknowledgments: CAPES, CNPq, FAPITEC-SE.



## P 8. Chemical composition of the leaf essential oils from lavandin transformed with limonene synthase gene

Asada S<sup>1</sup>; Tsuru M<sup>1</sup>

<sup>1</sup>Faculty of Agriculture, Meijo University, Nagoya City, Japan

113561001@ccalumni.meijo-u.ac.jp

Keywords: lavandin, limonene synthase, transgenic plant, cyclic monoterpenes

Lavender is a well known aromatic plant, whose essential oil from florets comprises several monoterpenes, including linalool, linalyl acetate, limonene, and lavandulol [1]. The leaves of lavender also produce monoterpenes such as 1,8-cineol and camphor, whose fragrance differs from that of florets. These compounds are produced via an enzymatic reaction from the common precursor geranyl diphosphate [2]. We hypothesized that it should be possible to produce floret-specific monoterpenes in the leaves using gene transfer. In this study, we transformed the limonene synthase gene driven by a constitutive promoter into lavender plants and analyzed essential oil components in the leaves.

Lavandin, the interspecific hybrid between true lavender and spike lavender, was used as the plant material. Leaf-derived callus was inoculated with *Agrobacterium tumefaciens* harboring the binary plasmid, pRI101AN/LIMS, into which the limonene synthase gene of true lavender (*LIMS*) was driven by the 35S promoter within the T-DNA region and then regenerated using the method of Tsuru et al [3]. The essential oils of the leaves from regenerates were extracted with *n*-hexane and analyzed by GC and GC-MS.

Three regenerated transformants, expressing 2.4- to 8.8-times higher levels of *LIMS* mRNA than non-transgenic control, were obtained. Comparative analysis of the essential oil components identified ten compounds: eight terpenoids and two aromatics. All the transformants produced significantly more borneol than did the non-transgenic control. Furthermore, significantly higher levels of the terpenoids of 1,8-cineol and limonene,  $\alpha$ -pinene and camphor, were detected in the two transformants than the non-transgenic control. From these results, limonene synthase was suggested to act as both a limonene synthase and a cyclic monoterpene synthase. We are now evaluating their growth habit and essential oil components of florets.

1. M Tsuru et al. (2001) *Sci. Hortic.* 88: 309-317.
2. L Alexander et al. (2010) *Planta* 231: 835-845.
3. M Tsuru et al. (2009) *J. Japan. Soc. Hort. Sci.* 78: 236-241.

### P 9. Antinociceptive effect of *Xylopi* *laevigata* leaf essential oil in mice – evidence of involvement of periaqueductal gray area

Quintans JSS<sup>1</sup>; Antonioli AR<sup>1</sup>; Cardoso RSD<sup>1</sup>; Prado DS<sup>1</sup>; Costa EV<sup>1</sup>; Silva TB<sup>1</sup>; Prata APN<sup>1</sup>; Lima PSS<sup>2</sup>; Bonjardim LR<sup>1</sup>; Quintans-Júnior LJ<sup>1</sup>

<sup>1</sup>Federal University of Sergipe – Brazil

<sup>2</sup>Federal University of Sergipe

jullyanas@yahoo.com.br

Keywords: *Xylopi* *laevigata*, pain, periaqueductal gray

*Xylopi* *laevigata* (Annonaceae) is commonly known as ‘meiú’ or ‘pindaíba’, and it is a plant used in folk medicine of Brazilian Northeast to treat painful and inflammatory disorders. Thus, we investigated the possible antinociceptive effect of *Xylopi* *laevigata* leaf essential oil (EOX) and trying to describe the possible involvement of central nervous system (CNS) areas.

EOX was extracted by hydrodistillation of fresh leaves of the *X. laevigata*. The identification of the components was made through comparison of substance mass spectrum with the database of the GC-MS, literature and retention index. The antinociceptive effect of EOX was examined using the acetic acid (0.85%) writhing reflex and formalin (20 µl of 1%)-induced nociception tests. The motor coordination was also evaluated using Rota rod (7 rpm, 180 s). Animals were pretreated with EOX at the doses of 12.5, 25 or 50 mg/kg (i.p.) 1 h before injection or vehicle (saline + 2 drops of Tween 80 0.2%). Experimental protocols were approved by the animal care and use Committee (CEPA/UFS: 16/11) at the Federal University of Sergipe. The obtained data were evaluated by one-way analysis of variance (ANOVA) followed by Tukey’s test.

The phytochemical analysis of EOX showed the presence of g-murolene (17.78%), δ-cadinene (12.23%), bicyclogermacrene (7.77%), α-copaene (7.17%), germacrene D (6.54%) and (*E*)-caryophyllene (5.87%) as the main compounds. In mice, the EOX also significantly produced ( $p < 0.05$  or  $p < 0.001$ ) an antinociceptive effect by reduction in the early and late phases of paw licking and by a reduced writhing reflex (formalin and writhing tests, respectively) (Table 1). Such results were unlikely to be provoked by motor abnormality (Data no shown). The immunofluorescence essay demonstrated that acute treatment with EOX produces a significant ( $p < 0.01$  or  $p < 0.001$ ) increase in the number of Fos positive cells in the piriform cortex and periaqueductal gray CNS areas (Figure 1).

Altogether, the results provide evidence for the use of *X. laevigata* by traditional medicine practitioners in the management of pain and inflammatory disorders.

Acknowledgments: FAPITEC-SE, CNPq and CAPES, Brazil.

## P 10. Composition of the essential oil from the flowers of *Solandra maxima* (Sessé & Moc.) P.S. green

Radoias G<sup>1</sup>; Bosilcov A<sup>1</sup>

<sup>1</sup>Brüder Unterweger GmbH, 9911 Thal-Assling, Austria

[alin.bosilcov@unterweger-oils.com](mailto:alin.bosilcov@unterweger-oils.com)

Keywords: *Solandra maxima*; Solanaceae; essential oil composition

*Solandra* is a genus of flowering plants in the nightshade family (Solanaceae) having phytochemical, ethno botanical and ornamental importance. They have very large, attractively scented flowers and are native to the Caribbean, Mexico and South America. To the best of our knowledge, no data dealing with its volatile compounds have been yet reported. In this study, the dried flowers of *S. maxima* cultivated in Austria under greenhouse conditions were hydrodistilled yielding 0,28% (w/w) of essential oil; alternatively, the flowers were subjected to enzymatic hydrolysis using beta-glucosidase prior to hydrodistillation, yielding 0,34% (w/w) of essential oil. The two samples were subjected to a comparative investigation by GC-FID and GC/MS. More than 70 compounds were identified in both samples, representing more than 90% of the total. The oil is dominated by oxygenated compounds: alcohols, phenols and esters were the major classes, the concentration of terpene alcohols being generally higher in the sample obtained after enzymatic hydrolysis. The main constituents in the two samples were linalool (12,4% and 17,9%), methyl salicylate (9,1% and 11,4%), safranal (3,8% and 5,7%), alpha-terpineol (2,0% and 4,5%), geraniol (2,6% and 3,6%), carvacrol (17,8% and 11,5%), (E)-nerolidol (3,4% and 4,3%) and benzyl salicylate (11,1% and 7,1%). This for a tropical flower rather unusual composition corresponds to a complex odour profile that can be described as heavy-floral, spicy-phenolic and rounded up by tobacco notes.

1. LM Bernardello, AT Hunziker (1987) *A synoptical revision of Solandra (Solanaceae)*. Nord. J. Bot. 7(6): 639-652
2. LM Surhone, MT Tennoe, SF. Henssonow (ed.) (2010) *Solandra*. Betascript Publishing, Beau Bassin-Rose Hill, Mauritius
3. C Rättsch (2004) *Enzyklopädie der psychoaktiven Pflanzen*, 7<sup>th</sup> ed. AT Verlag, Aarau, Switzerland
4. RP Adams (2007) *Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry*, 4<sup>th</sup> ed. Allured Publ. Corp., Carol Stream, IL.

## P 11. Antimicrobial activity of essential oil of *Satureja montana* spp. *pisidica* (Lamiaceae)

Markovic S<sup>1</sup>; Sukilovic B<sup>1</sup>; Kundakovic T<sup>1</sup>; Milenkovic M<sup>2</sup>; Lakusic B<sup>3</sup>

<sup>1</sup>Department of Pharmacognosy, Faculty of Pharmacy, University of Belgrade, Serbia

<sup>2</sup>Department of Microbiology and Immunology, Faculty of Pharmacy, University of Belgrade, Serbia

<sup>3</sup>Department of Botany, Faculty of Pharmacy, University of Belgrade, Serbia

smark.kraljevo@yahoo.com

Keywords: *Satureja montana* spp. *pisidica*; essential oil; antimicrobial activity

*Satureja montana* spp. *pisidica* (Wettst.) Šilic (Lamiaceae) is an aromatic semi-shrub that grows only in FYR of Macedonia as an endemic species. All species of the genus *Satureja* contain more than 0.5% of essential oil with strong antimicrobial activity [1,2]. The aim of this work was to study the composition and antimicrobial activity of essential oils of *S. montana* spp. *pisidica* from two different locations.

Plant material was collected on the mountains Galičica and Korab in July 2011 and subjected to hydrodistillation on Clevenger apparatus. The composition of essential oils was determined using GC/MS and GC/FID, while antimicrobial activity was evaluated using broth microdilution method and expressed as minimal inhibitory concentration (MICs, µg/ml).

Thirty six components were identified in the essential oil of *S. montana* from Galičica, which represents about 95% of the oil. Forty two components were identified in the sample from Korab or about 94% of the oil. In both oils, carvacrol was the most abundant component (Galičica 37.6%; Korab 20.9%). Antimicrobial activity was tested against selected bacteria and fungi: *Staphylococcus aureus*, *S. epidermidis*, *Micrococcus luteus*, *Bacillus subtilis*, *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, and *Candida albicans*. Tested essential oils exhibited strong antibacterial activity, especially against G(+) bacteria, and lower antifungal activity. MICs values ranged from 12.5-50.0 µg/ml (bacteria), and 25.0-50.0 µg/ml (fungi). The essential oil of the sample from Galičica was demonstrated the best activity against *S. epidermidis* (12.5 µg/ml). The most resistant was *P. aeruginosa* (50.0 µg/ml). This effect can be attributed to the monoterpene components thymol, carvacrol and p-cymene.

Acknowledgments: Ministry of Education and Science of Republic of Serbia (P173021).

1. T Kundaković et al. (2011) Nat. Prod. Com. 6: 1353-1356.

2. J Antić Stanković (2008) J. Serb. Chem. Soc. 73: 703-711.

**P 12. The volatile oil profile of *Salvia officinalis* cultivated in Arad county, Romania**

Munteanu MF<sup>1</sup>; Socaci S<sup>2</sup>; Hanganu D<sup>3</sup>; Olah NK<sup>4</sup>

<sup>1</sup>Faculty of Pharmacy, "Vasile Goldis" West University of Arad

<sup>2</sup>Department of Food Science and Technology, University of Agricultural Sciences and Veterinary Medicine, Cluj – Napoca

<sup>3</sup>Department of Pharmacognosy, Faculty of Pharmacy, "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj – Napoca

<sup>4</sup>Department of Pharmaceutical Industry, Faculty of Pharmacy, "Vasile Goldis" West University of Arad

anaionescu@yahoo.com

Keywords: *Salvia officinalis*; GC-MS; cineole; bornyl acetate

The volatile oil is the main compound of the species of *Salvia officinalis* L. (Lamiaceae). This study presents the evaluation of chemical composition of the *Salvia* essential oil obtained from plants cultivated and harvested in Arad county, Romania.

The volatile oil of *Salvia officinalis* was obtained by hydrodistillation (57 ml/kg) by means of the special device described in the European Pharmacopoeia. The distillation took place in 4 hours at a speed of 2-3 mL/minute. 0.6 mL volatile oil was brought to 10 mL with hexane (Merck, Germany). A TLC and GC-MS determination followed on the thus obtained volatile oil. The qualitative analysis by thin layer chromatography was performed using the chromatographic conditions mentioned in the monographs for *Salvia officinalis* from the German Homeopathic Pharmacopoeia respectively the European Pharmacopoeia. Based on the R<sub>f</sub> values and the aspect of the bands, compared to those of the standards applied on the plate, we could identify in the volatile oil of common sage: thujon, cineole and bornyl acetate. In the chromatogram there are still other bands in different shades of violet and of different intensities which may correspond to some terpenic compounds.

We can notice that the volatile oil contains small quantities of thujon, a neurotoxic compound. For a more precise determination of the terpenic compounds, the GC-MS analysis was performed. The obtained results confirm the data of the TLC analysis and show the possibility of identifying 16 terpenic compounds having different concentrations from 0,09 to 13,40 %. In the volatile oil of common sage we could identify as main compounds: 1,8-cineole (13.40 %), camphor (8.08 %); alpha-pinene (5.81 %); camphene (3.70 %) and thujon (2.76 %).

The results shown that the studied common sage contains less thujon (2.76 %) and camphor (8.08%), than is found in reference data (20-60 % for thujon, 14-37% for camphor) respectively a higher concentration of 1,8-cineole (13.40 %), that is almost at the maximum found in reference data (6-16%) [1].

1. PDR for Herbal Medicine (2004) Third Ed., Thomson PDR, Montvale, p. 698-700.

**P 13. Morphoanatomical and chemical composition of leaf essential oil of two *Lippia alba* morphotypes**

Costa LCB<sup>1</sup>; Silva DC<sup>1</sup>; Oliveira RA<sup>1</sup>; Jezler CN<sup>1</sup>; Oliveira ARMF<sup>1</sup>; Batista RS<sup>1</sup>

<sup>1</sup>Universidade Estadual de Santa Cruz

larissacbc@hotmail.com.br

Keywords: morphology; leaf anatomy; Verbenaceae

*Lippia alba* is commonly known in south of Bahia, Brazil, as "cidreira" or "erva-cidreira" and used as a tea in the treatment of respiratory infections and stomach problems. In this same region is possible to distinguish two morphotypes of *L. alba* with a degree of polymorphism, used irrespective by the population for the same indications, although only one is chemically known. The objective of this work was to characterize and compare the morphology of these plants, the leaf secretory structures, and the essential oil produced. Studies on medicinal and aromatic plants indicate that it is common in some species the occurrence of accesses with morphological and phytochemicals variations (1).

The morphotypes were called "cidreira" and "melissa" due to their popular name originally associated. "Cidreira" is more high, erect and has larger leaves than "melissa". Although some authors (2) reported there are not important anatomic variations between chemotypes of this specie, we found differences in the number of layers of palisade parenchyma and the types of trichomes. They present identical anatomical pattern except for the number of layers of palisade parenchyma, which is biserial in "cidreira" and uniseriate on "melissa". The two morphotypes have glandular and no glandular trichomes on both leaf surfaces, and the last confirmed the accumulation of essential oils by histochemical test using Nadi reagent (3). The glandular trichomes were described and classified into six types distributed differently in the two morphotypes. There were differences in the content and chemical composition of essential oil, with citral as the major component of "cidreira" and citral, limonene and carvone in "melissa". This comparative study allowed us to distinguish two morphotypes of *Lippia alba* through morphological differences as plant size, leaf area, number of layers of palisade parenchyma and types of trichomes. Differences were also found in the content and chemical composition of essential oil.

1. J Hadian et al. (2010) Industrial Crops and Products. 32: 62-69.

2. MRA Santos et al. (2004) Revista Ciência Agronômica. 35: 377-383.

3. R David and JP Carde (1964) Comptes Rendus de l' Academie des Sciences. 258: 1338-1340.

**P 14. Allelopathic effect of *Cymbopogon winterianus* Jowitt essential oil on respiratory metabolism of *Euphorbia heterophylla***

Reis B<sup>1</sup>; Kern K<sup>1</sup>; Marques RM<sup>1</sup>; Ishii-Iwamoto EL<sup>1</sup>; Salgueiro L<sup>2</sup>; Cavaleiro C<sup>2</sup>; Ferrarese-Filho O<sup>1</sup>; Bonato CM<sup>1</sup>

<sup>1</sup>Universidade Estadual de Maringá

<sup>2</sup>Universidade de Coimbra

[cmbonato@hotmail.com](mailto:cmbonato@hotmail.com)

Keywords: Geraniol; citronellal; citronellol; bioherbicide

Allelopathic effect of *Cymbopogon winterianus* Jowitt essential oil on respiration of *Euphorbia heterophylla* (dairy) was evaluated. Seeds were treated with essential oil of *C. winterianus* (0 to 0.25% with 2% DMSO) and placed randomly on germination chamber (Type BOD) with 12h photoperiod at 25°C for 7 days. Effects of essential oil on respiration of embryos and mitochondria isolated pretreated with essential oil were measured. Direct effect of essential oil on mitochondrial respiration also was determined. The data were analyzed by ANOVA and the variables analyzed by regression at 5% probability. Total respiration and cyanide-sensitive respiration (SHAM) were affected by pretreatment of embryos with the essential oil of all species. Essential oil of *C. winterianus* significantly reduced mitochondrial oxidative phosphorylation measured by O<sub>2</sub> consumption during ADP phosphorylation (state III) and for ADP/O when using NADH or malate as respiratory substrate. Basal, state IV and respiratory control (RC) did not differ significantly from control. The results demonstrate that the essential oil dramatically affects both the total respiration and the sensitive to KCN one. It's probably that essential oil interferes on ATP-synthase complex activity and/or adenylate (e.g ADP) supply. Therefore less adenylates may reach mitochondrial matrix to be phosphorylated. Thus, essential oil of citronella has high potential to be used as bioherbicide.

Acknowledgments: The authors thank to Fundação Araucária, CAPES and Fundação para a Ciência e Tecnologia (FCT), POCI 2010/FEDER for financial support to this work.

1. KA Kern, EM Pergo, FL Kagami, LS Arraes, MA Sert, EL Ishii-Iwamoto (2009). The phytotoxic effect of exogenous ethanol on *Euphorbia heterophylla* L. *Plant Physiology and Biochemistry*, 47: 1095-1101.

**P 15. Effect of *Pinus elliottii* essential oil on initial growth of *Ipomoea grandifolia* L.**Medina AF<sup>1</sup>; Marques RM<sup>1</sup>; Ishii-Iwamoto EL<sup>1</sup>; Salgueiro L<sup>2</sup>; Cavaleiro C<sup>2</sup>; Bonato CM<sup>1</sup><sup>1</sup>Universidade Estadual de Maringá<sup>2</sup>Universidade de Coimbra

cmbonato@hotmail.com

Keywords:  $\alpha$ -pinene,  $\beta$ -pinene; geraniol; citronellol; allelopathy

The weed *Ipomoea grandifolia* cause large reductions in productivity in some crops such as soybeans. The control of this weed with herbicide may cause several problems to the environment. Thus, aim of this study was to evaluate the effect of essential oil of *Pinus elliottii* (bioherbicida) on germination and growth of weed *Ipomoea grandifolia* L. In petri dish containing 15 seeds of sorghum in a circular distributed were added different petri dishes were placed in a growth chamber (type BOD), temperature of 25°C (2 ±). The concentrations of pine essential oil of each petri dish varied from 0 to 1% (v/v). Control was consisted of DMSO (dimethyl sulfoxide – 2%). The germination variables were analyzed during period of 72h. Germination percentage ( $\%G = \frac{\sum ni.Ni}{\sum ni} \cdot 100$ ), germination average time ( $GAT = \frac{\sum ni.ti}{\sum ni}$ ), germination average speed (GAS) and germination speed index (GSI) were evaluated. In addition, biometric variables such as root system and shoot growth, fresh and dry weight of seedlings were evaluated.

The essential oil of pine affected the kinetic variables and the percentage of germination. Concentrations of 0.6 and 1% dramatically inhibited the germination percentage (%G). One hundred percent germination was inhibited in a concentration of 1%. GAT, GAS and GSI variables were highly affected by the increase in the concentration of essential oil. The effect on these variables was proportional to the concentration of essential oil tested. All biometric variables were also affected by the essential oil in a dose-dependent mode. Major compounds in pine essential oil are  $\alpha$ -pinene (14,7%) and  $\beta$ -pinene (8,4%), geraniol (8,3%) and citronellol (7,9%). These compounds seem to act on the plant energy metabolism so that could explain the results presented here<sup>1,2</sup>. Thus, it appears that this essential oil has great potential as bioherbicide. Our results suggest that the pine essential oil demonstrates high potential as bioherbicide to control the weed *Ipomoea grandifolia* L.

Acknowledgments: The authors thank to Fundação Araucária, CAPES and Fundação para a Ciência e Tecnologia (FCT), POCI 2010/FEDER for financial support to this work.

1 D Abraham, L Takahashi, AB Kelmer-Bracht, EL Ishii-Iwamoto. (2003). Allelopathy Journal, 11: 21-30.

2 D Abraham, AC Francischini, EM Pergo, AB Kelmer-Bracht, EL Ishii-Iwamoto (2003). Plant Physiology and Biochemistry: 41, 985-991.



**P 16. Essential oil of *Baccharis dracunculifolia*: antitumoral and antiviral activities**

Bufalo MC<sup>1</sup>; Sforcin JM<sup>1</sup>

<sup>1</sup>Biosciences Institute, UNESP, Botucatu, Brazil

michelle\_bufalo@yahoo.com.br

Keywords: essential oil; *Baccharis dracunculifolia*; HEp-2 cells; poliovirus

*Baccharis dracunculifolia* D. C., a native plant from Brazil commonly known as “alecrim-do-campo”, has been indicated as the most important botanical source of a Brazilian propolis [1]. Since propolis is known for its antiviral and antitumoral activities [2], the goal of this work was to evaluate the effect of essential oil of *B. dracunculifolia* against poliovirus and human laryngeal epidermoid carcinoma (HEp-2) cells in vitro.

The essential oil of the dried leaves (500 g) was extracted by hydro-distillation using a Clevenger type apparatus. HEp-2 cells were incubated with different concentrations (5, 10, 25, 50 and 100 µg/well) of essential oil for different time periods (24, 48 and 72h), and the percentage of viable cells were analysed by crystal violet test. The effects of 0.2% DMSO were also analysed (essential oil solvent). For the antiviral assay, HEp-2 cells were incubated with 100DICT50 of virus suspension and stimulated with non-cytotoxic concentrations of essential oil for 48h. The viral quantification of poliovirus was evaluated using real-time PCR.

Data showed that 5, 10 and 25 µg/well of essential oil of *Baccharis* extracts were not cytotoxic to HEp-2 cells ( $p > 0.05$ ). However, higher concentrations of *Baccharis* extracts (50 and 100 µg/well) had an efficient cytotoxic effect against these cells ( $p < 0.05$ ). 0.2% DMSO showed no effects in the number of viable cells, evidencing that the cytotoxic effects were exclusively due to essential oil components. For the antiviral activities, 25 µg/100 µl the essential oil showed the lower relative viral quantification in real-time PCR.

Essential oil of *B. dracunculifolia* showed an antitumoral activity in a dose- and time-dependent manner, and important antiviral action. Further assays in vivo are still needed in order to explore a better comprehension of antiviral and antitumoral properties of *Baccharis dracunculifolia*.

Acknowledgments: Financial support: CNPq

1. Park et al. (2004) J. Agricult. Food. Chem. 52: 1100-1103.
2. Salatino et al. (2005) Evid. Based. Complement. Alternat. Med. 2: 33-38.

### P 17. Essential oil content and composition of some oregano species cultured in Denizli district, oregano production center of Turkey

Baydar H<sup>1</sup>; Arabacı O<sup>2</sup>

<sup>1</sup>University of Süleyman Demirel-Turkey

<sup>2</sup>University of Adnan Menderes-Turkey

hasanbaydar@sdu.edu.tr

Keywords: Turkey, Denizli district, Oregano, cultivation, essential oil

Turkey supplies more than 50% of the world's exported oregano (*Origanum* types). Denizli province is the primary Turkish oregano (*Origanum onites* L.) cultivation center in Turkey with a ratio of over 80% in both production and exporting. According to the registrations of Denizli Provincial Directorate of Agriculture, oregano cultivation area and the amount of production in 2009 were 7839 ha and 10660 tones, respectively. Recently *Origanum vulgare* subsp. *hirtum* (Link) leestwaart (Greek or Istanbul oregano) has also started to cultivate in limited field areas. *O. sipyleum* L. (Anatolian oregano) is an endemic species of the Western Anatolia, Turkey. In this research, essential oil content and composition of these species grown in Denizli district at about 1200 m altitude during flowering period on July, 2011 were determined by Clevenger type hydro-distillation apparatus and by GC/MS system, respectively. Plant samples from the locations were analyzed in terms of leaf, flower and stem ratios in the dried herb. The soil samples were analyzed in terms of their physical and chemical properties. According to the results of chemical soil analysis, oregano soils in the region were generally medium-textured and non-salinity, but high alkaline and excessively calcareous, and with low levels of organic matter. The main compound in the essential oil of the all oregano species examined in this study was carvacrol. Carvacrol percentages in the leaf and flower essential oils were 84.03 and 87.83% in *O. onites*, 78.23 and 81.96% in *O. vulgare* subsp. *hirtum*, 60.74 and 85.71% in *O. sipyleum*, respectively. Essential oils distilled from the dried flowering tops contained more carvacrol than that of the essential oils distilled from the dried leaves. Essential oil ratio of oregano should be at least 2% according to ASTA standards. When these standards are taken into consideration, Turkish and Greek oreganos grown in non-irrigated, loamy and alkali soils of Denizli district were found to be higher than the standards.

**P 18. Yield and chemical composition of the essential oil of Moroccan chamomile [*Cladanthus mixtus* (L.) Chevall.] growing wild in different areas of Morocco**

El Amrani A<sup>1</sup>; Anass E<sup>1</sup>; Jamal Eddine J<sup>1</sup>; Barroso JG<sup>2</sup>; Pedro LG<sup>2</sup>; Figueiredo AC<sup>2</sup>

<sup>1</sup>Laboratoire Synthèse, Extraction et Etude Physico-Chimique des Molécules Organiques, Faculté des Sciences Aïn -Chock, B.P 5366 Mâarif – Casablanca – Morocco

<sup>2</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portugal

aelamrani77@yahoo.fr

Keywords: *Cladanthus mixtus*; *Ormenis mixta*; *Chamaemelum mixtum*; *Anthemis mixta*; Asteraceae; Compositae; Moroccan chamomile

Recent floristic literature adopted new names for genera of Anthemideae (Compositae, Asteraceae). Based on phylogenetic analysis of *Chamaemelum* Mill. [1,2], the genus *Ormenis* is now named *Cladanthus* and therefore *Cladanthus mixtus* (L.) Chevall. is the newly adopted scientific name for the Moroccan chamomile [2].

Moroccan chamomile essential oil production ranks 9th in the Top 20 of Moroccan produced essential oils [3]. Given its commercial worth, the chemical variability of the essential oils isolated from Moroccan chamomile was evaluated.

Flowering aerial parts from *C. mixtus* populations were collected at nine Moroccan regions, Benguerir (Be), Bouznika (Bo), Chefchaouane (C), Kenitra (K), Meknes (M), Oujda (O), Settât (S), Sidi Alal Ibrahroui (SAI) and Tamesna (T). The essential oils were isolated by hydrodistillation and analyzed by GC and GC-MS as in [4]. The essential oils yields ranged between 0.1-0.8% (v/d.w.). Only five out of the nine essential oil samples analyzed showed a good correlation after agglomerative cluster analysis based on the essential oils chemical composition. These samples (Be, K, S, M and T) were characterized by the dominance of camphor (14-27%),  $\beta$ -myrcene (3-17%) and santolina triene (3-15%). All these, and C essential oils showed blue colour, due to the presence of chamazulene (<1%), whereas O, Bo and SAI essential oils were yellow.  $\beta$ -Myrcene (3-17%), *trans*- $\beta$ -farnesene (18%) and 2-tridecanone (16%) dominated C essential oil, whereas *trans*- $\beta$ -farnesene (43%) was the main component in O essential oil. 2-Methyl-2,*trans*-butenyl methacrylate (34%) dominated Bo sample, and santolina alcohol and 1,8-cineole (17% and 12%, respectively) were the main components in SAI essential oil.

The presence of some characteristic volatile components suggests that Moroccan chamomile essential oils variability may result from the fact that under the same common chamomile name different species may be collected, and reinforces the importance of avoiding the wild population's harvest. Being the chemical composition so diverse, this will reflect negatively in the essential oil quality and biological activity.

Acknowledgments: Partially funded by the Transnational Cooperation research contract FCT/CNRST - 6818.

1. C Oberprieler (2002) *Bot. J. Linn. Soc.* 138: 255–273.
2. W Greuter et al. (2003) *Willdenowia* 33: 37-43.
3. B Lawrence (2009) *Perfumer & Flavorist* 34: 38-44.
4. SM Albano et al. (2012) *Rec. Nat. Prod.* 6: 35-48.

**P 19. New esters of long chain alcohols and isobutanoic and isovaleric acids from the essential oil of *Scandix pecten-veneris* L. (Apiaceae)**

Radulović NS<sup>1</sup>; Mladenović MZ<sup>1</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science and Mathematics, University of Niš

markohem87@gmail.com

Keywords: *Scandix pecten-veneris* ssp. *brachycarpa*; volatiles; synthesis; C13-C18, C21 and C23 alcohols; isobutanoate; isovalerate

*Scandix pecten-veneris* L. (Apiaceae) is a wild-growing plant species native to Eurasia. This species is commonly divided into three different subspecies - *brachycarpa* (Guss.) Thell, *pecten-veneris* L. and *macrorhyncha* (C. A. Meyer) Rouy & Camus.[1]. The young stem tops of the plant have been used to stop bleeding, amenorrhea and for bladder affections [2]. Since the volatiles of this plant species have been the subject of only two previous studies, that identified 12 and 10 constituents in total [2,3], we decided to perform a more detailed analysis of the aerial parts and roots essential oil of *S. pecten-veneris* ssp. *brachycarpa* from Southeastern Serbia (in total 7 samples). Among one hundred-nine constituents identified by GC (FID) and GC/MS, tridecane (8.5-52.8%), pentadecane (1.3-23.5%) and  $\beta$ -caryophyllene (0.1-18.5%) were found to be the most abundant ones. The analysis also pointed to the presence of 14 additional minor compounds (up to 0.9%), that were, according to their mass spectral fragmentation, tentatively identified as the isomeric tricosanyl-, heneicosanyl-, tri-, tetra-, penta-, hexa-, hepta- and octadecyl pentanoates and butanoates. In order to corroborate the tentative identification of these esters, and obtain sufficient quantities for biological activity assays, a synthesis was undertaken of 17 esters of isobutanoic, isovaleric and 2-methylbutanoic acids with appropriate alcohols. All synthesized esters (of which 9 entirely new and 13 found for the first time as plant secondary metabolites) were spectrally characterized (IR, MS, NMR) and separately GC co-injected with the essential oils. Except for tri-, penta- and heptadecyl 2-methylbutanoates, all other synthesized compounds were detected in the analyzed oils of *S. pecten-veneris*. The content of the new esters was determined from the appropriate GC-FID calibration curves of excellent linearity ( $R^2=0.9999$ ) and varied within the range 0.01-9.33 mg / 100 g of dry plant material. The testing of the oils and of the herein synthesized compounds for antimicrobial activity is under way.

Acknowledgments: This work was funded by the Ministry of Education and Science of Serbia (Project 172061).

1. TG Tutin, VH Heywood, NA Burges, DM Moore, DH Valentine, SM Walters, DA Webb (1968) Flora Europaea, University press, Cambridge.
2. M Ashraf, R Ahmad, MK Bhatti (1979) Pakistan J. Sci. Ind. Res. 6: 320-321.
3. KH Kubeczka (1982) Aromat. Plants, 7: 165-73.

**P 20. Volatile secondary metabolites of *Senecio vernalis* Waldst. & Kit.**Radulović NS<sup>1</sup>; Blagojević PD<sup>1</sup>; Mladenović MZ<sup>1</sup><sup>1</sup>Department of Chemistry, Faculty of Science and Mathematics, University of Niš

markohem87@gmail.com

Keywords: *Senecio vernalis* Waldst. & Kit.; Asteraceae; essential oil; roots; above-ground parts; chemotypification

*Senecio vernalis* Waldst. & Kit., Asteraceae, (local Serbian name: žutenika) is a plant species widely distributed in Asia and SE Europe [1]. However, limited data on the chemical composition of the corresponding essential oil exist [1,2]. For this reason, detailed chemical analyses of the volatile secondary metabolites of several populations of *S. vernalis* were conducted in order to provide an insight into the chemotypification of this species in Serbia. Analyses (by GC and GC-MS) of 8 essential oils hydrodistilled from dry above-ground parts and rhizomes of *S. vernalis* (4 different populations) enabled the identification of 60 different constituents, representing 95.5-99.4% of the total oils. The most abundant constituents of the oils obtained from the above-ground parts (all populations) were  $\alpha$ -pinene (26.7-38.5%) and  $\beta$ -pinene (23.0-24.8%), whereas the most dominant constituents of the rhizome oils were either  $\alpha$ -humulene (40.9-57.0%, 3 samples) or nonacosane (33.3%, 1 sample). Hence, the biosynthesis and/or accumulation of volatile metabolites in *S. vernalis* is strongly plant organ dependent. This was additionally corroborated by agglomerative hierarchical cluster analysis (AHC) and principal component analysis (PCA). Both AHC and PCA analyses of 30 different essential oils (18 different *Senecio* taxa; data from the literature and the present study) were performed using four different types of variables.

Acknowledgments. This work was funded by the Ministry of Education and Science of Serbia (Project 172061). PB was supported by UNESCO-L'OREAL National Fellowships Programme for Young Women in Life Sciences.

1. Nori-Shargh et al. (2008) Flavour. Fragr. J. 23: 357-359.
2. Usta et al. (2009) Asian J. Chem. 21: 6369-6374.

**P 21. Long chain 3-methyl-2-alkanones from the essential oil of *Inula helenium* L. (Asteraceae)**

Radulović N<sup>1</sup>; Denić M<sup>1</sup>; Stojanović-Radić Z<sup>2</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science and Mathematics, University of Niš

<sup>2</sup>Department of Biology and Ecology, Faculty of Science and Mathematics, University of Niš

denijum@yahoo.com

Keywords: *Inula helenium* L.; Asteraceae; root essential oil; 3-methyl-2-alkanones; antistaphylococcal activity; 3-methyl-2-tetradecanone

*Inula helenium* L. (Compositae) is a perennial herb originally native to Southeastern Europe. In Serbia, it is a widespread plant, used in folk medicine mostly for the treatment of respiratory conditions, disorders of digestion, urinary infections, and for skin disorders. Very recently, we reported that *I. helenium* root essential oil possesses a very potent antistaphylococcal activity, with obvious membrane-damaging effects. Bioassay guided fractionation of the oil pointed to alantolactone, isoalantolactone and diplophyllin as the main carriers of the observed activity [1].

Herein, we report the identification of additional nine constituents of the oil from a fraction with a low MIC value (0.8 mg/ml). Mass spectral fragmentation pattern (e.g. base peak at  $m/z$  72) of these compounds suggested their possible identity - long chain 3-methyl-2-alkanones. The difference of 14 amu (one CH<sub>2</sub> group) between the molecular ions of two consecutive compounds in the gas chromatogram, separated by ca. 100 RI units, pointed to the existence of a C<sub>10</sub>-C<sub>18</sub> homologous series. These 3-methyl-2-alkanones were synthesized in three reaction steps with the overall yields of 45-50% starting from an appropriate halogen alkane. Gas co-chromatography of the obtained compounds with the fraction of *I. helenium* root oil unequivocally corroborated the original assumption. The construction of GC-FID calibration curves ( $R^2=0.9997-0.9999$ ) enabled us to determine the content of the ketones in the plant material: 0.05-25.1 mg/100 g of dry roots. The synthetic 3-methyl-2-alkanones were also screened for in vitro antimicrobial activity. These long chain 3-methyl-2-alkanones have a rather restricted occurrence in samples of natural origin: 3-methyl-2-decanone was found in the territorial marking fluid of the male Bengal tiger, *Panthera tigris* [2], while 3-methyl-2-tridecanone, 3-methyl-2-tetradecanone and 3-methyl-2-pentadecanone were detected in Dufour glands secretions of the species of desert-dwelling ants of the *Cataglyphis bicolor* group [3]. Hence, this is the very first report of long chain 3-methyl-2-alkanones as the plant secondary metabolites.

Acknowledgements: The authors acknowledge the Ministry of Education and Science of Serbia for the financial support (Project No 172061).

1. Z Stojanović-Radić et al. (2012) Eur. J. Clin. Microbiol. Infect. Dis. 31:1015–1025.
2. BV Burger et al. (2008) J. Chem. Ecol. 34: 659–671.
3. O Gökçen et al. (2002) J. Chem. Ecol. 28: 71–87.

**P 22. Chemical composition of tuber essential oil from *Helianthus tuberosus* L. (Asteraceae)**

Radulović NS<sup>1</sup>; Đorđević MR<sup>1</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science and Mathematics, University of Niš, Serbia

*miljanadjordjevic11@gmail.com*

Keywords: *Helianthus tuberosus*; Jerusalem artichoke; tubers; essential oil; helianthol A; dihydroeuparin

The genus *Helianthus* L. (Asteraceae) comprises about 50 annual and perennial sunflower species. *Helianthus tuberosus* L. (Jerusalem artichoke) is cultivated in Europe and other parts of the world as a crop and ornamental plant. The volatile oils of the aerial parts of *H. tuberosus* were analyzed in 1982, but no study has been carried out to date to determine the constituents of the tuber essential oil [1]. Herein, we describe the first analysis by GC and GC-MS of the hydrodistilled essential oil of the Jerusalem artichoke tubers. The fresh plant material, collected in Serbia, yielded only a small amount of the oil (0.0014%, w/w). One hundred-one constituents were identified in total, representing 89% of the oil. The main constituents were  $\beta$ -bisabolene (18.8%), undecanal (10.4%),  $\alpha$ -pinene (6.2%), kauran-16-ol (5.7%), 2-pentylfuran (4.7%) and (2*E*)-tetradecenal (4.0%). Several rare compounds characteristic for *Helianthus* sp. were also detected: helianthol A (0.9%), desmethoxy enecalinal (0.2%), desmethyl isoencecalinal (trace amount), enecalinal (trace amount), dihydroeuparin (1.9%), euparin (0.2%) and eupatoriochromene (0.3%). The tuber essential oil and the aerial parts one share a common major component -  $\beta$ -bisabolene.

Acknowledgments: This work was funded by the Ministry of Education and Science of Serbia (Project 172061).

1. A MacLeod et al. (1982) *Phytochemistry*, 21: 1647-1651.

**P 23. Antimicrobial volatile oil of *Lycopus europaeus* L. (Lamiaceae)**Radulović N<sup>1</sup>; Denić M<sup>1</sup>; Stojanović-Radić Z<sup>2</sup><sup>1</sup>Department of Chemistry, Faculty of Science and Mathematics, University of Niš<sup>2</sup>Department of Biology and Ecology, Faculty of Science and Mathematics, University of Niš

denijum@yahoo.com

Keywords: *Lycopus europaeus* L.; gypsywort; Lamiaceae; essential oil; antimicrobial activity; hotrienol

*Lycopus europaeus* L. (Lamiaceae), commonly known as gypsywort, is a perennial plant native to Eurasia. Gypsywort's most important medicinal properties come from the stem and leaves and these were traditionally used as astringents, sedatives, tonics and narcotics, as well as in the treatment of hemorrhages, nervous heart palpitations, goiter and Graves' disease [1, 2]. On only one previous occasion the essential oil of this taxon was investigated [3]. However, this work was undertaken some 40 years ago and is lacking the cutting edge of modern analytical techniques. This provoked us to carry out a detailed study of *L. europaeus* essential oils from the blooming (sample A) and fruit forming stages (sample B), and to screen these for *in vitro* antimicrobial activity in order to determine if some of these constituents may contribute to the plant's renowned ethnomedicinal uses.

GC and GC/MS analysis of the oils allowed the identification of 199 constituents. The oils consisted mainly of sesquiterpenoids, the main one being  $\beta$ -caryophyllene (13.9% and 25.7%) in both cases. Other important contributors were shyobunol (9.7% of sample A) and caryophyllene oxide (2.4% and 12.5%, in samples A and B, respectively). In addition to this rather unequal distribution of the main sesquiterpenoids, another significant feature that distinguished the two samples was the inverted ratio of the mono- (15.6% against 1.6%) and diterpenoids (24.6% against 8.8%). The monoterpene fraction was characterized by the high content of oxygenated acyclic derivatives among which the most predominant one was the rare monoterpene (*E*)-hotrienol. Screening of *L. europaeus* essential oil for its *in vitro* antimicrobial activity showed that this oil possesses a selective action against two Gram negative bacterial strains, *Escherichia coli* and *Klebsiella pneumoniae*.

Acknowledgements: The authors acknowledge the Ministry of Education and Science of Serbia for the financial support (Project No 172061).

1. JK Crellin, J Philpott (1997) Herbal Medicine Past and Present: A Reference Guide to Medicinal Plants. 2nd Duke University Press, Durham.
2. G Pinn (2005) In: Z Yaniv, U Bachrach (Eds.) Handbook of Medicinal Plants. The Haworth Press, Inc., Binghamton, 383-398.
3. ShN Sharipov et al. (1969) Khim. Prir. Soedin+ 5:316.



**P 24. Determination of the absolute stereochemistry of limonene and  $\alpha$ -santalol by both vibrational circular dichroism and raman optical activity spectroscopies**

Sakamoto A<sup>1</sup>; Ohya N<sup>1</sup>; Hasegawa T<sup>1</sup>; Izumi H<sup>1</sup>; Tokita N<sup>2</sup>; Hamada Y<sup>2</sup>

<sup>1</sup>Department of Chemistry, Graduate School of Science and Engineering, Saitama University

<sup>2</sup>The Open University of Japan

toshihas@mail.saitama-u.ac.jp

Keywords: absolute configuration; Vibrational circular dichroism (VCD); Raman optical activity (ROA); limonene;  $\alpha$ -santalol

Chirality is an important factor affecting the odor characteristics of fragrance compounds. Thus, it is fundamentally important to determine the absolute configuration of chiral odor compounds. Although X-ray crystallographic analysis is a useful and powerful method for determining the stereochemistry of crystalline organic compounds, it is necessary to obtain the good-quality crystals of the target molecule. Many types of terpenoids, especially monoterpenes and sesquiterpenes, with chiral centers are known. These compounds are important in fragrance, medicinal, and food chemistry. However, most of these compounds are liquids, and so evaluating their absolute configuration is difficult. Vibrational circular dichroism (VCD) and Raman optical activity (ROA) spectroscopies should be useful in solving this problem. Recently, we reported the preliminary results by using these two methods [1]. We measured infrared, Raman, VCD, and ROA spectra of (*R*)-(+)-limonene, (*S*)-(-)-limonene, and (*E*)-(+)- $\alpha$ -santalol. The optimized structures and infrared, Raman, VCD, and ROA spectra were calculated for these compounds by DFT method at B3LYP/6-31G\* and B3LYP/6-311+G\*\* levels. We compared the observed infrared, Raman, VCD, and ROA spectra of the compounds with those calculated by DFT method. We succeeded in the determination of the absolute configurations of (*R*)-(+)-limonene, (*S*)-(-)-limonene, and (*E*)-(+)- $\alpha$ -santalol.

1. A Sakamoto, N Ohya, T Hasegawa et al. (2012) Natural Product Communications, 7, 419-421.

**P 25. Antimicrobial activity of essential oil isolated from young twigs of Macedonian pine (*Pinus peuce* Griseb., Pinaceae)**

Karapandzova M<sup>1</sup>; Stefkov G<sup>1</sup>; Trajkovska-Dokik E<sup>2</sup>; Kadifkova-Panovska T<sup>1</sup>; Kaftandzieva A<sup>2</sup>; Kulevanova S<sup>1</sup>

<sup>1</sup>Department for Pharmacognosy, Faculty of Pharmacy, Skopje, The Former Yugoslav Republic of Macedonia

<sup>2</sup>Department for Microbiology, Faculty of Medicine, Skopje, The Former Yugoslav Republic of Macedonia

marijakarapandzova@yahoo.com; marija\_karapandzova@ff.ukim.edu.mk

Keywords: *Pinus peuce*; essential oil; antimicrobial activity.

Essential oils isolated from pine species have shown variable antimicrobial activity against different microorganisms [1, 2, 3]. Up to present, there is a lack of data that are related to the antimicrobial activity of *Pinus peuce*, thus the aim of this study was to assess antimicrobial activity of the essential oil isolated from young twigs of *Pinus peuce* from Macedonian flora. Essential oil was isolated by hydrodistillation in Clevenger type apparatus. Analysis of the essential oil composition by GC/FID/MS method revealed  $\alpha$ - and  $\beta$ -pinene as dominant components, beside limonene+ $\beta$ -phellandrene, bornyl acetate, *trans*-caryophyllene and germacrene D. Antimicrobial screening of the essential oil was made by hole-plate diffusion and broth dilution method against 13 bacterial isolates representing both Gram positive and Gram negative bacteria and one strain of *Candida albicans*. The most sensitive bacteria against tested *Pinus peuce* essential oil were *Streptococcus agalactiae*, *Streptococcus pyogenes*, *Enterococcus*, *Candida albicans*, encompassing *Staphylococcus pneumoniae*, *Haemophilus influenzae*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Acinetobacter* spp., *Escherichia coli* and *Salmonella enteritidis*. Minimal inhibitory concentrations (MICs) of the oil ranged from 7.5 - 125  $\mu$ l/ml. As this essential oil has shown promising activity towards the above mentioned microorganisms, it could be considered as antimicrobial agent for inhalation for the infections of upper respiratory tract, for treatment of urinary and gastrointestinal infection as well as different skin conditions.

1. O Oluwadayo Sonibare, K Olakunle (2008) African Journal of Biotechnology 7(14): 2462-2464.
2. K Youg-Suk, Sh Dong-Hwa (2005) Food Microbiology 22: 37-45.
3. G Sacchetti, S Maietti et al (2005) Food Chemistry 91: 621-632.

## P 26. Essential oils composition of *Hyptis* species with occurrence in the Brazilian Amazon

Monteiro OS<sup>1</sup>, Carvalho NCC<sup>1</sup>, Carreira LMM<sup>2</sup>, Andrade EHA<sup>3</sup>, Maia, JGS<sup>3</sup>, Lucas FCA<sup>4</sup>

<sup>1</sup>Programa de Pós-Graduação em Química, Universidade Federal do Maranhão, 65080-040 São Luís, MA, Brazil

<sup>2</sup>Coordenação de Botânica, Museu Paraense Emílio Goeldi, 66040-170 Belém, PA, Brazil

<sup>3</sup>Programa de Pós-Graduação em Química, Universidade Federal do Pará, 66075-900, Belém, PA, Brazil

<sup>4</sup>Departamento de Ciências Naturais, Universidade Estadual do Pará, 66113-200 Belém, PA, Brazil

eloisandrade@ufpa.br

Keywords: *Hyptis crenata*; *H. recurvata*; *H. parkeri*; *H. dilatata*; *H. suaveolens*; *H. alutacea*; essential oil composition

The genus *Hyptis* (Lamiaceae) is composed of 400 species occurring in tropical America. The species *H. suaveolens*, *H. mutabilis*, *H. goyazensis* and *H. crenata* were previously studied by us [1-4]. The essential oils of leaves and fine stems of *Hyptis crenata* Pohl ex Benth. (1), *H. recurvata* Poit. (2), *H. parkeri* Benth. (3), *H. dilatata* Benth. (4), *H. suaveolens* (L.) Poit (5). and *H. alutacea* Pohl ex Benth. (6) were analyzed by GC and GC-MS. The plants were collected in Eastern Amazon, in the municipality of São Geraldo do Araguaia (1,2), Pará state, and in the municipalities of São Felix de Balsas (3) and Mirador (4-6), Maranhão state. The main constituents identified in the oil of *H. crenata* were 1,8-cineole (43.8%),  $\alpha$ -pinene (20.7%) and  $\beta$ -pinene (8.0%); in the oil of *H. recurvata* were  $\beta$ -elemeno (15.5%), (E)-caryophyllene (14.0%),  $\alpha$ -selinene (10.2%),  $\beta$ -selinene (5.6%),  $\gamma$ -elemene (4.9%), germacrene D (4.8%) and caryophyllene oxide (4.6%); in the oil of *H. parkeri* were spathulenol (27.9%),  $\delta$ -elemene (7.5%), bicyclogermacrene (7.2%) and caryophyllene oxide (6.1%); in the oil of *H. dilatata* were camphor (32.6%), 1,8-cineole (19.9%), p-cymene (5.2%),  $\gamma$ -terpinene (5.4%) and  $\alpha$ -pinene (4.3%); in the oil of *H. suaveolens* were (E)-caryophyllene (18.1%), bicyclogermacrene (11.2%),  $\delta$ -elemene (12.3%), 1,8-cineole (9.1%) and spathulenol (8.1%); and in the oil of *H. alutacea* were  $\gamma$ -cadinene (19.1%),  $\delta$ -cadinene (10.5%),  $\alpha$ -muurolene (8.0%) and (E)-caryophyllene (7.2%).

Acknowledgments: This work was supported by BIONORTE/CNPq and FAPESPA/PA.

1. Gottlieb et al. (1981) *Acta Amazonica* 11: 143-148.
2. Luz et al. (1984) *J. Nat. Prod.* 47: 745-747.
3. Luz et al. (1989) *Acta amazonica* 19: 365-370.
4. Rebelo et al. (2009) *Braz. J. Pharmacogn.* 19: 23-235

## P 27. Essential oils composition of some Myrtaceae species from Eastern Amazon, Brazil

Andrade EHA<sup>1</sup>, Monteiro OS<sup>2</sup>, Carvalho NCC<sup>2</sup>, Ribeiro AF<sup>3</sup>, Carreira LMM<sup>4</sup>, Maia JGS<sup>1</sup>, Luca FCA<sup>5</sup>

<sup>1</sup>Programa de Pós-Graduação em Química, Universidade Federal do Pará, 66075-900, Belém, PA, Brazil

<sup>2</sup>Programa de Pós-Graduação em Química, Universidade Federal do Maranhão, 65080-040 São Luís, MA, Brazil

<sup>3</sup>Faculdade de Ciências Exatas e Naturais, Universidade Federal do Pará, 68501-970, Marabá, PA, Brazil

<sup>4</sup>Coordenação de Botânica, Museu Paraense Emílio Goeldi, 66040-170 Belém, PA, Brazil

<sup>5</sup>Departamento de Ciências Naturais, Universidade Estadual do Pará, 66113-200 Belém, PA, Brazil.

eloisandrade@ufpa.br

Keywords: Myrtaceae species; essential oil variation; Eastern Amazon

The Myrtaceae is comprised of about 80 genera and 3,000 species distributed in warm climates, with centers of occurrence in Australia and America. As part of the survey of Amazon aromatic flora, this work presents the chemical composition of essential oils of the following species of Myrtaceae: *Psidium myrsinites* Mart. ex DC. (1), *Myrcia fallax* (Rich.) DC. (2), *M. atramentifera* Barb. Rodr. (3), *M. bracteata* (Rich.) DC. (4), *Calycolpus goetheanus* (DC.) O. Berg (5), *Eugenia puniceifolia* (Kunth) DC. (6), *E. patrisii* Vahl (7) and *E. piauiensis* O. Berg (8). These species were collected in eastern Amazon, comprising the municipality of São Geraldo do Araguaia (1-4), Pará state, and the municipalities of São Domingos do Azeitão (5) and Mirador (6-8), Maranhão state. The essential oils of these species were hydrodistilled and analyzed by GC and GC-MS. The main constituents (over 4%) of *P. myrsinites* oil were (*E*)-nerolidol (18.6%), 1,8-cineole (14.6%),  $\alpha$ -pinene (11.0%), (*E*)-caryophyllene (10.4%), *p*-cymene (4.5%) and carvacrol (4.0%); in the *M. fallax* oil were  $\alpha$ -cadinol (11.1%), 1-*epi*-cubenol (7.7%), *epi*- $\alpha$ -muurolol (6.9%), guaiol (6.5%), viridiflorol (5.9%),  $\delta$ -elemene (4.9%) and selin-11-en-4 $\alpha$ -ol (4.8%); in the *M. atramentifera* oil were  $\alpha$ -cadinol (12.9%),  $\delta$ -cadinene (10.6%), *epi*- $\alpha$ -muurolol (6.5%),  $\alpha$ -copaene (6.3%) and *epi*- $\alpha$ -cadinol (4.3%); in the *M. bracteata* oil were caryophyllene oxide (10.1%),  $\alpha$ -cadinol (10.0%),  $\delta$ -elemene (6.8%), (*E*)-caryophyllene (6.6%), *epi*- $\alpha$ -muurolol (5.5%),  $\alpha$ -calacorene (5.0%) and germacrene D (4.5%); in the *C. goetheanus* oil were (*E*)-caryophyllene (31.0%), caryophyllene oxide (14.7%) and  $\alpha$ -humulene (9.7%); in the *E. puniceifolia* oil were (*E*)-caryophyllene (18.8%), caryophyllene oxide (12.3%), 1-*epi*-cubenol (8.3%),  $\alpha$ -humulene (5.3%), *trans*-calamenene (4.7%),  $\alpha$ -cadinol (4.4%) and  $\alpha$ -copaene (4.2%); in the *E. patrisii* oil were germacrene D (23.5%), (*E*)-caryophyllene (10.9%),  $\delta$ -elemene (7.9%),  $\beta$ -copaene (5.7%), bicyclogermacrene (5.7%) and  $\delta$ -cadinene (5.6%); in the *E. piauiensis* oil were (*E*)-caryophyllene (19.7%),  $\gamma$ -elemene (14.7%) and  $\delta$ -elemene (4.5%).

Acknowledgments: This work was supported by BIONORTE/CNPq and FAPESPA/PA.

## P 28. Nematotoxic effect of essential oils and their fractions against the pinewood nematode, *Bursaphelenchus xylophilus*

Faria JMS<sup>1</sup>; Barbosa P<sup>2</sup>; Mota M<sup>2</sup>; Figueiredo AC<sup>1</sup>

<sup>1</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portugal

<sup>2</sup>NemaLab-ICAAM & Departamento de Biologia, Universidade de Évora, 7002-554 Évora, Portugal

jmfaria@fc.ul.pt

Keywords: *Bursaphelenchus xylophilus*; *Pinus pinaster*; essential oils; monoterpenes; nematocidals; 2-undecanone

The pinewood nematode (PWN) *Bursaphelenchus xylophilus* is a highly pathogenic plant parasite that greatly affects pine forests. In Portugal, the most affected species is *Pinus pinaster* Aiton. Despite great efforts, since its first detection in 1999, the PWN has spread through the country, including Madeira Island, having been recently detected in Spain [1,2]. Containing this pest is of the utmost importance for European pine forest safeguard.

Since most synthetic chemicals used to control phytoparasites are toxic to humans and animals, and can accumulate in the soil and in food plants [3], in the present work, the nematotoxic potential of over 80 essential oils (EOs), isolated from the Portuguese flora, were assessed against the PWN. EOs were isolated by hydrodistillation and analysed by GC and GC-MS [3]. EOs hydrocarbon and oxygen-containing fractions were obtained as in [4]. Direct-contact assays, adapted from [3], were performed by adding EOs/methanol stock-solutions to 50-100 mixed-stage PWN suspensions. After 24h in darkness, dead and live nematodes were counted under an inverted microscope. Assays were repeated at least 10 times in two series.

Mortalities  $\geq 96\%$  were obtained with  $2\mu\text{L}/\text{mL}$  of the EOs isolated from *Cymbopogon citratus*, *Eucalyptus citriodora*, *Mentha arvensis*, *Origanum virens*, *Origanum vulgare*, *Ruta graveolens*, *Satureja montana*, *Syzygium aromaticum*, *Thymbra capitata*, *Thymus caespititius* (carvacrol and/or thymol-rich), *Thymus vulgaris* and *Thymus zygis*. These EOs were further tested at 1, 0.5 and  $0.25\mu\text{L}/\text{mL}$ . Minimum lethal concentrations (LC100)  $< 0.4\mu\text{L}/\text{mL}$ , were obtained for the 2-undecanone-rich *R. graveolens* EO and the carvacrol and  $\gamma$ -terpinene-rich *S. montana* and *T. capitata* EOs. Assays with EO fractions revealed that the monoterpene-rich nematotoxic EOs control PWN through their combined hydrocarbon and oxygen-containing fractions through additive and/or synergic relations.

As complex mixtures of active components, EOs may prove to be effective nematotoxic agents.

Acknowledgments: Jorge Faria is grateful to the Fundação para a Ciência e a Tecnologia (FCT) for the PhD grant SFRH/BD/43738/2008. This study was partially funded by FCT, under research contract PTDC/AGR-CFL/117026/2010.

1. M Mota et al. (1999) *Nematology* 1:727–734.
2. A Abelleira et al. (2011) *Plant Dis* 95: 776.
3. P Barbosa et al. (2010) *J Nematol* 42: 8-16.
4. Ferreira et al. (2006) *Flavour Frag J* 22: 1-9.

## P 29. Chemotypification of *Astrantia major* L. (Apiaceae) inferred from its fruit essential oil profile

Radulović NS<sup>1</sup>; Mladenović MZ<sup>1</sup>; Đorđević ND<sup>1</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science and Mathematics, University of Niš

nikoradulovic@yahoo.com

Keywords: Essential oils; *Astrantia major*; Apiaceae; Saniculoideae; Multivariate statistical analyses; Chemotaxonomy

*Astrantia major* L. (family Apiaceae, subfamily Saniculoideae) is a perennial plant species found in the Pyrenees, Carpathian Mountains, Balkan Peninsula and Western Asia [1]. The only previous study [2] dealing with the chemical composition of the essential oil of this plant species reported the isolation and structural elucidation of three oxygenated sesquiterpenes ( $\beta$ -sinensal,  $\beta$ -sinensol and  $\beta$ -sinensyl acetate). We decided to reanalyze, in much more detail, the essential oil of the up to now uninvestigated populations of *A. major*. In total, seventy-six constituents identified by GC and GC/MS accounted for 92.7-94.0% of the oils from two populations (Serbia and Poland). The oils differed significantly - the wild-growing population from Serbia contained zingiberene (47.9%),  $\beta$ -bisabolene (9.7%) and  $\beta$ -sesquiphellandrene (7.9%), whereas the one from Poland (botanical gardens) was sesquiterpene-poor with oleic acid (38.6%), nonacosane (15.4%) and linoleic acid (5.1%) as the major contributors. Motivated by the unresolved taxonomical relations between the Saniculoideae and Apioideae subfamilies, we performed multivariate statistical analyses (agglomerative hierarchical cluster analysis (AHC) and principal component analysis (PCA)) on the compositional data of these *A. major* samples and additional 15 Saniculoideae and 31 Apioideae taxa. This allowed us to assess the chemotaxonomical usefulness of such chemical data in differentiating taxa from these two Apiaceae subfamilies and to corroborate the existence of at least two *A. major* chemotypes.

Acknowledgement: This work was funded by the Ministry of Education and Science of Serbia (Project 172061).

1. WJ Manning, B Godzik (2004) Environ. Pollut. 130: 33-39.
2. H Buurma, R Bos, DHE Tattje, JH Zwaving (1978) Phytochemistry 17: 2129-2130.

**P 30. Phenological variability of the alkaloids content in the essential oil of *Conium maculatum* L.**

Radulovic NS<sup>1</sup>, Đorđević ND<sup>1</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science and Mathematics, University of Niš, Višegradska 33, 18000 Niš, Serbia

nevenka\_83@yahoo.com

Keywords: *Conium maculatum*, essential oil, conmaculatin, 2-pentyl-3,4,5,6-tetrahydropyridine, coniceine, coniine.

*Conium maculatum* L. (Apiaceae), hemlock, is a worldwide distributed, highly poisonous weed. Up to now ten volatile alkaloids have been identified from this plant species [1,2]. Coniine and  $\gamma$ -coniceine are generally the most abundant ones and they account for the most of the plant's acute and chronic toxicity [2]. Recently, we reported the existence of a dihomologue of coniine (2-pentylpiperidine, conmaculatin) in the fruit essential oil of hemlock [3]. In this work, the content of coniine,  $\gamma$ -coniceine and their dihomologues was determined by detailed GC-MS analyses of twelve samples of hemlock and tracked during different phenological phases. The highest levels of  $\gamma$ -coniceine and 2-pentyl-3,4,5,6-tetrahydropyridine were detected in the first stages of phenological development (11.0-47.8% and 0.5-3.7%, respectively), and at the same time coniine (7.0-15.8%) and conmaculatin (0-0.7%) were present in significantly lower amounts. During fruit ripening, coniine (15.4-35.6%) and conmaculatin (0.7-1.3%) content increased substantially, whereas that of  $\gamma$ -coniceine (0.4-7.1%) and its dihomologue (0.1-0.4%) dropped accordingly. These results indicate that an analogous biosynthesical relationship between conmaculatin and 2-pentyl-3,4,5,6-tetrahydropyridine exist as for the coniine/ $\gamma$ -coniceine pair.

Acknowledgement: This work was funded by the *Ministry of Education and Science of Serbia* (Project 172061).

1. DG Lang, RA Smith (1998) *In*: T Garland, AC Barr (Eds.) Toxic plants and other natural toxicants. CAB International, Oxford, pp. 419-422.
2. T Reynolds (2005) *Phytochemistry* 66: 1399-1406.
3. N Radulović, N Đorđević, M Denić, GPM Martins, PF Dias, F Boylan (2012) *Food Chem. Toxicol.* 50: 274-279.

**P 31. *Myrtus communis* from Algeria: chemical composition and variability**

Bouzabata A<sup>1</sup>; Casanova J<sup>1</sup>; Castola V<sup>1</sup>; Abed L<sup>2</sup>; Bighelli A<sup>1</sup>; Tomi F<sup>1</sup>

<sup>1</sup>Université de Corse-CNRS, UMR 6134 SPE, Equipe Chimie et Biomasse, Route des Sanguinaires, 20000 Ajaccio, France

<sup>2</sup>Laboratoire de Pharmacognosie et Matière Médicale, Université Ben Youcef Benkhedda, Alger, Algérie.

amelbouz2009@gmail.com

Keywords: *Myrtus communis*; Algeria; chemical composition; variability.

The aim of the present study was to evidence either an homogeneity or a chemical variability within oil samples isolated from aerial parts of *Myrtus communis* L. growing wild in Algeria.

The chemical composition of 55 oil samples of *Myrtus communis* isolated from aerial parts harvested, at the flowering stage, in 16 locations from East to West Algeria was investigated by GC (FID), in combination with retention indices and by <sup>13</sup>C NMR spectroscopy. The results were submitted to statistical analysis.

The major components were  $\alpha$ -pinene (27.4-59.2%) and 1, 8-cineole (6.1-34.3%), and were characterized by the lack of myrtenyl acetate. Statistical analysis allowed the distinction of two groups including each one two sub-groups. Groups I (78% of the samples) and II may be distinguished on the basis of the content of  $\alpha$ -pinene, linalool and linalyl acetate. Subgroups IA and IB could be distinguished by their contents of  $\alpha$ -pinene and 1,8-cineole. Subgroups IIA and IIB differed substantially by their contents in cineole and limonene.

According to the present and previous results, there is no correlation between pedo-climatic factors and the chemical composition of Algerian myrtle leaf oil. Taking into account these results as well as literature data on the composition of myrtle oils from North Africa, we remark that all the reported Moroccan oil samples contained myrtenyl acetate, and in contrast, oils samples from Algeria and Tunisia were characterized by the lack of myrtenyl acetate. As far the information was provided, it appeared that all the Moroccan myrtle oils were isolated from plants collected in the west part of Morocco and therefore, it could be hypothesized that Atlas Mountains constitute the natural limit between myrtle characterized by the presence or by the lack of myrtenyl acetate.

Acknowledgments: A. Bouzabata wishes to thank Dr G. De Belaire (Badji-Mokhtar University, Annaba, Algeria) for identification of the plant.



### P 32. Determination of minimal inhibitory and microbicidal concentrations of essential oils produced by Mediterranean plants

Nunes MJC<sup>1</sup>, Veríssimo CSC<sup>1</sup>, Valente F<sup>1</sup>, Santos MI<sup>2</sup>, Pedroso L.<sup>3</sup>, Sousa I<sup>1</sup>, Ferreira MASS<sup>1</sup>

<sup>1</sup>Instituto Superior de Agronomia, Universidade Técnica de Lisboa

<sup>2</sup>Instituto Superior de Agronomia, Universidade Técnica de Lisboa, Faculdade de Medicina Veterinária, Universidade Lusófona de Humanidades e Tecnologias

<sup>3</sup>Faculdade de Medicina Veterinária, Universidade Lusófona de Humanidades e Tecnologias

missantos@gmail.com

Keywords: essential oils; antimicrobial effect; pathogenic microorganisms

The aim of this work was to evaluate the antimicrobial effect of essential oils, for its possible application on salads to control the contaminant microbial flora. Methods: Essential oils of oregano (*Origanum vulgare*), three different sages (*Salvia officinalis*, *Salvia lavandulaefolia* and *Salvia sclarea*) and rosemary (*Rosmarinus officinalis*) were analyzed to evaluate its antimicrobial action. These oils were tested by two methods against *Yarrowia lipolytica*, *Pseudomonas aeruginosa* and *Listeria monocytogenes*. First it was determined the minimum inhibitory concentration (MIC) of -all the essential oils, by the micro dilution method on 96 well plates, at 25 or 37°C respectively. The determination of MIC was done by measuring the growth absorbance at 540 nm in all micro plates. To evaluate the most suitable oils for future work, statistical analyses were performed using ANOVA and Tukey tests with the R software. After MIC determination, for results scaling up purposes, it was necessary to study the minimum microbicidal concentration (MMC) by the dilution method in liquid medium. The MMC were tested using three of the five initial essential oils that presented the best performance: *Origanum vulgare*, *Rosmarinus officinalis* and *Salvia lavandulaefolia*. Spreading inoculations were done on agar plates for surviving cells count and the variability of the obtained calculations was evaluated. The inhibitory concentrations of the essential oils tested are discussed.

All the microbial strains demonstrated the most sensitivity to the *Origanum vulgare* essential oil and the least sensitivity to the *Salvia* spp. essential oils. The antimicrobial activity of *Origanum vulgare* oil is effective against *Yarrowia lipolytica*, its MMC correspond to 0.235% (v/v). However, it was found that *Pseudomonas aeruginosa* has a low sensitivity to the antimicrobial effect of these essential oils. The results obtained with *Listeria monocytogenes* for *Rosmarinus officinalis* showed a MMC of 3.75% (v/v).

The results obtained are promising for the intended application but further studies should be done.

### P 33. Chemical composition and antinociceptive properties of the essential oil of *Ferula ovina* (Umbelliferae)

Radulovic NS<sup>1</sup>; Zlatkovic DB<sup>1</sup>; Stojanovic NM<sup>2</sup>; Akhlaghi H<sup>3</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science and Mathematics, University of Nis, Visegradaska 33, Nis, Serbia

<sup>2</sup>Faculty of Medicine, University of Nis, Bulevar dr Zorana Djindjica 81, Nis, Serbia

<sup>3</sup>Department of Basic Sciences, Islamic Azad University, Sabzevar Branch, Sabzevar, Iran

dragan4@gmail.com

Keywords: *Ferula ovina*; essential oil; vanillates; Umbelliferae

*Ferula* L. is a genus of about 170 species of Umbelliferous flowering plants native to the Mediterranean region and east to central Asia. The Iranian flora comprises 30 species of this genus, one of them being *Ferula ovina* (Boiss.) Boiss. The aqueous extracts of *F. ovina* was found to possess antispasmodic, anticholinergic and smooth muscle relaxant activities [1]. Although the essential oil composition of *F. ovina* was previously reported [2,3], the potential pharmacological activity of this oil remains unknown up to date. Having this and the ethnopharmacological use of this plant species in mind, we decided to assess the antinociceptive activity of the essential oil of the aerial parts of *F. ovina* (collected in Iran) using the hot plate and tail immersion methods for central and the abdominal writhing test for peripheral analgesia in BALB/c mice. In the highest applied dose, the oil showed a strong peripheral analgesic activity by causing a 92.4% decrease in the number of writhes induced by acetic acid. On the other hand, the oil exerted only moderate central analgesic activity, reaching 55.8% of inhibition in the tail immersion test. These interesting results provoked us to perform detailed compositional analyses of the oil. GC and GC-MS analyses allowed the identification of 120 constituents. The oil consisted mainly of monoterpene hydrocarbons. The major constituents of this oil were camphene (18.0%), limonene (14.3%) and myrcene (13.5%). The oil also contained a number of rare esters of borneol. The structures of these esters were corroborated by synthesis. Bornyl *p*-methoxybenzoate turned out to be a new natural compound. Currently, additional work is underway to determine the active principles of this essential oil.

Acknowledgments: This work was funded by the Ministry of Education and Science of Serbia (Project 172061).

1. S Al-Khalil et al. (1990) J. Ethnopharmacol. 30(1): 35-42.
2. A Ghannadi et al. (2002) DARU 10(4): 165-167.
3. B Rahmani et al. (2008) J. Ess. Oil Res. 20(3): 232-235.

### P 34. Effect of *Origanum vulgare* essential oil on *Penicillium digitatum* using warm air flow

Smid J; Kloucek P; Finkousova A; Frankova A

Czech University of Life Sciences

smidj@af.czu.cz

Keywords: oregano; essential oil; antimicrobial; postharvest diseases; lemon; shelf-life

Antimicrobial activity of essential oils (EOs) is well known, but still is not widespread in crop products treatment. Because of high costs for treatment with EOs, new and cheaper methods are recommended. *Penicillium digitatum* is one of the main fungus causing storage losses on citruses and *Origanum vulgare* EO is known for its antimicrobial properties [1]. Method for treatment with essential oils vapours was greatly improved. Gas chromatograph (5890A GAS Chromatograph, Hewlett Packard) was modified as a warm air flow treating box (oven temperature: 40°C). Glass cube was inserted into the oven and the detector heater was ejected into the oven and used for vaporization of EO (detector temperature: 150°C). Antimicrobial activity of *O. vulgare* EO was tested in vitro and in model system on *Citrus limon* fruits. Inoculum was made from fourteen days old cultures of *P. digitatum* dissolving in Tris buffered saline with tween 80.

Six inoculated Petri dishes (55 mm diameter) or six *C. limon* fruits inoculated with *P. digitatum* in three artificial wounds (5 mm diameter) were placed on the bottom of glass cube. Both were kept in warm air flow for five minutes. EO from *O. vulgare* in concentrations 0.5; 2; 4 and 8  $\mu\text{l.l}^{-1}$  of air was used. Carvacrol (64,6%), para – cymene (5,2%) and thymol (2,9%) are the main components identified by GC-MS. The radial growth of *P. digitatum* on *C. limon* fruits were measured and Petri dishes were checked for *P. digitatum* growth after 4 days of incubation in 25°C. All tests were conducted in triplicate. Growth of *P. digitatum* on Petri dishes was totally inhibited by EO concentration 4  $\mu\text{l.l}^{-1}$ . EO in 8, 4, 2 and 0.5  $\mu\text{l.l}^{-1}$  concentrations reduced *P. digitatum* radial growth on *C. limon* fruits to 39, 46, 70 and 79 % respectively.

Previous in vitro experiments show inhibition in concentrations about 400  $\mu\text{l.l}^{-1}$ [2]. The method using activation of EO with temperature and air flow rapidly reduces effective concentration (100x). Moreover this method could be improved for mass production and is suitable for any fruits and vegetables.

Acknowledgement: This presentation is supported by University-wide internal grant agency - CIGA 20122008 – Plant products treatment with essential oil vapours under different physical conditions.

1. S Burt (2004) Int. J. Food Microbiol. 94/3, 223-253.
2. DJ Daferera et al. (2000) J. Agric. Food Chem. 48/6: 2576-2581.

**P 35. Essential oil composition of juniper berry (*Juniperus communis* L., Cupressaceae) from Macedonian flora**

Karapandzova M<sup>1</sup>; Sela F<sup>1</sup>; Stefkov G<sup>1</sup>; Kulevanova S<sup>1</sup>

<sup>1</sup>Department for Pharmacognosy, Faculty of Pharmacy, Skopje, The Former Yugoslav Republic of Macedonia

svku@ff.ukim.edu.mk

Keywords: *Juniperus communis*; essential oil composition; GC/FID/MS.

The berries of common juniper, *Juniperus communis* L. (Cupressaceae), is used for isolation of juniper oil, which has multi-purpose medicinal use as diuretic, antiseptic, stomachic, antireumatic, etc. [1]. Among medicinal, there is also a large commercial use in food industry and cosmetic and perfume production. The aim of the present study was determination of the chemical composition of Macedonian juniper berry oil in correlation to the areas of collection. The samples of berries were collected on twelve different locations in Republic of Macedonia. The essential oils were isolated by steam-distillation in the Clevenger-type apparatus using official method from European pharmacopoeia [2]. The chemical composition of essential oil samples was analyzed by gas chromatography system equipped with flame ionization detector and mass spectrometer as well as capillary flow technology. For that purpose, HP-5ms capillary column was used. Seventy-four components were identified representing 91.08 to 99.83% of the entire oils. The predominant fractions of the oils were monoterpene hydrocarbons presented from 39.11 to 73.38% of total oil composition. Great variability in the chemical composition and the content of some components was registered, depending of the area of collection. The most variable were  $\alpha$ -pinene (15.59-43.19%),  $\beta$ -pinene (1.65%-5.35%),  $\beta$ -myrcene (2.89%-26.50%), sabinene (2.80-11.77%) and limonene (2.90-4.46%). In the fraction of oxygen-containing monoterpenes the most abundant was terpene-4-ol (0.02-6.32%), while in the fraction of sesquiterpenes germacrene D (2.03-10.22%),  $\beta$ -elemene (1.43-6.40%) and *trans*-(E)-caryophyllene (1.8%-4.05%). The differences in the environmental conditions of plant growth and the characteristics of the locations where the collections were made, were considered as important factors for the variability in the essential oil composition of Macedonian juniper.

1. WC Evans (2002) *Trease and Evans Pharmacognosy*, 15<sup>th</sup> edition, W.B. Saunders, Edinburg, London, New York, Philadelphia, St Louis, Sydney, Toronto.
2. European Pharmacopoeia (2011), 7<sup>th</sup> edition, Council of Europe, Strasbourg.

### P 36. Glandular trichomes, histochemical localization of secretion and essential oil composition in *Plectranthus grandidentatus*

Mota L; Figueiredo AC; Pedro LG; Barroso JG; Ascensão L

Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, IBB, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portugal

lbmota@gmail.com

Keywords: *Plectranthus grandidentatus*; Lamiaceae; glandular trichomes; histochemistry; essential oils

A large number of *Plectranthus* species are well known for their medicinal uses, while others are very popular as ornamentals [1]. *Plectranthus grandidentatus* Gürke is an herbaceous perennial herb commonly used as groundcover due to its procumbent growth habit and pleasant scented leaves. Although this species was not reported as medicinal, their abietane diterpenoids showed activity against methicillin- and vancomycin-resistant bacteria [2].

As part of a ongoing study on *Plectranthus* species [3,4,5], we report herewith the morphology and histochemistry of the glandular trichomes, and the essential oils composition from *P. grandidentatus* growing in Portugal.

The types of glandular trichomes and their pattern of distribution on the leaves and flowers were studied by LM and SEM as in [3]. The stems and leaves of *P. grandidentatus* are covered by a tomentose indumentum composed of long, uniseriate and multicellular nonglandular trichomes. Five distinct types of glandular trichomes were identified. Peltate trichomes showed an orange characteristic colour and never developed a large subcuticular space. Capitulate trichomes were divided in three subtypes according to the dimension of the glandular head, length of stalk and secretion process. The flower exhibited also uncommon conoidal trichomes. The histochemical characterization of the secretion revealed the presence of terpenoids and flavonoids as the main constituents.

The essential oils, isolated by hydrodistillation of dried leaves and stems, were analyzed by GC and GC-MS as in [6]. The oils yields ranged from 0.02 to 0.05% (v/w). A total of 70 components were identified, representing more than 84% of the oils. Sesquiterpenes constituted the dominant fraction (58-68%), being  $\beta$ -caryophyllene,  $\alpha$ -santalene and  $\alpha$ -humulene the most abundant hydrocarbons and  $\beta$ -caryophyllene oxide and ledol the major oxygen-containing compounds. Monoterpene fraction did not exceed 10% and only  $\alpha$ -pinene and *trans*- $\beta$ -ocimene reached percentages above 1%.

The data obtained showed the high diversity of *P. grandidentatus* glandular trichomes and, as far as we know, this is the first report on their secretory structures and essential oils.

1. CW Lukhoba et al. (2006) *J. Ethnopharmacol.* 103: 1-24.
2. C Gaspar-Marques et al. (2006) *Phytomedicine* 13: 267-271.
3. L Ascensão et al. (1999) *Ann. Bot.* 84: 437-447.
4. S.Porfírio et al. (2010) *Food Chem.* 122:179-187.
5. PL Falé et al. (2011) *Food Function* 2: 130-136.
6. MD Mendes et al. (2011) *Ind. Crop. Prod.* 33: 710-719.

**P 37. Chemical and statistical analysis of *Achillea clypeolata* Sibth. & Sm. essential oils: plant organ-dependent biosynthesis/accumulation of the volatiles**

Blagojević P<sup>1</sup>; Stanković S<sup>1</sup>; Radulović N<sup>1</sup>

<sup>1</sup>University of Niš, Faculty of Science and Mathematics, Serbia

blagojevicpolina@gmail.com

Keywords: *Achillea clypeolata* Sibth. & Sm.; Asteraceae; essential oil; above-ground parts; multivariate statistical analysis

*Achillea* L. (Asteraceae) is a genus of about 85 flowering plants, commonly referred to as *yarrows* that occur in Europe and temperate areas of Asia. A number of *Achillea* species were the subject of extensive phytochemical studies, mainly due to their (ethno)pharmacological importance [1-4]. *Achillea clypeolata* Sibth. & Sm., an endemic species to the Balkan Peninsula and Asia Minor, is among those members of this genus that have received little attention and only several papers can be found in the literature dealing with *A. clypeolata* volatile secondary metabolites [1-3]. It has been previously reported that collections of *A. clypeolata* growing in different regions of Bulgaria differ in their chemical constituents to such an extent that the existence of chemotypes might be suggested [4]. Bearing this in mind, and in continuation of our chemical investigations of Asteraceae species used in folk medicine, we have now studied five Serbian populations of *A. clypeolata*. Analyses (GC and GC-MS) of 10 essential oil samples hydrodistilled from dry flowers and vegetative above-ground parts of *A. clypeolata* enabled the identification of more than 100 different constituents, representing 90-95% of the total oils. The most abundant constituents of the obtained oils were 1,8-cineole (0.1-34.8%), *trans*-verbenol (0-38.2%), camphor (0-17.9%), borneol (0-6.9%), *p*-mentha-1,5-dien-8-ol (0-9.5%), 4-terpineol (1.2-8.1%), germacrene D (1.3-9.8%), elemol (0.4-14.85%), g-eudesmol (trace-8.8%) and b-eudesmol (1.4-13.3%). As revealed by chemical analyses and a multivariate statistical treatment of the oils (agglomerative hierarchical cluster analysis and principal component analysis), the biosynthesis and/or accumulation of volatile metabolites in *A. clypeolata* is strongly plant organ-dependent. In addition, the obtained results showed that the studied populations do not belong to a single *A. clypeolata* chemotype. Moreover, they suggest that the volatile profile of the studied species is susceptible to environmental and climatic factors.

Acknowledgments: This work was funded by the Ministry of Education and Science of Serbia (Project 172061). PB was supported by UNESCO-L'OREAL National Fellowships Programme in Serbia.

1. M Maffei et al. (1994) *Biochem. Syst. Ecol.* 22: 679-687.
2. N Simic et al. (2005) *Flavour Fragr. J.* 20: 127-130.
3. JC Chalchat et al. (2005) *J. Essent. Oil. Res.* 17: 549-552.
4. MN Todorova, ET Asankova (1999) *Phytochemistry* 52: 1515-1518.

**P 38. Effects of methyl and isopropyl *n*-methylantranilates from the essential oil of *Choisya ternata* Kunth (Rutaceae) on experimental anxiety and depression in mice**

Radulovic NS<sup>1</sup>; Miltojevic AB<sup>1</sup>; Randjelovic PJ<sup>2</sup>; Stojanovic NM<sup>3</sup>; Boylan F<sup>4</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science and Mathematics, University of Niš

<sup>2</sup>Department of Physiology, Faculty of Medicine, University of Niš

<sup>3</sup>Faculty of Medicine, University of Niš

<sup>4</sup>Panoz Institute, School of Pharmacy and Pharmaceutical Sciences, Trinity College Dublin 23, Westland Row, Dublin 2, Ireland

nikola.st90@yahoo.com

Keywords: methyl N-methylantranilates; isopropyl N-methylantranilates; *Choisya ternata* Kunth; essential oil; light/dark test; tail suspension test

*Choisya ternata* Kunth (Rutaceae) is a plant species whose leaves are used in Mexican folk medicine for its antispasmodic and stimulative properties. Up to now, the essential oil of this plant species (Mexican orange) has been investigated on one previous occasion with only eighteen reported components [1]. This motivated us to perform detailed GC and GC-MS analyses of the hydrodistillate from *C. ternata* leaves. The essential oil turned out to be a complex mixture of compounds of which 157 were identified (the extent of identification of components being more than 98% of the total detected GC peak areas). A characteristic of the oil was the presence of two alkaloids - isopropyl and methyl *N*-methylantranilates. Their structures were confirmed by GC co-injection of the synthesized standards with the essential oil of Mexican orange. Motivated by the ethnopharmacological use and availability through synthesis, we decided to evaluate the anxiolytic and antidepressant effects of the two volatile anthranilates in open field, horizontal wire, light/dark, forced swimming and tail suspension tests, as well as their effect on the onset and duration of diazepam-induced sleep in BALB/c mice. The volatile alkaloids (50-200 mg/kg, *i.p.*), without having a muscle relaxant effect, caused a significant decrease in the time the animals spent in an unsecured and putatively dangerous area when compared with the control group, but had no effect on the number of crossings between the light/dark compartments. In addition to this anxiolytic activity, a significant antidepressant-like effect was apparent at all tested doses, which was not due to an increase in locomotive activity. The anthranilates administered on their own did not induce sleep in mice, but significantly prolonged the diazepam-induced sleep, in a dose dependent way, suggesting an interaction with the GABA receptor complex. The obtained data at least partially provide a justification for the ethnomedicinal usage of Mexican orange and point to the two volatile alkaloids as the active principles from the essential oil.

Acknowledgments - This work was funded by the Ministry of Education and Science of Serbia (Project 172061).

1. MJ Respaud et al. (1997) J. Essent. Oil. Res. 9: 475-476.

### P 39. Chemotaxonomy of Balkan *Geranium* and *Erodium* species inferred from the essential oil chemical composition

Radulović N<sup>1</sup>; Dekić M<sup>2</sup>; Stojanović-Radić Z<sup>3</sup>; Randelović V<sup>3</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science and Mathematics, University of Niš, Višegradska 33, 18000 Niš, Serbia

<sup>2</sup>Department of Chemical and Technological Sciences, State University of Novi Pazar, Vuka Karadžića bb, 36300 Novi Pazar, Serbia

<sup>3</sup>Department of Biology and Ecology, Faculty of Science and Mathematics, University of Niš, Višegradska 33, 18000 Niš, Serbia

mdekic@np.ac.rs

Keywords: Geraniaceae; *Geranium*; *Erodium*; Essential oil; Chemotaxonomy; Multivariate statistical analysis

The genus *Geranium* L. (Geraniaceae) comprises ca. 250 species, distributed mainly in the temperate region of the northern hemisphere, but no comprehensive taxonomic treatment is nowadays available [1]. A closely related genus of this plant family, *Erodium* L' Hérit. includes more than 50 species, of which, about 40 are widespread across the Mediterranean region. According to Janković [1], nineteen *Geranium* and two *Erodium* species can be found in the Serbian flora.

Very little attention was paid to the species of these genera with respect to their volatile constituents. In general, reports regarding the essential oils of only few species exist in the literature: *G. macrorrhizum* [2-3], *G. robertianum* [4], *G. phaeum* [2] and *E. cicutarium* (leaf hexane extract [5]).

Prompted by the lack of such compositional data, we decided to perform detailed GC-MS analyses of the essential oils isolated from seven *Geranium* (*G. macrorrhizum*, *G. robertianum*, *G. sanguineum*, *G. columbinum*, *G. lucidum*, *G. purpureum* and *G. phaeum*) and three *Erodium* (*E. cicutarium*, *E. ciconium* and *E. absinthoides*) species from Serbia and Macedonia. This resulted in the identification of 551 and 233 constituents in total, respectively. The significance of the variations in essential oil composition/production was determined using multivariate statistical analyses (agglomerative hierarchical clustering analysis and principal component analysis). The main conclusion, based on the dendograms and biplots obtained as the results of the performed analyses, using relative abundances of the identified oil constituents, consists in that there is no great intergeneric variability in the oil compositions, i.e. the chemical composition of the investigated *Geranium* and *Erodium* taxa was generally very similar. These results confirm the close phylogenetic relationship between the two genera.

Acknowledgment: This work was supported by the Ministry of Education and Science of Serbia [Project No. 172061].

1. MM Janković (1973) In: M Josifović (Ed.), Flora of Serbia V. SANU, Beograd, pp.136–160.
2. J-C Chalchat, SD Petrović, ZA Maksimović, MS Gorunović (2002) J. Essent. Oil Res. 14: 333–335.
3. NV Bozhkova, G Stoev, AS Orahovats, N Rizov (1984) Phytochemistry 23: 917–918.
4. LG Pedro, MSS Pais, JJC Scheffer (1992) Flavour Fragr. J. 7: 223–226.
5. M Lis-Balchin (1993) J. Essent. Oil Res. 5: 317–318.



**P 40. Antimicrobial volatile glucosinolate autolysis products from *Hornungia petraea* (L.) Rchb. (Brassicaceae)**

Radulović N<sup>1</sup>; Dekić M<sup>2</sup>; Stojanović-Radić Z<sup>3</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science and Mathematics, University of Niš, Višegradska 33, 18000 Niš, Serbia

<sup>2</sup>Department of Chemical and Technological Sciences, State University of Novi Pazar, Vuka Karadžića bb, 36300 Novi Pazar, Serbia

<sup>3</sup>Department of Biology and Ecology, Faculty of Science and Mathematics, University of Niš, Višegradska 33, 18000 Niš, Serbia

mdekcic@np.ac.rs

Keywords: Brassicaceae; *Hornungia petraea*; Glucosinolate; Isothiocyanate; Antimicrobial activity

The genus *Hornungia* comprises approx. 10 species, distributed in Europe, North Africa and west Asia. In Serbia, the genus is represented by only one species – *Hornungia petraea* (L.) Rchb. (syn. *Hutchinsia petraea* (L.) R. Br.) [1].

In general, only two previous references on the phytochemistry of *Hornungia* and the closely related *Hutchinsia* taxa can be found regarding the analysis of glucosinolates (GLS) and their volatile breakdown products in *Hutchinsia alpina* [2,3].

Recently we reported the results of a study of volatile GLS autolysis metabolites in plant samples of *H. petraea* [4]. GC–MS analysis of the autolysate and the synthesis of a series (12 compounds) of possible GLS breakdown products revealed/corroborated the presence of glucoaubrietin, glucolimnanthin, glucolepigramin and glucotropaeolin in this species as the most likely “mustard oil” precursors.

According to the classification of GLS-containing species based on GLS profiles (subdivision of the crucifers based on their GLS content into four groups) [3], this taxon and *Hutchinsia alpina*, a species closely related to *H. petraea*, the only studied from this genus, could be recognized as crucifers with the predominance of simple aromatic GLSs. The similarity of the GLS profile of the two phylogenetically close taxa provides support for the suggestion of Appel and Al-Shehbaz [5] to reduce the genus *Hutchinsia* (= *Pritzelago*) to synonymy of *Hornungia*, i.e. to treat *H. alpina* as a species of the genus *Hornungia*.

GLS degradation products identified in the autolysate of *H. petraea*, benzyl, 3- and 4-methoxybenzyl isothiocyanate, along with several other structurally related compounds were evaluated for antimicrobial activity in order to possibly pinpoint the role of the latter secondary metabolites in the plant tissues. The assays showed a very high antibacterial activity of the tested isothiocyanates against *Sarcina lutea* and an antifungal effect against *Aspergillus fumigatus* and *Candida albicans* with MIC values in the order of 1 mg/ml value.

Acknowledgment: This work was supported by the Ministry of Education and Science of Serbia [Project No. 172061].

1. R Jovanović-Dunjić (1972) In: M Josifović (Ed.) Flora of Serbia, vol. 3. SANU, Belgrade, pp. 339–340.
2. A Kjaer et al. (1953) Acta Chem. Scand. 7: 1276–1283.
3. RN Bennett et al. (2004) J. Agric. Food Chem. 52: 428–438.
4. NS Radulović, MS Dekić, ZZ Stojanović-Radić (2012) Phytochem. Lett. 5: 351–357.
5. O Appel, A Al-Shehbaz (1997) Novon 7: 338–340.

#### P 41. Essential oils composition of some Asteraceae species from Eastern Amazon, Brazil

Carvalho EEN<sup>1</sup>; Domingos VD<sup>2</sup>; Ribeiro AF<sup>3</sup>; Carreira LMM<sup>4</sup>; Lucas FCA<sup>5</sup>; da Silva JKR<sup>3</sup>; Andrade EHA<sup>3</sup>; Maia JGS<sup>3</sup>

<sup>1</sup>Universidade Federal do Tocantins

<sup>2</sup>Instituto Federal de Educação, Ciência e Tecnologia do Tocantins

<sup>3</sup>Universidade Federal do Pará

<sup>4</sup>Museu Paraense Emílio Goeldi

<sup>5</sup>Universidade Estadual do Pará

[gmaia@ufpa.br](mailto:gmaia@ufpa.br)

Keywords: Asteraceae species, essential oil variation, Eastern Amazon

In Brazil, mainly in the Amazon, the Asteraceae are represented by ca. 180 genera, most of which consists of herbaceous plants. The family is considered one of the most important sources of species with therapeutic value, many of which have been studied chemically and pharmacologically. Specimens of *Tilesia baccata* (L.) Pruski (1), *Vernonia scabra* Pers (2), *Ichthyothere cunabi* Mart. (3), *Blainvillea dichotoma* (Murray) A. Stewart (4), *Ageratum conyzoides* L. (5), *Unxia camphorata* L. (6), *Riencourtia angustifolia* Gardner (7), *Eupatorium squalidum* DC. (8) and *Ichthyothere rufa* Gardner (9) were collected in the municipality of São Geraldo do Araguaia (1-5) and Carajás National Forest (6,7), Pará state, and in the municipalities of Taquaruçu (8) and Palmas (9), Tocantins state. The essential oils of these species were hydrodistilled and analyzed by GC and GC-MS. The main constituents (over 4%) of *T. baccata* oil were  $\beta$ -bisabolene (16.0%),  $\beta$ -caryophyllene (7.9%),  $\gamma$ -elemene (7.6%) and spathulenol (6.8%); in the *V. scabra* oil were caryophyllene oxide (17.1%), (*E*)-caryophyllene (16.9%), germacrene D (8.7%),  $\alpha$ -humulene and (6.0%); in the *I. cunabi* oil were dehydrosesquiceneole (17.9%),  $\delta$ -elemene (11.0%), bicyclogermacrene (10.0%), spathulenol (5.7%), germacrene D (5.5%) and  $\beta$ -elemene (4.4%); in the *B. dichotoma* oil were (*E*)-caryophyllene (17.9%), spathulenol (5.9%), caryophyllene oxide (7.3%), (*E*)-nerolidol (5.5%),  $\delta$ -cadinene (5.1%),  $\gamma$ -elemene (4.5%) and bicyclogermacrene (4.1%); in the *A. conyzoides* oil were *epi*- $\alpha$ -cadinol (31.3%), *trans*- $\beta$ -guaiene (11.7%),  $\alpha$ -cadinol (10.7%) and  $\alpha$ -santalene (7.0%); in the *U. camphorata* oil were  $\alpha$ -phellandrene (27.6%), camphor (15.0%), (*E*)-caryophyllene (11.6%), thymol methyl ether (7.0%) and germacrene D (4.4%); in the *R. angustifolia* oil were  $\delta$ -elemene (15.4%), bicyclogermacrene (12.9%), geranyl acetate (10.0%), geraniol (9.5%), (*E*)-caryophyllene (7.3%), terpinolene (5.2%), germacrene D (5.2%) and *epi*- $\alpha$ -cadinol (4.9%); in the *E. squalidum* oil were  $\beta$ -elemene (21.6%), caryophyllene oxide (5.8%),  $\alpha$ -humulene (5.3%), humulene epoxide II (4.9%); and in the *I. rufa* oil were sabinene (33.1%), terpinen-4-ol (28.1%), limonene (6.7%) and  $\gamma$ -terpinene (6.4%).

Acknowledgments: This work was supported by BIONORTE/CNPq and FAPESPA/PA.

## P 42. The minor constituents of modern roses

Sakurai K; Atoji H; Kawaraya A

Takasago International Corporation

kazutoshi\_sakurai@takasago.com

Keywords: Ambridge rose, Cymbeline, Lady Hillingdon, Alkylpyridines, 6-hydroxydihydrothespirane

The volatile constituents of English roses (Ambridge rose & Cymbeline) were analyzed using Aromascope® technology followed by GC-MS analysis.

As the major components of both of the flowers, 4-methoxystyrene, 1,3-dimethoxy-5-methylbenzene(DMMB), geraniol, nerol, citronellol, cis-3-hexenyl acetate, anisaldehyde, and citral (neral & geranial) were detected and from Ambridge rose total 118 components then from Cymbeline total 154 components were identified.

Among them, three kinds of alkylpyridines, such as 3-ethylpyridine(1), 5-ethyl-2-methylpyridine (2) and 2,6-dimethylpyridine (3) the nitrogen containing components were identified from both of English roses for the first time. These alkylpyridines have already reported as the one of the key components from the basic fraction in peppermint oil<sup>1)</sup> and jasmine absolute<sup>2)</sup>.

Then, the small addition of these alkylpyridine derivatives 1-3 to the rosy fragrance made to enhance the natural flowerscent and a good effect for the rosy fragrance.

In addition, from Lady Hillingdon (the most popular representative tea-scented modern rose), 6-hydroxydihydrothespiranes (2,6,6,10-tetramethyl-1-oxaspiro[4,5]decan-10-ols) (4), considering as the precursor of the thespiranes or their intermediate from  $\beta$ -ionol,  $\beta$ -ionone and dihydro- $\beta$ -ionol, were identified for the first time from this rose flower. The compounds 4 have already found in the *Osmanthus* absolute<sup>3)</sup>, black tea<sup>4)</sup> and green tea<sup>5)</sup> as racemate.

These compounds 4 showed the woody camphorous and very strong earthy note. In this study, only two enantiomers were detected on GC and chiral GC analysis.

1. K.Sakurai, *et al.*, *Agric.Biol.Chem.*, 1983, 47, 2307- 2312.

2. T.Toyoda, *et al.*, *Agric.Biol.Chem.*, 1978, 42, 1901-1905.

3. R.Kaiser, *et al.*, The abstract paper of 7<sup>th</sup> International Congress of Essential oils, 1977, 395- 399.

4. T.Yamanishi, *et al.*, *Agric.Biol.Chem.*, 1966, 30, 1102-1105.

5. W.Renold, *et al.*, *Helv Chim Acta*, 1974, 57, 1301-1308.

**P 43. Influence of leaf drying and storage of the essential oil of *Ocimum basilicum* cultivar Maria Bonita on its chemical constituents**

Alves MF<sup>1</sup>; Blank AF<sup>1</sup>; Fontes SS<sup>1</sup>; Arrigoni-Blank MF<sup>1</sup>; Jesus HCR<sup>2</sup>; Alves PB<sup>2</sup>

<sup>1</sup>*Departamento de Engenharia Agrônômica, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil.*

<sup>2</sup>*Departamento de Química, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil.*

*arie.blank@gmail.com*

Keywords: basil; essential oil; post-harvest; linalool

The objective of this study was to evaluate the influence of leaf drying and oil storage on the content and chemical composition of the essential oil of the Maria Bonita cultivar of basil. In the first trial, the effect of the drying time of leaves in an oven with forced-air circulation was evaluated at a temperature of 40°C. In the second trial, the effect of storage time was evaluated (0, 15, 30, 60, 90, 120, 150, 210, 240 and 270 days) at two temperatures [room ( $\pm 27^\circ\text{C}$ ) and freezer ( $-20^\circ\text{C}$ ) temperature]. The essential oil was obtained from dry leaves by hydrodistillation using a Clevenger apparatus. Qualitative analysis of the chemical composition of the essential oils was performed using a gas chromatograph coupled to a mass spectrometer (GC-MS). A quantitative analysis of the essential oil components was conducted in a gas chromatograph equipped with a flame ionization detector (FID). The drying process at 40°C was efficient, reducing the moisture content of basil from 84.5% to 1.3% over a period of eight days. There was a linear reduction in the essential oil content from 6.0% to 3.9% during the process of leaf drying at 40°C. We also observed that the drying time proportioned decrease of linalool content from 76.38% to 74.09%, and increase of the content of  $\alpha$ -*trans*-bergamotene (from 1.1% to 1.8%) and epi- $\alpha$ -cadinol (from 1.57% to 1.77%). In the second trial, we noted increase of the linalool content from 76.99% to 79.40% after 210 days of storage at room temperature and to 79.82% after 240 days of storage in freezer. We can conclude that basil essential oil can be stored for up to seven months at room temperature and eight months in freezer.

Acknowledgments: The authors thank FAPITEC/SE, CNPq and CAPES for their financial support of this work.

**P 44. Influence of time and temperature of storage of the essential oil of *Pogostemon cablin* on its chemical constituents**

Santos PS<sup>1</sup>; Arrigoni-Blank MF<sup>1</sup>; Blank AF<sup>1</sup>; Brito AF<sup>1</sup>; Gomes Neto SC<sup>2</sup>; Santos DA<sup>3</sup>; Alves PB<sup>3</sup>

<sup>1</sup>*Departamento de Engenharia Agrônômica, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil.*

<sup>2</sup>*Raros Naturals Company, CEP 59280-000, Macaíba, Rio Grande do Norte, Brasil.*

<sup>3</sup>*Departamento de Química, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil.*

*fatima.blank@gmail.com*

Keywords: patchouli; aromatic plant; volatile oils; postharvest; patchoulol

The aim of this work was to investigate the effect of the time (0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, and 12 months) and temperature (room ( $\pm 27^{\circ}\text{C}$ ) and freezer ( $-20^{\circ}\text{C}$ ) temperature) of storage on the chemical composition of patchouli (*Pogostemon cablin* (Blanco) Benth.) essential oil. For this purpose, the essential oil of patchouli accession POG-002 was extracted from dry leaves at room temperature by steam distillation. The chemical composition was determined by gas chromatography-mass spectrometry (GC-MS), and the essential oil was quantified by gas chromatography with flame ionization detection (GC-FID). Fifteen of the components present in the essential oil were identified. Patchoulol was the main component of the oil; its concentration ranged from 36.36% to 42.12% at room temperature and from 35.19% to 42.93% at  $-20^{\circ}\text{C}$ . These results indicate that the essential oil of patchouli may be stored at room temperature and at  $-20^{\circ}\text{C}$  for 11 months without affecting its quality.

Acknowledgments: The authors thank FAPITEC/SE, CNPq and CAPES for their financial support of this work.

## P 45. Antifungal activity and analysis of volatile components of *Origanum virens* Hoffmans & Link

Silva LA<sup>1</sup>; Delgado FG<sup>2</sup>; Tinoco MT<sup>3</sup>; Rodilla JM<sup>1</sup>; Gomes AC<sup>1</sup>

<sup>1</sup>Departamento de Química, Universidade da Beira Interior, Rua Marquês d'Ávila e Bolama, 6200-Covilhã, Portugal

<sup>2</sup>Escola Superior Agrária, Quinta da Sra. de Mércules 6001-099 Castelo Branco, Portugal

<sup>3</sup>Departamento de Química, Universidade de Évora, Rua Romão Ramalho, 7000-Évora, Portugal

[mlas@ubi.pt](mailto:mlas@ubi.pt)

Keywords: *Origanum virens*; antifungal activity; essential oils

*Origanum virens* Hoffmans & Link is an aromatic plant which grows wild in some regions from Portugal. In the present paper, we analysed the chemical composition of essential oils from fresh and dried leaves of *O. virens* and their antifungal activity against strains of *Aspergillus niger*, *Penicillium* sp and *Fusarium oxysporum* isolated from soils.

*O. virens* was collected in Central Eastern of Portugal during the flowering period. Plant material was identified by a plant taxonomist and voucher specimens were deposited in the Herbarium of ESACB (Escola Superior Agrária de Castelo Branco). Aerial parts of the plant were separated into two lots, one of them was immediately submitted to hydrodistillation and other one was dried at room temperature during a month.

Yellowish essential oils were obtained from *O. virens* in a yields of 1.8% and 2.4% (v/W) to fresh and dry plant, respectively. The essential oil presented as major components linalool and thymol.

Our results showed that although the essential oil yield of dry plant is greater than those of fresh plant.

In order to the antifungal activity, both essential oils of *O. virens* studied by us, on doses of 10 µL, inhibited totally the growth of the tested fungi. Doses of 5 µL of each essential oil also showed activity against the fungi strains used in this work, in particular against *Penicillium* sp and *Aspergillus niger*.

**P 46. Identification of methyl 2-hydroxy-3-methylhexanoate - a new headspace constituent of *Galanthus nivalis* L. (Amaryllidaceae)**

Radulović NS<sup>1</sup>; Miltojević AB<sup>1</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science and Mathematics, University of Niš, Višegradska 33, 18000 Niš, Serbia

anamiltojevic@yahoo.com

Keywords: *Galanthus nivalis* L.; Amaryllidaceae; snowdrop; headspace; synthesis; methyl 2-hydroxy-3-methylhexanoate

*Galanthus nivalis* L. (Amaryllidaceae), snowdrop, is an early spring flower native to a large area of Europe, highly admired for its delicate beauty. It has attracted considerable attention due to its capability to biosynthesize pharmaceutically important compounds, especially alkaloids such as galanthamine [1]. Despite its popularity, the volatiles of this species have never been previously investigated.

Detailed GC and GC/MS analyses of the headspace constituents of intact *G. nivalis* flowers revealed, among others, the presence of a number of methyl esters of 2-hydroxycarboxylic acids known to be olfactively important trace components of some flowers absolutes, highly valued in perfumery [2]. In order to corroborate the tentative mass spectral identification of these headspace constituents, a synthesis of several methyl esters of different  $\alpha$ -hydroxycarboxylic acids was performed, utilising two different synthetic approaches. The stereospecific one involved the diazotisation and hydrolysis of L-amino acids to the corresponding L- $\alpha$ -hydroxycarboxylic acids, whereas the other approach included the addition of HCN to an aldehyde and the hydrolysis of the obtained cyanohydrin. In both cases the obtained  $\alpha$ -hydroxycarboxylic acids were esterified by ethereal diazomethane. One of the headspace constituents was shown to be the diastereoisomerically pure methyl 2-hydroxy-3-methylhexanoate (1). Compound 1, which can be regarded as a new plant secondary metabolite, was obtained in gram quantities starting from 2-methylvaleraldehyde (total yield 55%), in order to perform its full spectral characterisation (MS, IR, 1D and 2D – NMR, <sup>1</sup>H and <sup>13</sup>C) and for further testing of its biological activity and olfactory properties.

Acknowledgements: Authors are grateful to the Ministry of Education and Science of the Republic of Serbia (Project No. 172061) for financial support of this work.

1. B Sidjimova et al. (2003) Pharmazie 58: 935-936.
2. R Snowden et al. (2005) Flavour Fragr. J. 20: 372-380.

### P 47. Chemical composition and *in vitro* biological activity of the volatile and fixed oils of *Nigella sativa*

Piras A<sup>1</sup>; Rosa A<sup>2</sup>; Marongiu B<sup>1</sup>; Falconieri D<sup>3</sup>; Porcedda S<sup>1</sup>; Ozcelik B<sup>4</sup>; Koca U<sup>5</sup>

<sup>1</sup>Università di Cagliari, Dipartimento di Scienze Chimiche e Geologiche

<sup>2</sup>Università di Cagliari, Dipartimento di Scienze Biomediche

<sup>3</sup>ITI "M. Giua", Cagliari; <sup>4</sup>Gazi University, Dipartimento di Microbiology; <sup>5</sup>Gazi University, Dipartimento di Pharmacognosy

maronb@unica.it

Keywords: *Nigella sativa*; essential and fixed oil; antibacterial; antifungal; antituberculosis; supercritical extraction

*Nigella sativa* L. (Ranunculaceae) grows in Mediterranean and western Asian countries. *N. sativa* seeds have traditionally been used in folk medicine as a natural remedy for various diseases as well as a spice. The seeds of *N. sativa* have been subjected to a range of pharmacological, phytochemical and nutritional investigations. The seeds contain both fixed and essential oils, proteins, alkaloids and saponin. Much of the biological activity of the seeds has been shown to be due to thymoquinone, the major component of the essential oil. The fixed oil is composed mainly of unsaturated fatty acids, including the unusual C20:2 eicosadienoic acid. Isolation of volatile and fixed oils from *N. sativa* seed of Turkey and Egypt have been obtained by supercritical fractioned extraction with carbon dioxide. Extraction experiments were carried out at pressures of 90 and 250 bar and temperature of 40 °C. The extraction step performed at 90 bar produced a volatile fraction mainly formed by thymoquinone (79-86%) and o-cymene (5-11%). The oil yield relative to this step of the process was 0.1-0.3% by weight of the charge. The last extraction step at 250 bar produced a fixed oil. The yield of this step was 21-26% by weight. The most represented fatty acids of fixed oil from *N. sativa* were 18:2 n-6 (54-55%), 18:1 n-9 (22-23%), 16:0 (12-13%), 18:0 (3%), and 20:2 (2-3%). The fixed oils obtained from *N. sativa* were evaluated for the antibacterial activity by employing standard strains of *E. coli*, *P. aeruginosa*, *A. baumannii*, *S. aureus*, *E. faecalis* via broth microdilution method. *In vitro* antifungal activity of the derivatives against *C. albicans*, *C. tropicalis*, and *C. krusei* were screened by using ketoconazole, and fluconazole as control agents. Anti-mycobacterium activity the breakpoint concentration ( $\mu\text{g mL}^{-1}$ ) was determined against standard strains of *M. tuberculosis H37Rv* and *M. avium* (ATCC 15769) by using the colorimetric resazurin microtiter assay (REMA). The compounds displayed antimicrobial activity towards all of the standard (ATCC, RSKK) strains of the tested bacteria at MIC values of 8-64  $\mu\text{g mL}^{-1}$  and were revealed to be ineffective against isolated strains (MIC > 256  $\mu\text{g mL}^{-1}$ ). The fixed oils emerged as the most effective against the bacteria of *M. avium* with MIC values between 2 and 8  $\mu\text{g mL}^{-1}$ . On the other hand, the compounds exhibited better antifungal effect against *C. krusei* at 64  $\mu\text{g mL}^{-1}$ , which is close to the effect of the control fluconazole.



**P 48. Carvacrol attenuates cancer pain in mice**

Guimarães AG<sup>1</sup>; Melo NSR<sup>1</sup>; Oliveira MA<sup>1</sup>; Alves RS<sup>1</sup>; Bezerra DP<sup>1</sup>; Quintans JSS<sup>1</sup>; Quintans-Júnior LJ<sup>1</sup>

<sup>1</sup>Federal University of Sergipe - Brazil

jullyanas@yahoo.com.br

Keywords: Monoterpene, carvacrol, cancer, pain

The purpose of the present study was to evaluate the activity of carvacrol (CARV), a phenolic monoterpene, on hypernociception induced by Sarcoma 180 (S180) in mice.

Swiss male mice (20-30 g) were divided into groups (n = 12/grupo) which were treated (s.c.) after tumor induction with vehicle (saline + cremophor) CARV (12.5, 25 or 50 mg/kg) or morphine (15 mg/kg) daily for 15 days. For tumor induction a suspension of 10<sup>6</sup> cells (25 mL) was injected into the right hindpaw of the mice. Mechanical hypernociception was evaluated using digital Von Frey apparatus (Insight<sup>®</sup>, Brazil), and the paw withdrawal threshold was determined in grams and expressed as delta ( $\Delta$  = force (g) after cells implantation - the force (g) before cells implantation). The spontaneous nociception and limb use was assessed guarding and flinching behaviors were measured during a 10 min observation period. Tumor growth assessed by paw volume was measured using a plethysmometer. All of these parameters were evaluated at days 1, 3, 5, 7, 9, 13 and 15 after tumor induction. The data were statistically analyzed by ANOVA followed by Tukey test. Protocols were approved for the UFS Ethic Committee (CEPA: 71/10).

In mechanical hypernociception induced by S180, CARV (50 mg/kg) was able to increase significantly ( $p < 0.05$ ) the paw withdrawal threshold in 9 ( $0.37 \pm 0.50$  g), 11 ( $0.85 \pm 0.65$  g), 13 ( $1.48 \pm 0.65$  g) and 15 ( $1.63 \pm 0.62$  g) days compared to vehicle ( $2.50 \pm 0.34$  g,  $3.87 \pm 0.46$  g,  $3.82 \pm 0.69$  g,  $4.24 \pm 0.60$  g, respectively). This same dose was able to reduce ( $p < 0.05$ ) the spontaneous nociception on days 11 ( $0.17 \pm 0.11$  s), 13 ( $0.17 \pm 0.17$  s) and 15 ( $0.25 \pm 0.18$  s) days, different from vehicle ( $1.08 \pm 0.43$  s,  $1.17 \pm 0.37$  s,  $1.08 \pm 0.29$  s, respectively). Besides, all doses improved the limb use in 9 (12.5 mg/kg:  $3.91 \pm 0.08$ , 25 mg/kg:  $4.00 \pm 0.00$ ; 50 mg/kg:  $3.92 \pm 0.08$ ) and 11 (12.5 mg/kg:  $3.75 \pm 0.13$ , 25 mg/kg:  $3.75 \pm 0.13$ , 50 mg/kg:  $3.83 \pm 0.11$ ) days, when compared to vehicle (9 days:  $3.58 \pm 0.15$ , day 11:  $3.17 \pm 0.24$ ). In 13 ( $3.58 \pm 0.19$ ) and 15 days ( $3.50 \pm 0.19$ ) only the highest dose promoted significant changes ( $p < 0.05$ ) compared it with the control (day 13:  $2.91 \pm 0.23$ , day 15:  $2.83 \pm 0.24$ ). However, no significant change of paw volume was observed.

The results show that CARV have antinociceptive activity on nociception induced sarcoma 180, suggesting that this compound might be a promising candidate for the treatment of cancer pain.

Acknowledgements: CNPq and FAPITEC-SE

**P 49. Comparison of essential oil composition from *Artemisia* species: *A. dracunculus*, *A. absinthium*, *A. abrotanum*, *A. vulgaris***

Sponza S<sup>1</sup>; Chizzola R<sup>1</sup>

<sup>1</sup>University of Veterinary Medicine

*silvia.sponza@vetmeduni.ac.at*

Keywords: *Artemisia dracunculus*; *Artemisia absinthium*; *Artemisia abrotanum*; *Artemisia vulgaris*; essential oil compounds; GC

*Artemisia* is a large and diverse genus of plants belonging to the family *Asteraceae* [1]. *Artemisia* species are used as spices in the culinary tradition as well as herbal medicine, in particular its essential oil, in folk medicine [2, 3].

The objective of the study is to compare the essential oil composition of four *Artemisia* species wild grown in the Botanical Garden of the Institute for Applied Botany and Pharmacognosy (Vienna, Austria).

A representative sample of leaves of *A. dracunculus*, *A. absinthium*, *A. abrotanum*, and *A. vulgaris* were collected, air dried and distilled in the Eppendorf MicroDistiller. The analyses of the essential oils were performed by GC-MS and GC-FID. The compounds were identified according to their mass spectra and retention indices [4].

*A. dracunculus* and *A. absinthium* present a simple essential oil composition where the main compound is estragole with 79,5% and  $\beta$ -thujone with 78,1%, respectively. The main compounds of the essential oil of *A. abrotanum* are davone derivates, while the *A. vulgaris* essential oil composition shows cineol (36,0%), sabinene (17,4%) and terpinene-4-ol (11,0%) as main compounds.

The study highlights that each four *Artemisia* species investigated present different main compounds enabling their differentiation.

Acknowledgments: This project is carried out within the PlantLIBRA project ([www.plantlibra.eu](http://www.plantlibra.eu)) that has received funding from the European Community's FP7/2007-2013 under grant agreement n°245199.

1. D Obolskiy et al. (2011) J. Agric. Food Chem. 59: 11367-11384.
2. D Lopez-Lutz et al. (2008) Phytochemistry 69: 1732-1738.
3. S Kordali et al. (2005) J. Agric. Food Chem. 53: 9452-9458.
4. R P Adams (2007) Identification of Essential Oil Components by Gas chromatography / Mass Spectrometry, 4<sup>th</sup> ed. Allured Pub. Corp., Carol Stream, IL.

## P 50. Comparative evaluation of fruit volatiles from two pear (*Pyrus communis* L.) Portuguese varieties

Lucas SC<sup>1</sup>; Grego JAB<sup>1</sup>; Trindade H<sup>2</sup>; Figueiredo AC<sup>2</sup>

<sup>1</sup>Escola Superior Agrária de Santarém, Departamento de Ciências Agrárias e Ambiente, Quinta do Galinheiro, S. Pedro, 2000 904 Santarém, Portugal

<sup>2</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portugal

Susanalucas1@gmail.com

Keywords: *Pyrus communis*, Rosaceae, fruit, pear, Rocha pear, volatiles

With about 10969 ha of pears and an average annual production of about 176870 t, the pear (*Pyrus communis* L.) is the second most cultivated fruit crop in Portugal [1]. FAO statistics reveal that Portugal ranked 11 among the Top 20 pear-fruit exporters in 2009 [2]. Many local pear varieties are produced throughout Portugal, but one variety, Rocha pear (*pêra Rocha*), makes up about 95% of total production and is distinguished with a Protected Designation of Origin, or PDO [3]. As part of a wider research aiming to characterize a pear cultivar resulting from a chance seedling originated on a farm located at Tábua, Portugal, this work compares these pear fruit volatiles with those from Rocha pear.

The fruits from the pear under study (Tp) were collected and kept at 6°C in normal atmosphere for 4 months. Rocha pear fruits (Rp) were obtained commercially. The volatiles were isolated by hydrodistillation and analysed by Gas Chromatography and Gas Chromatography-Mass Spectrometry [4].

The same yield, <0.05% (v/f.w.) was obtained for both fruit volatiles, hexyl acetate (51% in Tp, 37% in Rp), butyl acetate (12% in Tp, 18% in Rp) and *trans,trans*- $\alpha$ -farnesene (4% in Tp, 15% in Rp) being the dominant components. Although Tp and Rp fruit volatiles shared similar main compounds, some major differences were detected in the minor compounds (<5%). The presence of hexyl acetate in both Portuguese fruits varieties, as well as in other pears varieties volatiles, such as William's [5], does not allow using this compound as a diagnostic character to differentiate among varieties. Further studies are being carried out to evaluate the differences in volatile profile at harvest date and after storage, as volatiles, in addition to appearance, flesh texture and grit stone content, are determinant in pear quality.

Acknowledgments: Eng. Justina Franco (ESAC), Eng. Rui Sousa (INRB), Eng. Arminda Lopes (ESAV), Sr Fernando Tavares e Dr Mário Vicente (Picos de Couto). Partially supported by Picos de Couto, Portugal.

1. INE (2011) Estatísticas Agrícolas 2010. Instituto Nacional de Estatística (INE, I.P.), Lisboa, Portugal.
2. FAOSTAT (2009) Food and Agriculture Organization of the United Nations - Key statistics of food and agriculture external trade, FAOSTAT, Statistics Division.
3. J Salta *et al.* (2010) *J Funct Foods* 2: 153-157.
4. P Barbosa *et al.* (2010) *Journal of Nematology* 42: 8-16.
5. F Rapparini, S Predieri (2003) In: *Horticultural Reviews*, Vol. 28. J Janick (Ed.). John Wiley & Sons, pp: 237-324.

**P 51. Composition of volatile compounds of horseradish roots (*Armoracia rusticana* L.) depending on genotype**

Tomsone L<sup>1</sup>; Kruma Z<sup>1</sup>; Talou T<sup>2</sup>; Galoburda R<sup>1</sup>

<sup>1</sup>Latvia University of Agriculture, Latvia; <sup>2</sup>Institut Polytechnique de Toulouse, France

[lolita.tomsone@llu.lv](mailto:lolita.tomsone@llu.lv)

Keywords: SPME; horseradish; allyl isothiocyanate; genotype

Horseradish (*Armoracia rusticana* L.) is a member of the family *Brassicaceae*. It is a perennial plant, with a particularly pungent flavour and significant antioxidant properties. Horseradish has about 0.2% to 1.0% essential oil, mainly sinigrin, sinigrin-derived allyl isothiocyanate, diallyl sulfide. The present study was carried out to determine composition of volatile compounds of *A. rusticana* depending on genotype. Nine genotypes of horseradish were collected at the end of August 2011 at Pure Horticultural Research Centre (Latvia) collection field. Volatiles from fresh horseradish roots were extracted using solid phase microextraction (SPME) with DVB/Car/PDMS fibre and analysed by gas chromatography-mass spectrometry. Compounds were identified by comparison of their mass spectra with mass spectral libraries (Nist98), and by calculation of linear retention indexes. The studied horseradish genotypes differed both in quantitative and qualitative content of aroma compounds and the highest content of horseradish volatile compounds in genotype 'Turku' was detected. A total of 15 volatile compounds were identified. The main aroma compound of all horseradish samples was allyl isothiocyanate forming 61-82% of total identified volatile compounds and these results are comparable with those found in literature. Isothiocyanates are formed during breakdown of sulfur-containing compounds – glucosinolates that are in cruciferous vegetables. Also all horseradish samples contained significant amounts of phenylethyl isothiocyanate (4-18%) that is formed from glucosinolate - gluconasturtiin. This study revealed the great influence of genotype on the content of volatiles in horseradish roots.

Acknowledgement This research has been performed within the framework of the "Osmose" between Latvia University of Agriculture and Institut Polytechnique de Toulouse. Authors acknowledge Pure Horticultural Research Centre for supply of horseradish roots.

**P 52. Alkamides from *Achillea serbica* Nym. (Asteraceae) root essential oil**

Radulovic N<sup>1</sup>; Đorđević M<sup>1</sup>; Blagojevic P<sup>1</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science and Mathematics, University of Niš

nikoradulovic@yahoo.com

Keywords: *Achillea serbica* Nym; Asteraceae; yarrow; root; essential oil; alkamide

An extensive amount of work has been done on the aerial parts volatiles of many yarrow species (*Achillea* L., Asteraceae) [1]. Contrary to that, and despite the fact that root oil ducts exist in a number of *Achillea* taxa, the underground parts were generally neglected as a potential source of essential oils [1]. The volatile oils of the above-ground parts of *Achillea serbica* Nym. (syn. *A. ageratifolia* (Sibth. & Sm.) Boiss. subsp. *serbica* (Nyman) Heimerl and *A. ageratifolia* (Sibth. & Sm.) Boiss. subsp. *ageratifolia* var. *serbica* (Nyman) Hayek) were analyzed on two previous occasions [2,3], but no data on the corresponding root oil exist. For that reason and due to the fact that the roots of this yarrow are a valued folk herbal remedy used as a general stimulant, we have performed detailed chemical analyzes (gas chromatography and gas chromatography/mass spectrometry) of the hydrodistilled *A. serbica* root oil. Interestingly, the oil was characterized by the presence of five different (alk)amides (pyrrolides) containing the C<sub>12</sub> and olefinic C<sub>16</sub> acid moieties: 3,6-epoxydodecanoic (1, 3.4%), (*E,Z*)-hexadeca-2,7-dien-10-ynoic (2, 21.8%), (*Z*)-hexadeca-7-en-10-ynoic (3, 8.6%), (*E,E,E*)-hexadeca-2,6,8-trien-10-ynoic (4, 3.6%), and (*E,Z*)-hexadeca-2,7-dienoic (5, 3.0%), which made 40.4% of the total oil. These were detected for the first time in an essential oil sample. The different patterns of unsaturation of the acid residues suggest close biogenetic connections with oleic and crepenynic acids [4]. Compounds 1-5 were previously reported as constituents of *A. serbica* root extract. Worth mentioning is the fact that alkamides are generally considered as unstable and that there are only two previous studies reporting this type of compounds as essential oil constituents, and both in *Achillea* species (*A. ptarmica* and *A. distans*) [5].

Acknowledgments: This work was funded by the Ministry of Education and Science of Serbia (Project 172061). PB was supported by UNESCO-L'OREAL National Fellowships Programme for Young Women in Life Sciences.

1. PML Lourenco et al. (1999) *Phytochemistry*, 51: 637-642.
2. J-C Chalchat et al. (1999) *J. Essent. Oil Res.* 11: 306-310.
3. N Simic et al. (2000) *Flavour Fragr. J.* 15 : 141-143.
4. H Greger et al. (1987) *Phytochemistry*, 26: 2289-2291.
5. J Lazarevic et al. (2010) *Nat. Prod. Res.* 24: 718-731.

**P 53. Genetic and chemical variation in Mexican oregano essential oil**Martínez-Natarén DA<sup>1</sup>; Parra-Tabla V<sup>2</sup>; Ferrer-Ortega MM<sup>2</sup>; Calvo-Irabién LM<sup>1</sup><sup>1</sup>Centro de Investigación Científica de Yucatán. Calle 43 #130 Chuburna Mérida 97200, Yucatan, Mexico<sup>2</sup>Departamento de Ecología Tropical, Campus de Ciencias Biológicas y Agropecuarias, Universidad Autónoma de Yucatán. 15.5, Carretera Mérida-Xmatkuil, 4-116, Mérida, Yucatán 97315, México

lumali@cicy.mx

Keywords: AFLP; chemotype; genetic diversity; genetic differentiation; *Lippia graveolens*.

Mexican oregano is an economically important aromatic plant. In order to understand the importance of environmental and genetic factors in essential oil production, in this study we estimated the genetic diversity and structure for 14 wild populations of *Lippia graveolens* located in four different bioclimatic regions in southeastern Mexico using AFLP (amplified fragment length polymorphism) markers. Essential oil composition from each individual (total n = 95) was characterized using GC and GC-MS [1]. Based on the definition of chemotype by previous authors [2] we identified three different chemotypes: (a) thymol and (b) carvacrol, characterized by high concentration of these monoterpenes and (c) sesquiterpene with high concentrations of a and b-caryophyllene. Two primer combinations produced a total of 87 polymorphic bands. The overall genetic diversity of *L. graveolens* described as the percentage of polymorphic loci (PPL= 60.9%) and Nei's gene diversity ( $H_j = 0.17$ ) was moderate but was not associated to bioclimatic conditions. Genetic variation was analyzed at different levels. Regarding chemotypes, thymol had the highest genetic diversity (PPL = 82.8% and  $H_j = 0.22$ ). PCoA revealed that chemotypes exhibit certain level of genetic differentiation. Maximum parsimony dendrogram showed that some individuals from the same chemotype clustered together. Bayesian analyses revealed a low but significant differentiation among chemotypes ( $q_{II} = 0.008$ ). Regarding populations, gene diversity showed significant differences ( $F_{13,1204} = 22.8$ ,  $P < 0.001$ ), the populations 4, 5 and 8 (predominantly thymol individuals) showed the highest gene diversity ( $H_j = 0.31-0.25$ ), while population 3 (exclusively sesquiterpene chemotype) showed the lowest value ( $H_j = 0.058$ ). Cluster (PCoA and maximum parsimony) and Bayesian analyses ( $q_{II} = 0.027$ ) revealed a low level of genetic differentiation among populations.

Acknowledgments: This study was financially supported by CONACyT, doctoral scholarship to DMN and grant given to LCI (CB-2007, 82873). We thank Sara V. Good, University of Winnipeg, for laboratory facilities and Dulce Linares and Felix Martínez-Nuñez for technical assistance in laboratory work.

1. V Acosta-Arriola (2011). Ms Thesis. CICY, Mérida, México.

2. H Hendriks et al. (1990). J. Essent. Oil Res. 2: 155-162.

## P 54. A survey of the essential oil composition of rose-scented geranium grown in South Africa

de Figueiredo M<sup>1</sup>; Stevens CM<sup>2</sup>; Withwell MI<sup>1</sup>

<sup>1</sup>Department of Agriculture and Environmental Affairs, P/Bag X9059, Pietermaritzburg, 3200, KwaZulu-Natal, South Africa

<sup>2</sup>Department of Agriculture and Environmental Affairs, P/Bag X9059, Pietermaritzburg, 3200, KwaZulu-Natal, South Africa

Maria.Figueiredo@kzndae.gov.za

Keywords: *Pelargonium* sp; chemical composition; variability; climatic diversity

Rose-scented geranium (*Pelargonium* sp) essential oil was received from different provinces of South Africa from 2003 to 2011 and its chemical composition determined by gas chromatography (GC) and GC/MS. The variability of the major constituents of the essential oil between provinces and within regions of the KwaZulu-Natal province was determined. South African climate presents great diversity between and within provinces [1]. The seasonal effect on the oil composition from the KwaZulu-Natal regions was investigated. With the exception of 6,9-guaiadiene and citronellol which varied significantly at  $P \leq 0.05$  between some of the provinces, the major constituents of the oil analysed were not significantly different. Within the KwaZulu-Natal province, linalool and citronellyl formate were found to differ significantly. Interactions between season and KwaZulu-Natal regions were not significant. The mean values of the major constituents of South African oil, expressed as percentages of the total constituents were: isomenthone 4.78, linalool 3.03, 6,9-guaiadiene 7.17, citronellyl formate 15.26, geranyl formate 8.25, citronellol 24.08 and geraniol 14.81. The rose-scented geranium essential oil received from different provinces of South Africa was found to have a lower linalool and a higher citronellyl formate than oils from other parts of the world [2]. These characteristics can be used to identify rose-scented geranium oil of South African origin.

1. South Africa weather website. <http://www.sa-venues.co./no/weather.htm> [26 April 2012]

2. EA Weiss (1997). *In*: Essential oil crops. CAB International (ed), New York and UK, pp. 24-58

## P 55. Essential oils composition of *Croton* species with occurrence in the Brazilian Amazon

Turiel NA<sup>1</sup>; da Silva JKR<sup>2</sup>; Carreira LMM<sup>3</sup>; Lucas FCA<sup>4</sup>; Monteiro OS<sup>5</sup>; Carvalho NCC<sup>5</sup>; Andrade EHA<sup>1</sup>; Maia JGS<sup>1</sup>

<sup>1</sup>Programa de Pós-Graduação em Química, Universidade Federal do Pará

<sup>2</sup>Programa de Pós-Graduação em Biotecnologia, Universidade Federal do Pará; <sup>3</sup>Coordenação de Botânica, Museu Paraense Emílio Goeldi

<sup>4</sup>Departamento de Ciências Naturais, Universidade Estadual do Pará

<sup>5</sup>Programa de Pós-Graduação em Química, Universidade Federal do Maranhão

nathalie@ufpa.br

Keywords: *Croton*; essential oil composition; cluster analysis

*Croton* is a genus of Euphorbiaceae comprising about 1300 species widespread in Africa, Asia and South America. Many species are used in the traditional medicine of these continents, especially to treat cancer, diabetes, malaria, ulcers, among other diseases [1]. The essential oils of leaves and fine stems of *Croton campestris* A. St.-Hil. (0.5%), *C. chaetocalyx* Mull. Arg. (1.1%), *C. draconoides* Mull. Arg. (0.8%), *C. eriocladus* Mull. Arg. (0.6%) and *C. glandulosus* Mull. Arg. (0.7%) were analyzed by GC and GC-MS. The plants were collected in Eastern Amazon and their compositions are rich in terpene compounds. The main constituents identified in the oil of *C. campestris* were (*E*)-caryophyllene (23.0%),  $\gamma$ -elemene (13.9%), germacrene D (13.7%),  $\beta$ -elemene (7.1%) and  $\delta$ -elemene (6.0%); in the oil of *C. chaetocalyx* were bicyclogermacrene (13.9%),  $\delta$ -elemene (13.5%), germacrene D (9.3%), spathulenol (9.0%),  $\delta$ -cadinene (8.0%) and (*E*)-caryophyllene (7.1%); in the oil of *C. draconoides* were  $\beta$ -pinene (16.9%),  $\alpha$ -pinene (16.5%), curzerene (12.8%) and germacrene D (9.0%); in the oil of *C. eriocladus* were (*E*)-caryophyllene (24.1%), germacrene D (17.9%),  $\alpha$ -humulene (6.2%), bicyclogermacrene (5.2%) and  $\delta$ -elemene (5.0%); and in the oil of *C. glandulosus* were spathulenol (19.7%), bicyclogermacrene (9.6%), (*E*)-caryophyllene (8.9%) and  $\delta$ -elemene (8.8%). A hierarchical cluster analysis using the compositional profile of the major constituents (below 5%) has been carried out. Complete linkage and absolute correlation coefficient distance was selected as measure of similarity. The oils were divided into three groups: *C. campestris* and *C. eriocladus* (92% similarity), *C. chaetocalyx* and *C. glandulosus* (88% similarity), and *C. draconoides* maintained separately.

Acknowledgments: This work was supported by BIONORTE/CNPq and FAPESPA/PA.

1. Salatino et al. (2007) J. Braz. Chem. Soc. 18: 11-33.



## P 56. Antifungal activity of *Piper* essential oils from the Amazon against the *Fusarium* disease in pepper

da Silva JKR<sup>1</sup>; Ramos AR<sup>2</sup>; Luz SFM<sup>2</sup>; Meireles ENM<sup>1</sup>; Andrade EHA<sup>3</sup>; Maia JGS<sup>4</sup>

<sup>1</sup>Programa de Pós-Graduação em Biotecnologia, Universidade Federal do Pará, 66075-900, Belém, PA, Brazil.

<sup>2</sup>Faculdade de Ciências Exatas e Naturais, Universidade Federal do Pará, 68501-970, Marabá, PA, Brazil.

<sup>3</sup>Programa de Pós-Graduação em Química, Universidade Federal do Pará, 66075-900, Belém, PA, Brazil.

<sup>4</sup>Programa de Pós-Graduação em Química, Universidade Federal do Pará, 66075-900, Belém, PA, Brazil.

joycekellys@ufpa.br

Keywords: *Piper nigrum*, *P. callosum*, *P. divaricatum*; *P. krukoffii*; *Fusarium solani* f. sp. *piperis*; antifungal activity

The pepper (*Piper nigrum* L.), originary from southeastern Asia, is a famous spice used in the entire world to aromatize many traditional and regional dishes. This species is very susceptible to the disease known as fusariosis, with restricted occurrence to Brazil. The attack of phytopathogen *Fusarium solani* f. sp. *piperis* Alb. causes root rot and large losses for the plant culture. The essential oils obtained from *Piper callosum* Ruiz & Pav, *P. divaricatum* G. Mey. and *P. krukoffii* Yunck. Were analyzed by GC and GC-MS and submitted to antifungal activity [2] against *Fusarium solani* f. sp. *piperis*. The oil compositions of *P. callosum*, *P. divaricatum* and *P. krukoffii* were rich in phenylpropanoid compounds. The main constituent identified in the oil of *P. callosum* were safrol (78.0%), methyeugenol (8.7%) and 1,8-cineole (2.6%); in the oil of *P. divaricatum* were methyleugenol (84.0%) and eugenol (8.7%); and in the oil of *P. krukoffii* were myristicin (40.3%), apiole (25.4%) and  $\alpha$ -elemene (8.2%). The inhibition percent of growth of *Fusarium solani* f. sp. *Piperis* in the oil of *P. callosum* was 53.0%. The inhibition percent varied from 78.9 to 100% in the oil of *P. divaricatum*. The oil of *P. krukoffii* was only able to inhibit the fungus growth in 24.5%. The inhibition percent data were submitted to analysis of variance (ANOVA,  $p < 0.05$ ). The differences were found to be extremely significant to *P. divaricatum* ( $P = 0.000$ ), moderate to *P. callosum* ( $P = 0.0023$ ) and not significant to *P. krukoffii* ( $P = 0.733$ ).

Acknowledgments: This work was funded by BIONORTE/CNPq and FAPESPA/PA and supported by UFPA/New Doctors Program.

1. SY Wang (2005) *Bioresource Technology* 96: 813–818.

## P 57. Essential oils composition of Annonaceae species with occurrence in the Brazilian Amazon

Maia JGS<sup>1</sup>; Andrade EHA<sup>1</sup>; Turiel NA<sup>1</sup>; Ribeiro AF<sup>2</sup>; Carreira LMM<sup>3</sup>; Lucas FCA<sup>4</sup>

<sup>1</sup>Programa de Pós-Graduação em Química, Universidade Federal do Pará

<sup>2</sup>Faculdade de Ciências Exatas e Naturais, Universidade Federal do Pará, Marabá, PA

<sup>3</sup>Coordenação de Botânica, Museu Paraense Emílio Goeldi

<sup>4</sup>Departamento de Ciências Naturais, Universidade Estadual do Pará

nathalie@ufpa.br

Keywords: essential oil composition; Annonaceae

The Annonaceae comprises tropical and sub-tropical species that are widespread in South and Central America, Africa, Asia and Australia. Early phytochemical review has estimated ca 120 genera and more than 2000 species in this family [1]. Economically, it is of appreciable importance as a source of edible fruits, raw material for cosmetics and perfumery and medicinal plants. As part an ongoing survey of the Amazon aromatic plants we are reporting the volatile composition of some Annonaceae species collected in Eastern Amazon, municipality of São Geraldo do Araguaia, Pará state, Brazil. The oils were analyzed by GC and GC-MS. The main constituents identified in the oil of *Xylopiá aromática* (Mart.) Lam. were spathulenol (21.7%), limonene (17.8%),  $\delta$ -elemene (10.2%) and bicyclogermacrene (9.4%); in the oil of *Annona paludosa* Aubl. were spathulenol (18.0%), caryophyllene oxide (14.7%), (*E*)-caryophyllene (7.6%) and germacrene D (5.8%); in the oil of *Annona crassiflora* Mart. were  $\gamma$ -elemene (21.0%),  $\delta$ -elemene (14.3%), (*E*)-caryophyllene (14.0%), germacrene D (13.4%) and bicyclogermacrene (9.6%); in the oil of *Guattteria schomburgkiana* Mart. were  $\delta$ -cadinene (14.7%), spathulenol (11.2%),  $\alpha$ -cadinol (10.9%), caryophyllene oxide (4.9%) and zonarene (4.0%); in the oil of *Duguetia echinophora* R.E.Fr were spathulenol (37.4%), caryophyllene oxide (7.1%), (*E*)-caryophyllene (5.8%) and germacrene D (5.7%). Eearly papers indicated that spathulenol and caryophyllene oxide could be considered as chemotaxonomic markers for Annonaceae species occurring in the Amazon [2,3]. Excepting the oil of *A. crassiflora*, the other oils showed significant amounts of spathulenol and caryophyllene oxide, which confirm this hypothesis.

Acknowledgments: This work was supported by BIONORTE/CNPq and FAPESPA/PA.

1. Lebouef et al. (1982) Phytochemistry 21: 2783-2813.

2. Andrade et al. (2003). Flavour Fragr. J .

3. Maia et al.(2003) Flavour Fragr. J .

**P 58. Antioxidant and antifungal activity and composition of the oils from *Lippia gracilis* and *L. origanoides***

Franco CS<sup>1</sup>; da Silva JKR<sup>2</sup>; Monteiro OS<sup>3</sup>; Carvalho NCC<sup>3</sup>; Ribeiro AF<sup>4</sup>; EHA<sup>4</sup>; Maia JGS<sup>4</sup>

<sup>1</sup>Programa de Pós-graduação em Ciências Farmacêuticas, Universidade Federal do Pará, 66075-900, Belém, PA, Brazil.

<sup>2</sup>Programa de Pós-Graduação em Biotecnologia, Universidade Federal do Pará, 66075-900, Belém, PA, Brazil.

<sup>3</sup>Programa de Pós-Graduação em Química, Universidade Federal do Maranhão, 65080-040 São Luís, MA, Brazil

<sup>4</sup>Programa de Pós-Graduação em Química, Universidade Federal do Pará, 66075-900, Belém, PA.

joycekellys@ufpa.br

Keywords: *Lippia gracilis*; *L. origanoides*; antioxidant and antifungal activity; oil composition

The present study was conducted to evaluate the antioxidant and antifungal properties of the essential oil of *Lippia gracilis* Schauer and *L. origanoides* Kunth, collected in the locality of São Felix das Balsas (February 2011), Maranhão state, and in the National Forest of Carajás (May 2011), Pará state, respectively. The chemical composition of hydrodistilled oils was analyzed by GC and GC-MS. The main constituents in the oil of *L. gracilis* were thymol (72.5%), *p*-cymene (9.3%) and thymol methyl ether (5.4%) and in the oil of *L. origanoides* were 1,8-cineol (31.9%), nerolidol (16.2%) and  $\alpha$ -terpineol (7.5%). The antioxidant potential of the samples was evaluated using the method of inhibition of free radical 2,2-diphenyl-1-picrylhydrazyl (DPPH). The essential oils were able to reduce the stable radical DPPH showing a significant antioxidant activity to *L. gracilis* ( $IC_{50} = 35.7 \pm 3.3 \mu\text{g.mL}^{-1}$ ), while for *L. origanoides* it was very low ( $IC_{50} > 500 \mu\text{g.mL}^{-1}$ ). The antifungal assay showed that both essentials oils inhibited the growth of the fungus *Cladosporium sphaerospermum*, with detection limit of 25  $\mu\text{g}$  to *L. gracilis* and 50  $\mu\text{g}$  to *L. origanoides*. The results showed that, at least, the essential oil of *L. gracilis* could be used as a natural preservative ingredient in food and pharmaceutical industries.

Acknowledgments: This work was supported by BIONORTE/CNPq and FAPESPA/PA

**P 59. Composition and antifungal activity of *Sphagneticola trilobata* Pruski (Asteraceae) volatile oils collected in different biomes**

Raggi L<sup>1</sup>; Cordeiro I<sup>2</sup>; Moreno PRH<sup>3</sup>; Young MCM<sup>1</sup>

<sup>1</sup>Núcleo de Pesquisa em Fisiologia e Bioquímica, Instituto de Botânica, São Paulo, Brazil

<sup>2</sup>Núcleo de Pesquisa em Curadoria do Herbário, Instituto de Botânica, São Paulo, Brazil

<sup>3</sup>Instituto de Química, Universidade de São Paulo, São Paulo, Brazil

*ludraggi@gmail.com*

Keywords: Wedelia; essential oil; Atlantic Rain Forest; Brazilian Cerrado; *Cladosporium sphaerospermum*

*Sphagneticola trilobata* (Asteraceae) is a native Brazilian herbaceous plant, commonly known as vedélia, used in folk medicine to treat a variety of illnesses, such as headaches, fevers, infections and respiratory pathologies. The geographical origin, soil characteristics and climate differences may influence in quantitatively and/or qualitatively the essential oils chemical composition. The present work analysed the composition and antifungal activity of the volatile oils from *S. trilobata* collected at two different biomes from São Paulo state: two populations from Atlantic Rain Forest (São Paulo [SP] and Paranapiacaba [PARN]) and one from the Brazilian Cerrado (Mogi-Guaçu [MGÇ]). Fresh aerial parts were hydrodistilled in a Clevenger apparatus during four hours and analysed by GC and GCMS. The antifungal activity was determined by the bioautography method as described by Homans & Fuchs [1] with the fungus *Cladosporium sphaerospermum* (Penzig). The yields (w/w) were 0.21, 0.11 and 0.09% for SP, PARN and MGÇ respectively. The chemical composition of the volatile oils obtained from these three locations was quantitatively distinct. The main components of the oils were: SP -  $\beta$ -Pinene 45.2%,  $\alpha$ -Phellandrene 16.0%,  $\alpha$ -Pinene 15.7% and Limonene 8.8%; PARN -  $\alpha$ -Pinene 29.8%,  $\beta$ -Pinene 22.0% and  $\alpha$ -Phellandrene 14.3% and MGÇ -  $\alpha$ -Pinene 28.7%,  $\alpha$ -Phellandrene 18.4% and Limonene 15.5%. The oil from the population of the Brazilian Cerrado showed strong inhibitory activity against the fungus tested. The oil from the Atlantic Rain Forest at PARN showed lower activity and SP did not show growth inhibition. The highest percentage of limonene in the composition of the volatile oil from MGÇ plants might have had influence in the activity found, either for intrinsic activity or synergism with other compounds in the oil.

Acknowledgments: Pos-graduate program in Biodiversidade Vegetal e Meio Ambiente and CNPq

1. L Homans, A Fuchs (1970) J. Chromatogr., 51: 327-329.

**P 60. Antioxidant activity of essential oils from *Satureja* growing in Spain**

García-Rellán D<sup>1</sup>; Verdeguer M<sup>1</sup>; Ortuño C<sup>2</sup>; Blázquez MA<sup>3</sup>; Boira H<sup>1</sup>

<sup>1</sup>Instituto Agroforestal Mediterráneo (Universidad Politécnica de Valencia)

<sup>2</sup>Grupo de Análisis y Simulación de Procesos Agroalimentarios, Dpto. Tecnología de Alimentos (Universidad Politécnica de Valencia)

<sup>3</sup>Farmacología, Facultat de Farmàcia (Universitat de València)

dagarrel@upvnet.upv.es

Keywords: *Satureja*, essential oil, antioxidant activity, total phenolic compounds

The phenolic content and the antioxidant activity of five essential oils obtained from *S. montana*, *S. innota*, *S. cuneifolia* and *S. intricate* growing in Spain were measured in order to find new potential sources of natural antioxidant.

*Satureja* is a genus of the perennial or annual semi-bushy species native to warm temperate regions. The aerial parts of five wild perennial populations of *Satureja* species were collected from the phytogeographic area of this genus in the Comunidad Valenciana (Spain) in September 2010. One population of each species was studied, except in the case of *S. innota*, for which two populations were analysed due to their different ecological conditions.

Essential oils were obtained by hydro-distillation using a Clevenger-type apparatus. Total phenolic compounds and antioxidant activity of the essential oils were determined according to Folin-Ciocalteu [2] and to FRAP assays [3, 4], respectively.

The total phenolic content of the essential oils varied between *Satureja* species. Significant differences were found between *S. montana* and others *Satureja* species, being its content in total phenolic compounds much higher. Data also provided no significant differences between both populations of *S. innota*.

The results showed a relationship between total phenolic compounds and antioxidant activity.

*S. montana* had the highest antioxidant activity. The GC/MS analysis [5] indicates that this essential oil was characterized by a higher carvacrol content compared to the others *Satureja* essential oils.

1. R Morales et al. (2010) Flora Ibérica 12: 414-421.

2. VL Singleton et al. (1999) Meth. Enzymol. 299: 152-78.

3. IFF Benzie, JJ Strain (1996) Anal. Biochem. 239: 70-76.

4. R Pulido et al. (2000) J. Agric. Food Chem. 48: 3396-3402.

5. D García-Rellán (2009). Tesis de Máster UPV.

## P 61. Individual variability of wormwood (*Artemisia absinthium* L.) essential oil composition

Zámboriné-Németh E; Bernáth J; Kindlovits S; Sárosi S

*Corvinus University of Budapest*

*eva.nemeth@uni-corvinus.hu*

Keywords: chemotype, diversity, thujone, myrcene, sabinene

Wormwood has gained a contradictory reputation by its thujone content and potential toxicity which is heavily disputed until today (1). In the last decades, composition of the essential oil of several populations of different origin has been studied and numerous chemotypes identified (e.g. 2,3,4). However, these results always represented the quality of bulk samples which is not able to reveal the natural variability of the species. In the recent investigations we studied the essential oil content and composition of 30 individual plants in a common cultivated genotype in Hungary. Shoot samples were taken at the beginning of flowering (July). The dried herb was hydrodistilled (PhEur) and the oil analysed by GC-MS. The components were identified by MS libraries and their LRIs.

The essential oil content varied between 0.110 and 0.962% d.w. We identified 49 compounds in the oil.  $\beta$ -Thujone was the main component (in 55-71% of the oil) in 53% of the plants. However, in 9 plants it has been found only in traces. The linear correlation coefficient between essential oil content and its  $\beta$ -thujone proportion was  $r = 0.5764$ .

The second and third most abundant components were  $\beta$ -myrcene (4-34% in oil) and sabinene (2-32% in oil) both being main compounds in 13% of the samples. Besides, in 20% of the oils, they were found in appr. equal proportions. A single sample contained trans-chrysanthenol and another one an unidentified sesquiterpene (RT: 44.63; LRI 1985) as main component. Sabinene and trans-chrysanthenol chemotypes of wormwood have not been reported before. On the other side, compounds described as main components in the populations of other regions like p-cymene, bornyl-acetate, cis-epoxyocimene, cis-chrysanthenyl-acetate (2,3,4) have not been found in our samples. Minor compounds present in each sample were p-cymol,  $\beta$ -caryophyllene, linalyl-butanoate,  $\alpha$ -curcumene, neryl-isobutanoate, lavandulyl-isovalerate, neryl-isovalerate, caryophyllene-oxide, geranyl-isovalerate. Each oil consisted higher number of sesquiterpene compounds than monoterpenes.

The results demonstrate the significance of individual sampling for chemotaxonomical surveys. Selection of unique chemotypes may contribute to development of new products.

1. DW Lachenmeyer, T Kuballa (2007) J. Sci. Food and Agricult. 87: 2147-2151.
2. A Arino et al (1999) J. Essent. Oil Res. 11: 619-622.
3. JA Pino et al. (1997) J. Essent.Oil res. 9: 87-89.
4. A Basta et al. (2007) J. Essent. Oil Res. 19: 316-318.

**P 62. Chemical composition, antioxidant and antibacterial activity of the essential oils of *Pimpinella pruatjan*, an endemic species from Java Island, Indonesia**

Nurcahyanti ADR; Al-ani I; Sporer F; Wink M

*Institute of Pharmacy and Molecular Biotechnology, Heidelberg University*

*agustina.nurcahyanti@gmx.de*

Keywords: antioxidant activity, antibacterial activity, *Pimpinella pruatjan*, essential oil, chemical composition

The essential oil from root and aerial parts of native *Pimpinella* species growing in a restricted mountain area in Java was studied concerning chemical composition, antioxidant and antibacterial activities. Using high resolution GLC-MS, the essential oil isolated by hydrodistillation was separated and components were identified [1]. The antioxidant activity was determined by conjugated diene assay and analysis of scavenging free radicals activity [2,3]. The antibacterial activity of the essential oil was assessed against some pathogenic bacteria [4]. The essential oil contains monoterpenes, sesquiterpenes, and phenylpropanoids. Antioxidant and antibacterial properties (MIC and MBC) against Gram negative and Gram positive bacteria will be presented. The composition of the essential oils from *Pimpinella* species growing in tropical climate region will be compared with known *Pimpinella* species which are widely distributed in the temperate zone of Europe and Iran.

Acknowledgments: We are grateful to Dr. Ireng Darwati Kadarso (Indonesian Medicinal and Aromatic Crops Research Institute, Indonesia) for providing *Pimpinella pruatjan*.

1. ML Ashour et al. (2009) J. Pharm. Pharmacol. 61: 1079-87.
2. MB Farhat et al. (2009) J. Agric. Food Chem. 57: 10349-56.
3. A Wei, T Shibamoto (2010) J. Agric. Food Chem. 58: p. 7218 - 7225.
4. NCCLS (2006) *In: Approved Standards*. 9th ed. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically. Wayne, Pennsylvania.

**P 63. The composition of the essential oil from “candeia da serra” (*Eremanthus erythropapus*) determined by injection of the oil and by SPME of the entrapped vapor, as well as its antibacterial activity**

Lima FWJ<sup>1</sup>; Dantas-Barros AM<sup>2</sup>; Braga TV<sup>3</sup>; Siqueira EP<sup>4</sup>; Nelson DL<sup>5</sup>

<sup>1</sup>Faculdade de Farmácia, Universidade Federal de Minas Gerais

<sup>2</sup>Depto. de Produtos Farmacêuticos, Faculdade de Farmácia, Universidade Federal de Minas Gerais

<sup>3</sup>Faculdade Pitagoras

<sup>4</sup>Laboratório de Química de Produtos Naturais, Centro de Pesquisas René Rachou

<sup>5</sup>Depto. de Alimentos, Faculdade de Farmácia, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

[dlnelson@ufmg.br](mailto:dlnelson@ufmg.br)

Keywords: candeia; composition; essential oil; antibacterial activity

The essential oil from candeia da serra (*Eremanthus erythropapus*) was isolated by hydrodistillation. The composition of the oil was determined by injection of the oil onto a 30 m X 0.25 mm DB-5 column in a Shimadzu model GC-17 gas chromatograph coupled to Shimadzu QP5050 mass selective detector. The composition of the headspace above the essential oil was also determined by adsorption onto a DVB/PDMS-coated fiber and desorption in the injection chamber of the chromatograph. The potential of the essential oil from the leaves of candeia to inhibit the growth of *Escherichia coli* (ATCC 25922) and *Staphylococcus aureus* (ATCC 25923) in foods was also assessed. The composition of the essential oil was confirmed by the Kovats index and the NIST library. A total of 43 compounds were isolated, mostly terpenes. The study of the antimicrobial activity was performed employing the disk diffusion method standardized by the Subcommittee for Antimicrobial Susceptibility Tests (NCCLS) [1]. Isolated colonies of bacteria were standardized in sterile saline solution according to the 0.5 McFarland scale. After immersing the swab in this solution, the microorganism was inoculated on Mueller-Hinton agar. Filter paper discs (6 mm), previously coated with 10  $\mu$ L of sample at 50 mg/mL were placed on the surface of the agar. A disk impregnated with ethanol was used as a negative control, and a disk impregnated with 10 mg of Ampicillin was used as a positive control. Tests were performed in six replicates. After incubation of the plate in an oven at 35 °C for 18 hours, the formation of inhibition zones around the disks was assessed. An inhibition zone  $\geq 7$  mm was considered to be positive for herbal extracts [2]. The *Escherichia coli* (ATCC 25922) and *Staphylococcus aureus* (ATCC 25923) showed resistance in the in vitro test, indicating that the essential oil from candeia leaves does not inhibit the growth of these microorganisms.

Acknowledgements: The Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for financial support and scholarships.

1- NCCLS. (2003) Padronização dos Testes de Sensibilidade a Antimicrobianos por Disco-difusão: Norma Aprovada – Oitava Edição. Substitui a Norma M2-A7. 23, n. 1.

2- Nascimento, G. G. F. *et al.* (2000) Braz. J. of Microbiol. 31, 247-256.



**P 64. Seasonal variations in essential oil composition of roots and plants of *Artemisia absinthium* L in a population from Teruel (Spain)**

Llorens-Molina JA<sup>1</sup>; Vacas S<sup>2</sup>; García-Rellán D<sup>1</sup>; Boira H<sup>1</sup>

<sup>1</sup>Instituto Agroforestal Mediterráneo, Universidad Politécnica de Valencia

<sup>2</sup>Centro de Ecología Química Agrícola – Instituto Agroforestal del Mediterráneo, Universidad Politécnica de Valencia

juallom2@qim.upv.es

Keywords: *Artemisia absinthium*; essential oil; seasonal variations; root; chemotype

The essential oil of *Artemisia absinthium* L has been extensively studied because of its many applications as to prepare absinth beverages as for the ethnopharmacological point of view [1, 2]. Seven chemotypes have been characterized in the Iberian Peninsula, three of them lacking in the neurotoxic compound beta-thujone [1]. Nevertheless, less attention has been paid to root essential oil composition [3, 4, 5]. The aim of this research is to study the plant and root essential oil composition of a (Z)-epoxyocimene-chrysantenyl acetate chemotype of an *Artemisia absinthium* L population from Teruel (Spain) over its vegetative period (may-august). For this purpose, three samplings were performed. In each one of them, five plants were collected and the essential oils from the fresh aerial parts and roots were obtained by means of a Clevenger type and a Likens-Nickerson apparatus, respectively. They were analyzed by GC and GC-MS with the same column (DB-5) and operating conditions.

Concerning the aerial parts, the results confirm their chemotype and show a predominance of individual variations with regard to seasonal ones. The sesquiterpenoid fraction is noticeable (4-10 %) and a significant occurrence of C12 compounds, such as (8E,10E)-dodeca-8,10-dien-1-ol, was found at the beginning of vegetative period. Regarding the root extracts composition, their main components are monoterpenes (35-77 %), mainly, alpha-fenchene and beta-myrcene, and monoterpene esters (15-55 %), mainly bornyl acetate, while the sesquiterpenoid fraction and other compounds amount is almost negligible. These results are similar to the ones obtained in a previous research [5] whose aerial part chemotype was different (with a high amount of beta-thujone). The conclusions of these results lead to define two trends to pursue this research: firstly, to evaluate separately individual and seasonal variability, monitoring the aerial parts composition in individual plants over the vegetative cycle. Secondly, to go deeply into the secondary metabolites role of plants and roots to explain their qualitative differences.

1. A Ariño (1999) Variabilidad química en los aceites esenciales de *Artemisia absinthium* de la Península Ibérica, tesis doctoral, Universidad del País Vasco (Spain).
2. A Orav et al (2006) Proc. Estonian Acad. Sci. Chem., 55(3): 155-165.
3. Al Kennedy et al. (1993) Phytochemistry, 32(6): 1449
4. S Nin et al. (1997) Plant Cells Reports, 16: 725-730
5. P Blagojevic et al. (2006) J. Agric. Food Chem. 54: 4780-4789

**P 65. Diurnal and environmental variations in essential oil composition of *Mentha suaveolens* L and *Mentha longifolia* L populations from the Eastern Iberian Peninsula**

Llorens-Molina JA<sup>1</sup>; Vacas S<sup>2</sup>; López S<sup>3</sup>; Rivero CF<sup>3</sup>; García-Rellán D<sup>1</sup>; Boira H<sup>1</sup>

<sup>1</sup>Instituto Agroforestal Mediterráneo, Universidad Politécnica de Valencia

<sup>2</sup>Centro de Ecología Química Agrícola–Instituto Agroforestal del Mediterráneo, Universidad Politécnica de Valencia

<sup>3</sup>Escuela Técnica Superior de Ingeniería Agronómica y del Medio Natural, Universidad Politécnica de Valencia

juallom2@qim.upv.es

Keywords: *Mentha longifolia*, *Mentha suaveolens*; essential oil; chemotypes; diurnal variations; environmental factors

Many researches dealing with the geographical chemical variability [1, 2] and biological applications [3, 4] of *Mentha longifolia* L and *Mentha suaveolens* L have been developed. Nevertheless, a limited attention has been paid to environmental factors affecting their essential oil composition [5]. The goal of this research is to contribute to knowledge of diurnal and organ variations and soil characteristics in essential oil of *Mentha suaveolens* L and *Mentha longifolia* L from six populations in the Eastern Iberian Peninsula.

For this purpose, a total amount of 79 samples were processed and the following variables were studied: location and climate (Coastal or Continental Mediterranean), morning or evening harvest. organ (leaves or flowers) and soil characteristics. The plant material was frozen and kept at -40°C until its extraction by means of a Likens-Nickerson apparatus. The extracts were analyzed by GC and GC-MS with the same column (DB-5) and operating conditions. In addition, 24 soil samples were obtained and pH, conductivity, organic matter, carbonates, etc. were determined.

Piperitenone oxide was the main compound (around 80 %) in two populations of *Mentha suaveolens* L, although the occurrence of pulegone instead of piperitenone oxide was found in some samples. In the other location, its composition is closer to the reported for *Mentha suaveolens* ssp. *Insularis* (Req.) [3], with piperitone oxide as a main compound. Some diurnal significant differences ( $P < 0.05$ ) have been found regarding the total oxygenated monoterpenes, both for leaves and flowers. With regard to *Mentha longifolia* L, the main compounds were alpha -terpineol acetate (30-45 %) and carvone acetate (30-40 %). This composition has not been found from literature [6]. This fact should be object of further investigations.

Despite of these significant differences, these results show a great variability among samples which can be interpreted as a greater predominance of genetic factors over environmental ones. Consequently, this study will be extended to other populations by means of individual analysis.

1. A M Viljoen et al. (2006) J. Essent. Oil Res. 18: 60-65.

2. M Maffei (1988) Flavour Fragr. J. 3: 23-26.

3. S Soutour et al. (2008) Flavour Fragr. J. 23: 107-114.

4. AM Dzamic et al. (2010) Botanica Serbica, 34(1): 57-61.

5. AC Figueiredo et al. (2008) Flavour Fragr. J. 23: 213-226.

6. BM Lawrence (2006) In: BM Lawrence (Ed.) Mint: The Genus Mentha. Taylor & Francis Group, Boca Raton, US, pp. 325-346.

**P 66. Chemical characterization of the essential oil from the leaves and blossoms of “escova de garrafa” (*Callistemon viminalis*) from Minas Gerais, Brazil**

Oliveira CM<sup>1</sup>; Nelson DL<sup>2</sup>; Cardoso MG<sup>1</sup>; Miranda CASF<sup>1</sup>; Gomes MS<sup>1</sup>; Queiroz VT<sup>3</sup>; Barroso JG<sup>4</sup>; Pedro LG<sup>4</sup>; Figueiredo AC<sup>4</sup>

<sup>1</sup>Universidade Federal de Lavras, Departamento de Química, Caixa Postal 3037, Campus Universitário, Lavras, 37200-000, Minas Gerais, Brazil

<sup>2</sup>Universidade Federal de Minas Gerais, FAFAR, Departamento de Alimentos, Belo Horizonte, 31270-901, Minas Gerais, Brazil

<sup>3</sup>Universidade Federal do Espírito Santo, Departamento de Química e Física, Alegre, 29.500-000, ES, Brazil

<sup>4</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portugal

dlnelson@ufmg.br

Keywords: escova de garrafa; *Callistemon viminalis*; essential oil

*Callistemon viminalis* (Myrtaceae; Callistemon) is an ornamental plant from Australia and is commonly known as “escova de garrafa” (bottle brush). It is a shrub that reaches 3-7 feet in height. The hydroalcoholic extract has antimicrobial and insecticidal activity [1,2]. The aim of the present work was to identify and quantify the chemical constituents present in the essential oils from the leaves and flowers of *C. viminalis* by GC. The collection of plant material was performed in the Garden of Medicinal Plants of the Federal University of Lavras (UFLA/MG). The essential oil was isolated by hydrodistillation from the wood and the aerial parts of *C. viminalis*. The characterization of the essential oil was performed at the Department of Plant Biology, Faculty of Science, University of Lisbon, Portugal, by gas chromatography coupled with mass spectrometry and gas chromatography with a flame ionization detector as in [3]. In the analysis of the essential oil from the leaves (fo) and flowers (fl) of *C. viminalis*, 99.82 and 99.46%, respectively, of the constituents were identified. The same major constituents were found in both leaves and flowers in different proportions (%): 1,8-cineole (fo - 69.14, fl - 66.93),  $\alpha$ -pinene (fo - 18.91, fl - 16.00), limonene (fo - 5.89, fl - 10.04) and  $\alpha$ -terpineol (fo - 1.70; fl - 2.19). The components or combination of components responsible for the biological activities are yet to be determined.

Acknowledgments: Partially supported by CNPq, CAPES, Fapemig.

1 AF Ndomo et al. (2010) Journal of Applied Entomology 134: 333–341

2. OO Oyedeji et al. (2009) Molecules 14: 1990-1998

3. MD Mendes et al. (2011) Ind. Crop. Prod. 33: 710-719.

**P 67. Composition of volatile oil of *Mosiera bullata* species grown in Cuba.**

Saucedo Y<sup>1</sup>; Jorge E<sup>1</sup>; Ramis G<sup>2</sup>; Vander Heyden I<sup>3</sup>; Monteagudo U<sup>1</sup>; Simó EF<sup>2</sup>; Herrero JM<sup>2</sup>; Lerma MJ<sup>4</sup>; De Armas Y<sup>1</sup>; Vicet L<sup>1</sup>

<sup>1</sup>Pharmacy Department, Faculty of Chemistry, Central University "Marta Abreu" of Las Villas, C-54830 Santa Clara, Cuba

<sup>2</sup>Department of Analytical Chemistry, Faculty of Chemistry, University of Valencia, E-46100 Burjassot (Valencia), Spain

<sup>3</sup>Department of Analytical Chemistry and Pharmaceutical Technology, Faculty of Medicine and Pharmacy, Vrije Universiteit Brussel-VUB, Belgium

<sup>4</sup>Departamento de Química Analítica y Tecnología de los Alimentos, Universidad de Castilla – La Mancha, Avda. Camilo José Cela 10, E-13004, Ciudad Real, Spain.

ysaucedo@uclv.edu.cu

Keywords: *Mosiera bullata*; Myrtaceae; essential oil composition; cineol

Myrtaceae family (121 genera, 3800–5800 spp.) is one of the most important families in tropical forests. Several members of this family are used in folk medicine as antidiarrheal, antimicrobial, antioxidant, cleanser, antirheumatic, and anti-inflammatory agent and to decrease the blood cholesterol. [1-5] Objective: The aim of the study was to determine the composition of essential oil of *Mosiera bullata* leaves collected in botanical garden, Cuba, being examined by GC/MS and to evaluate the antioxidant activity of essential oil by in vitro techniques. Methods: The volatile oil was obtained from dried *Mosiera bullata* obtained by hydrodistillation and was carried out the identification of its main component by GC–MS analyses. Kováts indices, mass spectra and standard compounds were used to identify a total of 14 individual compounds. The major component found was 1.8 cineol (ca. 62%). The potential antioxidant activity was also investigated and the sample tested by 2, 2'-diphenyl-1-picrylhydrazyl (DPPH), reducing power and linoleic acid assay. Results: The sample tested was found to interact slightly with the stable free radical DPPH in a time dependent manner and this antioxidant activity was supported by the complementary antioxidant assay in linoleic acid system and TBA method. The total amount of phenolic compounds in the essential oil was determined as gallic acid equivalents. Conclusions: This research is essential to identify quality parameters of this oil and is useful for quality control in the future industrial production and for the use of essential oil as antioxidant in the pharmaceutical industry.

1. ME Alves et al. (2011). Chemistry & Biodiversity. 8: 73-94
2. R. P. Limberger et al. (2002). J. Essent. Oil Res. 2002. 14: 302
3. R. P. Limberger et al. (2002). Flavour Fragrance J. 17: 341
4. R. A. Cole et al. (2008). Chem. Biodiversity
5. 1327. 5. J. R. Silva et al (2009). Chem. Nat. Comp.45: 565.

## P 68. Essential oil and glandular trichomes of *Ocimum campechianum* in South Eastern Mexico

Calvo-Irabién LM<sup>1</sup>; Canul-Puc K<sup>1</sup>; Dzib G<sup>1</sup>

<sup>1</sup>Centro de Investigacion Cientifica de Yucatan. Calle 43 #130 Chuburna Mérida 97200, Yucatan. Mexico

lumali@cicy.mx

Keywords: aridity index; *Ocimum campechianum*; sesquiterpenes; Yucatan

*Ocimum campechianum* Mill. is an important aromatic herb traditionally used for different medicinal purposes in Mayan communities [1], particularly gastrointestinal disorders [2]. Despite the economic and therapeutic importance of *Ocimum campechianum* there is no information regarding secretory structures and the effect of environmental factors on the essential oil characteristics. The main goal of this study was to describe the morphology of glandular trichomes in *Ocimum campechianum*, and assess the variation in glandular trichome and essential oil characteristics across six populations of this species, found at sites with varying levels of aridity at southeastern Mexico. The morphology and density of glandular trichomes were investigated using a combination of stereo microscopy and SEM. Essential oil was extracted using steam distillation and was characterized with GC, GC-MS and commercial standards. Correlation analysis between essential oil yield and aridity was performed. *Ocimum campechianum* presented one type of glandular trichome, peltate, on both sides of the leaf: Peltate trichomes were clearly distinct in size and morphology. Higher size trichomes (ca 50 µm diameter) with 8 cell head, were more abundant than smaller peltate trichomes (ca 10 µm de diámetro) with a two cell head. Trichomes were uniformly distributed on both leaf surfaces. Average essential oil yield was considerable, 2.2% (w/w) and showed important variation among populations. Spearman rank correlation showed no clear effect of aridity on essential oil yield. *Ocimum campechianum* essential oil was dominated by sesquiterpenes, the main components were isoeugenol, caryophyllene (E-), camphor, germacrene D, bicyclogermacrene, α-humulene, spathulenol and β-elemene.

Acknowledgments: This study was financially supported by CONACyT grant given to LCI (YUC-2008 C06-108231). We thank Fabiola Escalante Erosa, Lilia Can and Cindy Cachón, for technical assistance in laboratory work.

1. Roys R.L. (1931). The Ethnobotany of the Maya. Toulane University. New Orleans.
2. Ankli A et al. (1999). Econ. Bot. 53:144-160

**P 69. Combined treatment of mung sprouts contaminated with salmonella with vapors of oregano essential oil and low pressure**

Frankova A; Pavel Kloucek P; Jakub Smid J

University of Life Sciences Prague

frankovaa@af.czu.cz

Keywords: mung sprouts; *Salmonella enteritidis*; oregano essential oil; low pressure treatment

Seed sprouts are in general recognised as a good source of vitamins and minerals in human diet. On the other hand, due to the way they are processed, they present potentially high health risk for the consumers as also identified in 2073/2005 EC [1]. High temperature, humidity, and lack of special treatment during the seed sprouting makes ideal conditions for the growth of bacterial flora including the pathogenic one (e.g. *Escherichia coli*, *Salmonella enteritidis*, *Listeria monocytogenes*). Moreover, recent outbreak (*E. coli* in sprouts) has proved the need for the introduction of the new steps in the production chain, which will minimize the risk of pathogenic bacteria presence in final products. Attempts to treat the seeds before sprouting with different types of chemicals as chlorine or essential oils (EOs), high temperature and pressure sometimes brought promising results, however, so far there is no reliable method which would lead to the reduction of all pathogenic bacteria in the sprouts[2].

Thus the aim of this study was to evaluate if treatment of mung sprouts with vapor phase of oregano EO in combination with low pressure can significantly reduce the presence of *Salmonella* in the sprouted seeds.

Seeds of mung bean (*Vigna radiata*) were inoculated with 7 log CFUg<sup>-1</sup> of *Salmonella enteritidis* ATCC 13076 and sprouted for 4 days. After that they were treated with oregano EO (1 and 0.1 µl/ml) in combination with low pressure and fast evaporation of the EO supported by heating. Time of the treatment was 5, 15 and 60 min. The same tests series was also done under atmospheric pressure. The numbers of *S. enteritidis* was investigated immediately and fourth day after the treatment. Reduction of the *S. enteritidis* ranged from 0.3 to 3 log CFUg<sup>-1</sup>. The most significant reduction of the bacteria was observed when the sprouts were treated for 5 and 15 min under the low pressure with heated EO.

The results have shown that combination of the EO, low pressure and heat can lead to significant and in compare to other studies quick reduction of *S. enteritidis* in final product. Thus it has a potential to be used in the real sprout production.

Acknowledgements: This study was supported by CIGA 21160/1313/3103 and S grant of MSMT CR

1. Commission Regulation (EC) No 2073/2005 of 15 November 2005 on microbiological criteria for foodstuffs.
2. W. R. Weissinger, K. H. MCWattres, L. R. Beuchat (2001) J. Food Protect. 64, 442–450

## P 70. Effect of *Cymbopogon winterianus* essential oil on germination and growth of *Euphorbia heterophylla*

Reis B<sup>1</sup>; Kern KA<sup>1</sup>; Marques RM<sup>1</sup>; Ishii-Iwamoto EL<sup>1</sup>; Salgueiro L<sup>2</sup>; Cavaleiro C<sup>2</sup>; Filho OF<sup>1</sup>

<sup>1</sup>Universidade Estadual de Maringá

<sup>2</sup>Universidade de Coimbra

brunoblew@hotmail.com

Keywords: Geraniol; citronellal; citronellol; germination inhibition; bioherbicide

*Euphorbia heterophylla* L. weed is an annual cycle plant, belonging to the Euphorbiaceae family widely distributed in tropical and subtropical America. This weed has a center of origin in South America (Brazil-Paraguay), which confers great genetic variability. It is currently one of the plants that cause most losses in soybeans, and show resistance to some herbicides as inhibitors of acetolactase synthase enzyme currently employed in agriculture [1]. The objective of this study was to evaluate the potential bioherbicide the essential oil of *Cymbopogon winterianus*. Fifty seeds of *E. heterophylla* L. were placed in Petri dishes containing two germitest sheets previously autoclaved and moistened with solutions (10 mL) of essential oil of *C. winterianus* ((0.03, 0.06, 0.12 and 0.25% [v / v]) diluted in Dimethyl Sulfoxide (DMSO [2% v/ v]) [2]. Four repetitions were placed randomly on germination chamber (Type BOD) with 12h photoperiod at 25°C for 7 days. Essential oil analysis was carried out by gas chromatography (GC) and by gas chromatography-mass spectroscopy (GC/MS). Major compounds were geraniol (23,9%), citronellal (32,4%), citronellol (14,6%), geranial (3,3%) and limonene (3,3%). Germination percentage ( $\%G = \sum ni.Ni \cdot 100$ ), germination average time ( $GAT = \sum ni.ti / \sum ni$ ), germination average speed (GAS) and germination speed index (GSI), root system and shoot growth, fresh and dry weight of seedlings were all affected by essential oil of *Cymbopogon winterianus*. The results indicate that one or more monoterpenes present in essential oil directly inhibits the germination of *E. heterophylla* and has potential bioherbicide effect.

Acknowledgments: The authors thank to Fundação Araucária, CAPES and Fundação para a Ciência e Tecnologia (FCT), POCI 2010/FEDER for financial support to this work.

1. KA Kern, EM Pergo, FL Kagami, LS Arraes, MA Sert, EL Ishii-Iwamoto (2009). The phytotoxic effect of exogenous ethanol on *Euphorbia heterophylla* L. Plant Physiology and Biochemistry, 47: 1095-1101.
2. HA Silva, AP Valderrama, FC Moreira, RM Marques, B Reis, CM Bonato (2012). The effect of high dilutions of *Pulsatilla nigricans* on the vigour of soybean seeds subjected to accelerated aging. Acta Scientiarum. Agronomy. 34: 201-206.

**P 71. Biological properties and essential oil composition of leaves of *Syzygium cumini* (L.) skeels from São Luís-Maranhão, Brazil**

Dias CN<sup>1</sup>; Rodrigues KAF<sup>2</sup>; Carneiro SMP<sup>2</sup>; Carvalho FAA<sup>2</sup>; Andrade EHA<sup>3</sup>; Maia JGS<sup>3</sup>; Moraes DFC<sup>1</sup>

<sup>1</sup>Programa de Pós-Graduação em Saúde e Ambiente, Universidade Federal do Maranhão, 65085-580 São Luís, MA, Brazil

<sup>2</sup>Núcleo de Pesquisas em Plantas Medicinais, Programa de Pós-Graduação em Farmacologia, Universidade Federal do Piauí, 64049-550 Teresina, PI, Brazil

<sup>3</sup>Programa de Pós-Graduação em Química, Universidade Federal do Pará, 66075-900 Belém, PA, Brazil

cla\_noleto@yahoo.com.br

Keywords: larvicidal; molluscicidal; leishmanicidal; *Syzygium cumini*

*Syzygium cumini* (Myrtaceae) is usually known as black plum or jambolan. In this work the essential oil of leaves of *S. cumini* was tested against *Biomphalaria glabrata* snails, *Aedes aegypti* larvae, that are schistosomiasis and dengue vectors, respectively and an aquatic non-target organism. In addition, potential anti-*Leishmania* was evaluated as well as the hemolytic activity and the chemical composition of this essential oil. The essential oil was obtained by hydrodistillation and then analyzed by GC/MS. For the larvicidal activity, solutions of 25 to 200mg/L from the oil were tested against 4th instar larvae of *A. aegypti* [1]. For the molluscicidal activity, the snails were exposed to solutions of 25 to 100 mg/L [2]. For toxicity evaluation on a non-target organism, 2nd instar larvae of *Artemia salina* were used [3]. For leishmanicidal activity, the oil was tested against *Leishmania amazonensis* at concentrations of 3.1 to 400 mg/L [4]. The hemolytic activity was evaluated using human erythrocytes [5]. All the tests were carried out in triplicate and the results are shown as LCs and percent of hemolysis. The leaves of *S. cumini* provided oil, with a yield of 0.5%. The analysis of this oil showed the identification of 11 compounds, representing 99.98% of the total oil fraction, with monoterpenes comprising 92.46% and sesquiterpenes 7.52%. The major compounds were the monoterpene hydrocarbons  $\alpha$ -Pinene (31.85%), *cis*- $\beta$ -Ocimene (28.98%) and *trans*- $\beta$ -Ocimene (11.71%). In the larvicidal test, the oil was not lethal even at the highest concentration. In the molluscicidal test, toxicity tests to a non-target organism and leishmanicidal tests, the LC<sub>50</sub> were 90.4mg/l, 175.4mg/l and 60.34mg/l, respectively. In the hemolytic test, 7.5, 4.7 and 3.7% of hemolysis were observed at the concentrations of 400, 200 and 100 mg/l, respectively. The essential oil of the leaves of *S. cumini* showed promising activity as a molluscicide and leishmanicide.

Acknowledgments: CAPES, FAPEMA

1. World Health Organization (2005) WHO/CDS/WHOPES/GCDPP/2005/3.
2. WHO (1965) Bul. of the WHO 33:567-576.
3. J de S Luna et al. (2005) J. Ethnopharmacol. 97:199-206.
4. F Oliveira-Silva et al. (2008) Am. J. Trop. Med. Hyg. 78:745-749.
5. DG Valadares et al. (2011) Parasitol. Int. 60: 357-363.



## P 72. Chemical composition and larvicidal activity of leaf essential oils from *Mentha pulegium* and *Foeniculum vulgare*

Rocha DK<sup>1</sup>; Matos OC<sup>2</sup>; Novo MT<sup>1</sup>; Delgado M<sup>3</sup>; Moiteiro C<sup>4</sup>

<sup>1</sup>Instituto de Higiene e Medicina Tropical, UNL, Rua Junqueira, 100, 1349-008 Lisboa, Portugal. Unidade de Parasitologia e Microbiologia Médicas, IHMT, Rua Junqueira, 100, 1349-008 Lisboa, Portugal.

<sup>2</sup>Instituto Nacional de Recursos Biológicos, INIAV. Quinta do Marquês, Av. República, 2784-505 Oeiras, Portugal. Unidade de Parasitologia e Microbiologia Médicas, IHMT, Rua Junqueira, 100, 1349-008 Lisboa, Portugal.

<sup>3</sup>Delegação do Ministério do Desenvolvimento Rural, MDR. Porto Novo, Santo Antão, Cabo Verde.

<sup>4</sup>Departamento de Química e Bioquímica, Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1149-016, Lisboa, Portugal.

oliviamayos@sapo.pt

Keywords: *Aedes aegypti*; dengue; vector control; aromatic plants

Dengue is a potentially fatal mosquito-borne infection with 50 million cases annually, and 2.5 billion of people vulnerable to the disease, transmitted mainly by *Aedes aegypti* (Linnaeus, 1762). This major public health problem has epidemics in Brazil and recently in Cape Verde [1]. The lack of anti-viral treatment or vaccine against dengue makes mosquito vector control the only option to prevent or reduce virus transmission. The constituents of plant essential oils can affect insect's behavior being potentially effective in pest control [2,3,4]. In our studies we intended to contribute to control the development of earlier *Ae. Aegypti* instars. Essential oils (EOs) obtained by hydrodistillation of the aerial parts of *M. pulegium* and *F. vulgare*, were assayed on third instar larvae.

For EOs larvicidal activity four concentrations were assayed. Tween 20 was used to obtain emulsions of the EOs in water. The lethal concentrations LC<sub>50</sub> and LC<sub>90</sub> were determined by probit regression (SPSS for Windows®). The <sup>13</sup>C-NMR technique provides reliable qualitative information of the main chemical constituents of the total oil content. The <sup>13</sup>C-NMR analysis of *M. pulegium* oil showed that pulegone was the major monoterpene ketone identified with traces of menthone, whereas trans-anethole and limonene were the main constituents of *F. vulgare*. The EOs of these plants showed strong larvicidal activity 24h after exposure. 100 % mortality of 3<sup>rd</sup> instar larvae of *Ae. aegypti* was achieved with 0.060 ml/L of the wild fennel (*F. vulgare*) essential oil. Preliminary bioassays showed that 3<sup>rd</sup> instar larvae of *A. aegypti* are more susceptible to *F. vulgare* EO than to that of *M. pulegium*.

Acknowledgments: Authors are grateful to Fundação Calouste Gulbenkian for the PhD grant and to FCT for financial support of UPMM and project PEst-OE/QUI/UI0612/2011.

1. WHO, 2009. Dengue: guidelines for diagnosis, treatment, prevention and control New edition.
2. Burfield T, Reekie SL., 2005. Mosquitoes, malaria and essential oils. *Int. J. Aromather*; 15: 30–41.
3. Nerio LS, Olivero-Verbel J, Stashenko E, 2010. Repellent activity of essential oils: a review. *Bioresour Technol*; 101: 372-378.
4. Shaaya E and Rafaeli A., 2007 Essential Oils as Biorational Insecticides–Potency and Mode of Action. In, *Insecticides Design Using Advanced Technologies*. Eds: Ishaaya I., Nauen R and Horowitz AR. Springer-Verlag Berlin Heidelberg.

### P 73. Thyme and juniper essential oils screening for light dependent antifungal properties

Matos OC<sup>1</sup>; Cavaleiro C<sup>2</sup>; Salgueiro L<sup>2</sup>

<sup>1</sup>Instituto Nacional de Recursos Biológicos, INIAV. Quinta do Marquês, Av. República, 2784-505 Oeiras. Portugal. Unidade de Parasitologia e Microbiologia Médicas, IHMT, Rua Junqueira, 100, 1349-008 Lisboa, Portugal.

<sup>2</sup>Centro de Estudos Farmacêuticos, Faculdade de Farmácia, Pólo das Ciências da Saúde, Azinhaga de Santa Comba, Universidade de Coimbra, 3000-548 Coimbra, Portugal

oliviamatos@sapo.pt

Keywords: Essential oils; photoactivity; phytopathogenic fungi, antifungal activity

Plant screening for light activated bioactivity is an issue increasing importance. Essential oils of *Juniperus communis* spp. *alpina* and *Thymus zygis* spp. *zygis* were analyzed for photoactive antifungal properties. Phytopatogenic fungi *Botrytis cinerea*, *Cladosporium cucumerinum* and *Fusarium culmorum*, reported to be harmful toxin-producers, were used as biological targets.

For both plant oils 1:10, 1:50 and 1:100 v/v solutions were prepared in n-hexane. Fungi were grown and maintained in culture on PDA (Difco Lab) at 25°C. For the antifungal tests fungi suspensions were prepared by disintegrating the mycelium in a specific liquid medium. Each fungus suspension was filtered under vacuum through a nylon net (100 µm pores) on a Buckner funnel, adding liquid media until a spore count of about  $5-7,5 \times 10^4$  per ml was achieved for each fungus. Stock solutions (10<sup>-4</sup> M) of 8-methoxypsoralen (8-MOP) and 8-methoxytione-psoralen (8-MOPT) were used as standards respectively for the irradiation with the UVA and VIS spectrum of light. For the antifungal tests performed on TLC plates 25 and 50 µl of each concentration of the plant oil, and 25 µl of the solution of the standards were applied. Control spots only with the solvent were also assayed. Fungi were inoculated by spraying the TLC plates with the fungal suspension. Incubation was performed in humid chambers at 23±2°C for 4-5 days. Inhibition areas were revealed as spots without fungus growth. The quantitative evaluation was based on the area of the spots free of fungus growth. Photoactivity was determined by comparing the results achieved in the dark to those of the correspondent sample irradiated. A set of five fluorescent lamps (Osram L18w/10 Daylight, emission range ~350-730 nm,  $\lambda_{max}$ = 485nm with 0,62W/m<sup>2</sup> fluence) or a set of five UVA lamps (Phylips TLK 40W/09N; emission range ~310-440nm;  $\lambda_{max}$ =354nm, 0,44W/m<sup>2</sup> fluence) were used as VIS and UVA light sources, respectively.

Relevant antifungal effects were obtained with both plant oils on the different fungi, however EO were slightly less efficient than the standards 8-MOP and 8-TMOP in relation to UVA or VIS light irradiation, respectively. As a whole EO photoactivity was higher after irradiation with UVA light than with VIS light. Best photoactivity was observed with Juniper oil (~72.3% UVA and VIS) on *C.cucumerinum* and *B.cinerea*, and Thyme oil (80.3% (UVA) and 73.8 (VIS) on *C.cucumerinum*.

Acknowledgements: Authors thank Prof. António Maçanita, from IST-UTL, for supplying the standards.

**P 74. Citral, a natural acyclic monoterpene, reduces orofacial nociception in mice**

Barreto RSS; dos Santos MRV; Quintans JS; Bonjardim LR; Barreto AS; Quintans Júnior LJ; Lima PSS; Antonioli AR

*Federal University of Sergipe - Brazil*

*rosanasfisio@hotmail.com*

Keywords: Monoterpenes, citral, orofacial pain, analgesic

The purpose of the present study was to evaluate the antinociceptive effect of citral (CIT) using orofacial nociception induced by -formalin, -capsaicin and -glutamate on mice.

Swiss male mice (26-32g) were pretreated with CIT (50, 100 or 200 mg/kg, i.p.), morphine (5 mg/kg, i.p.) or vehicle (distilled water + one drop of Tween 80 0.2%) 1 h before formalin (20 µl, 2%), capsaicin (20 µl, 2.5 µg) or glutamate (40 µl, 25 mM) injection into the right upper lip of animals. The motor coordination was also evaluated using Rota rod apparatus (8 rpm, 180 s). Experimental protocols were approved for the UFS Ethic Committee (CEPA # 26/09). The data were statistically analyzed by ANOVA followed by Dunnett test ( $p < 0.05$ ).

Our results revealed that i.p. pretreatment with CIT was effective in reducing nociceptive face-rubbing behaviour in both phases on formalin test ( $p < 0.01$  or  $P < 0.001$ ) and also produced significant antinociceptive effect at all doses in the capsaicin- and glutamate- induced orofacial nociception tests ( $p < 0.001$ ). Such results were unlikely to be provoked by motor abnormality.

Our results suggest that citral, a natural acyclic monoterpenes, might represent important tool for management and/or treatment of orofacial pain.

Acknowledgements: FAPITEC/SE/Brazil; CNPQ/Brazil.

**P 75. Chemical characterization of the essential oil of *Hyptis pectinata* germplasm**

Jesus AS<sup>1</sup>; Blank AF<sup>1</sup>; Arrigoni-Blank MF<sup>1</sup>; Lima RN<sup>2</sup>; Alves PB<sup>2</sup>

<sup>1</sup>Departamento de Engenharia Agrônômica, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil.

<sup>2</sup>Departamento de Química, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil.

arie.blank@gmail.com

Keywords: Lamiaceae; medicinal and aromatic plant; accessions; chemical constituents; volatile oil

The aim of this study was to characterize chemically the essential oil of 14 accessions of *Hyptis pectinata* L. Poit from the Active Germplasm Bank of the Federal University of Sergipe, Brazil. The field experiment was conducted at the "Campus Rural da UFS" Research Farm of the Federal University of Sergipe, located in São Cristóvão, Sergipe, Brazil. The essential oil was obtained from dry leaves by hydrodistillation using a Clevenger apparatus. Qualitative analysis of the chemical composition of the essential oils was performed using a gas chromatograph coupled to a mass spectrometer. A quantitative analysis of the essential oils was conducted in a gas chromatograph equipped with a flame ionization detector. We do not observed significative differences between accessions for essential yield. Variations for chemical constituents of the essential oil from *H. pectinata* accessions were observed. Compounds encountered in major quantity in the essential oils of the accessions were  $\beta$ -elemene, (E)-caryophyllene, biciclogermacrene, trans cadina-1,4-diene, espatulenol, caryophyllene oxide and calamusenone. Major percentages of (E)-caryophyllene were observed in the accessions SAM-014 (43.38%) and SAM-015 (31.70%). Highest percentages of caryophyllene oxide were observed in the accessions SAM-010 (44.89%) and SAM-008 (44.69%) and major percentages of calamusenone were observed in the accessions SAM-017 (37.00%) and SAM-018 (29.04%). The multivariate analysis showed the formation of four groups according to the chemical composition of the essential oil of *H. pectinata* accessions.

Acknowledgments: The authors thank FAPITEC/SE, CNPq and CAPES for their financial support.

**P 76. Phytochemical analyses and antinociceptive effect of *Aristolochia trilobata* L. stem essential oil and 6-methyl-5-hepten-2yl acetate in rodents**

Santos DA; Alves PB; Barreto RSS; Costa EV; Quintans JS; dos Santos MRV; Quintans-Júnior LJ

Federal University of Sergipe - Brazil

rosanasfisio@hotmail.com

Keywords: *Aristolochia trilobata*, essential oil, 6-methyl-5-hepten-2yl acetate (sulcatyl acetate), pain

*A. trilobata* L. (Lamiaceae), popularly known as 'mil-homens', produces an essential oil that is generally sold at popular Brazilian markets to treatment of colic, diarrhea, and dysentery. Present study evaluated phytochemical composition of *A. trilobata* stem essential oil (EO) and antinociceptive property of EO and 6-methyl-5-hepten-2yl acetate (sulcatyl acetate), mainly compound, in two models of pain.

EO was extracted by hydrodistillation of dried stem of the *A. trilobata* (UFS Herbarium voucher specimens, ASE#23.161). The identification of the components was made through comparison of substance mass spectrum with the database of the GC-MS, literature and retention index. Sulcatyl acetate (SA) was isolated and identified from OE. The antinociceptive effect of EO and SA were examined using the acetic acid (0.85%) writhing reflex and formalin (1%)-induced nociceptive behavior in mice. Male mice (34-38g) (n=8, per group) were pretreated with vehicle (saline + tween 80 0.2%) or EO (25, 50 or 100 mg/kg, i.p.) or SA (25 or 50 mg/kg, i.p.) 1 h before experiments. Protocols were approved by the animal care and use Committee (CEPA/UFS # 16/12) at the UFS. Data were evaluated by one-way ANOVA followed by Tukey's test ( $p < 0.05$ ).

The phytochemical analysis of EO showed the presence of 6-methyl-5-hepten-2yl acetate (sulcatyl acetate) (25.45%), limonene (12.07%), linalool (7.34%) and p-cymene (7.14%) as the main compounds. The presence of sulcatyl acetate in EO was confirmed by IR, MS and 1D (1H and 13C) and 2D HSQC and HMBC NMR experiments. Our results revealed that i.p. pretreatment with EO and SA were effective in reducing nociceptive behavior in writhing reflex test ( $p < 0.001$ ) and both phases on formalin test ( $p < 0.01$  or  $p < 0.001$ ). Such results were unlikely to be provoked by motor abnormality.

Our results suggest that EO and SA might represent important tool for management and/or treatment of pain.

Acknowledgments: FAPITEC/SE/Brazil, CAPES/Brazil, CNPq/Brazil.

**P 77. Impact of hybridization in *Ocimum basilicum* on the volatile and sensory profile.**

Costa AS<sup>1</sup>; Arrigoni-Blank MF<sup>1</sup>; Silva MAAP<sup>2</sup>; Blank AF<sup>1</sup>; Santos DA<sup>3</sup>; Alves PB<sup>3</sup>

<sup>1</sup>Departamento de Engenharia Agrônômica, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil.

<sup>2</sup>Departamento de Tecnologia de Alimentos, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil.

<sup>3</sup>Departamento de Química, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil.

fatima.blank@gmail.com

Keywords: basil; essential oil; quantitative descriptive analysis; principle component analysis

The aim of this study was to develop the volatile and sensory profile by Quantitative Descriptive Analysis (QDA) using essential oil of three basil (*Ocimum basilicum* L.) hybrids ('Cinnamom' x 'Maria Bonita'; 'Sweet Dani' x 'Cinnamom'; 'Sweet Dani' x 'Maria Bonita'). The plants were cultivated at the "Campus Rural da UFS" Research Farm of the Federal University of Sergipe, located in São Cristóvão, Sergipe, Brazil. The essential oils were obtained from dry leaves by hydrodistillation using a Clevenger apparatus. Qualitative analysis of the chemical composition of the essential oils was performed using a gas chromatograph coupled to a mass spectrometer (GC-MS). A quantitative analysis of the essential oil components was conducted in a gas chromatograph equipped with a flame ionization detector (FID). Twelve descriptive terms were developed by a team of selected judges, which also led to the definition of each term and the reference samples. Data were analyzed by ANOVA, Tukey test and principle component analysis. The sample 'Cinnamom' x 'Maria Bonita' had stronger overall aroma intensity than the others and less citrus aroma. Hybridization favored the appearance of new compounds in the essential oil of hybrid 'Cinnamom' x 'Maria Bonita', such as trans-linalool oxide, (E)-methyl cinnamate, (Z)-methyl cinnamate, which does not exist in the parent's essential oil. For the hybrids 'Sweet Dani' x 'Cinnamom' and 'Sweet Dani' x 'Maria Bonita' the new compound was trans-linalool oxide.

Acknowledgments: The authors thank FAPITEC/SE, CNPq and CAPES for their financial support.

## P 78. Comparative study of volatile oil profile of different *Rosmarinus officinalis* L. extracts

Olah NK<sup>1</sup>; Hanganu D<sup>2</sup>; Socaci S<sup>3</sup>; Morgovan C<sup>4</sup>; Mos L<sup>5</sup>; Toma CC<sup>6</sup>

<sup>1</sup>Department of Pharmaceutical Industry, Faculty of Pharmacy, "Vasile Goldis" West University of Arad, 1 Feleacului Street, 300041 and SC PlantExtrakt SRL, 407059, Rădaia, Jud. Cluj, Romania

<sup>2</sup>Department of Pharmacognosy, Faculty of Pharmacy, "Iuliu Hatieganu" University of Medicine and Pharmacy, Cluj – Napoca, 15 I.Creangă Street, Romania

<sup>3</sup>Department of Food Science and Technology, University of Agricultural Sciences and Veterinary Medicine, 64 Calea Floresti street, Cluj-Napoca, Romania

<sup>4</sup>Department of Pharmaceutical Management, Marketing and Legislation, Faculty of Pharmacy

<sup>5</sup>Faculty of Medicine

<sup>6</sup>Department of Pharmacognosy, Faculty of Pharmacy,

olahdr@aol.com

Keywords: *Rosmarinus officinalis*; volatile oil profile; GC-MS; fresh plant; dry plant

The Rosemary (*Rosmarinus officinalis*) is a well known medicinal and culinary herb [1]. The purpose of this study was to compare the volatile oil profile of different Rosemary extracts obtained from dry and fresh herb, highlighting the differences between the extracts used in aroma-, phyto- and gemmotherapy. The main topic of present study is to highlight the differences between the dry and fresh Rosemary's volatile oil profile and the specific volatile oil profile of the special gemmotherapeutic Rosemary extract.

There were studied the volatile oil separated by hydrodistillation from dry plant, the hydroalcoholic extracts (1:5 – dry plant:solvent) obtained from fresh respectively dry plant and the gemmotherapeutic extract obtained from fresh plant (1:20 – dry plant:solvent) [2]. The volatile oil profile was evaluated by GC-MS using headspace injection at 85°C, for 15 min, ZB-5MS 50 m x 0,32 mm x 0,25 microm capillary column respectively helium as carrier gas. The separated compounds were identified using an MS spectra library. The quantitative determination was performed by normalization.

The results shown the separation of 38 compounds from volatile oil, 33 from fresh and 13 from dry plant hydroalcoholic extracts respectively 28 from the gemmotherapeutic extract. The main separated compounds were alpha-pinene, 1,8-cineol, camphene, d-limonene and cymene. It could be observed a significant difference between the 4 volatile oil profile. The gemmotherapeutic extract has higher content in 1,8-cineol than in alpha-pinene. Generally, the fresh plant extracts contain more 1,8-cineole, cymene, d-limonene and less alpha-pinene or camphore than the dry plant extract.

This study showed a significant difference between the extracts from dry, respectively fresh Rosemary. The fresh plant extracts are more rich in volatile oil components than the dry plant extracts. The gemmotherapeutic Rosemary extract has a high content of volatile oil in comparison with its extraction ratio. These differences in the volatile oil profiles of the studied extracts are due by the degradation processes that occur during the drying of plants or the extraction. This shows the importance of using fresh plants in the extraction, having undestroyed active compounds.

1. Czygan I. et Czygan F.C. (1997) Rosmarin – *Rosmarinus officinalis* L. In ZPZ, 18(3): 182-186.

2. European Pharmacopoeia 7 (2011) Medpharm Scientific Publisher, Stuttgart.

## P 79. Chemical composition of *Xylopi*a *rubescens* leaf oil: structure elucidation of new compounds

Tomi F<sup>1</sup>; Yapi TA<sup>1</sup>; Boti JB<sup>1</sup>; Ahibo AC<sup>1</sup>; Bighelli A<sup>2</sup>; Casanova J<sup>2</sup>

<sup>1</sup>Université de Cocody-Abidjan

<sup>2</sup>Université de Corse-CNRS

*felix.tomi@univ-corse.fr*

Keywords: furanoguai-1,4-diene; (8Z,11Z,14Z)-8,11,14-heptadecatrien-2-ol; structure elucidation; *Xylopi*a *rubescens*; leaf oil

The aim of this work was to investigate the chemical composition of *Xylopi*a *rubescens* leaf oil. The structure of two new compounds has been elucidated.

Analysis of the essential oil and structural elucidation were carried out using a combination of chromatographic (CC, GC with retention indices) and spectroscopic techniques (MS, <sup>13</sup>C NMR, 2D NMR).

Fifty components, accounting for 91.9% of the whole composition, were identified including various compounds whose spectroscopic data were absent on commercial computerized MS libraries. The structure of two new compounds, furanoguai-1,4-diene A and (8Z,11Z,14Z)-8,11,14-heptadecatrien-2-ol B, has been elucidated using two-dimensional NMR spectroscopy (Figure 1). Among the minor compounds, 1βH,7αH,10βH-guaia-4,11-diene is reported for the first time as natural compound.

The composition of *Xylopi*a *rubescens* oil, dominated by furanoguai-1,4-diene (36.4%) A, differs drastically from the composition of leaf oils of other *Xylopi*a species.

Acknowledgements: Thanks to Pr Aké Assi (identification of the plant), Dr Casabianca (MS-TOF spectra) and the Ministère de l'Enseignement Supérieur de Côte d'Ivoire (Research grant to T.A.Yapi).



**P 80. Identification and quantification of germacrenes A, B and C in *Cleistopholis patens* (Benth.) Engler & Diels essential oils**

Ouattara ZA<sup>1</sup>; Boti JB<sup>2</sup>; Ahibo CA<sup>2</sup>; Tomi F<sup>1</sup>; Casanova J<sup>1</sup>; Bighelli A<sup>1</sup>

<sup>1</sup>Université de Corse-CNRS

<sup>2</sup>Université de Cocody-Abidjan

[ange.bighelli@univ-corse.fr](mailto:ange.bighelli@univ-corse.fr)

Keywords: *Cleistopholis patens*; essential oil; heat-sensitive compounds; quantification; <sup>13</sup>C NMR

Identification of individual components of essential oils is generally achieved by GC-MS, in combination with retention indices. Some difficulties may occur when heat-sensitive compounds are present in the oil. For instance, it has been reported that 1,5-germacradiene derivatives isomerise thermally to the corresponding elemene through a [3.3] sigmatropic reaction. During GC(FID) and GC-MS analysis, isomerisation of germacradiene may occur or not, depending of temperature of injector, the length of the column (25-60m) and the temperature program (2-5°C/min). The aim of this study was to quantify three heat-sensitive germacrenes A, B and C using <sup>13</sup>C NMR in combination with GC(FID).

The chemical composition of one leaf oil and two trunk bark oil samples from *Cleistopholis patens* was investigated by GC (RI), by GC-MS and by <sup>13</sup>C NMR.

In the leaf oil sample, 35 compounds accounting for 90.9% of the composition have been identified.  $\beta$ -Elemene was identified by GC-MS and by <sup>13</sup>C NMR and quantified by GC(FID) (6.4%). According to GC(FID), germacrene A accounted for 0.1%. With respect to NMR analysis (mean intensity of signals), the ratio germacrene A/ $\beta$ -elemene was evaluated as 35/65 and the compounds accounted for 2.2/4.2% in the essential oil. Obviously, in our hands (GC columns, BP-1 and BP-20, 50m, 2°C/min), germacrene A rearranged to  $\beta$ -elemene and therefore, the content of  $\beta$ -elemene was overestimated.

Concerning trunk bark oils, 45 compounds accounting for 94.9% (sample I) and 98.7% (sample II) have been identified. In sample I, germacrene B and  $\gamma$ -elemene (13.5% and 6.1%, GC(FID), respectively) were identified by GC-MS, but only the signals of germacrene B were observed on the <sup>13</sup>C NMR spectrum. The thermal rearrangement of germacrene B to  $\gamma$ -elemene during GC analysis is evidenced.

In the sample II,  $\delta$ -elemene (8.2%, GC(FID)) was identified by GC-MS and by <sup>13</sup>C NMR, while germacrene C was identified only by <sup>13</sup>C NMR. One again, germacrene C is not detected by GC due to its thermal rearrangement to  $\delta$ -elemene who is overestimated ( $\delta$ -élémente: 5.6%, germacrène C: 2.6%).

From these results, it may be concluded that germacrenes A, B and C may undergo a thermal rearrangement to  $\beta$ -,  $\gamma$ - and  $\delta$ -elemene, respectively, during GC(FID) or GC-MS analyses. Accurate quantification of the three heat-sensitive compounds was achieved by the combination of GC(FID) and <sup>13</sup>C NMR spectroscopy.

## P 81. Application of dosy for the structural characterization of bergamot essential oil

Zdunic G<sup>1</sup>; Godjevac D<sup>2</sup>; Vajs V<sup>2</sup>

<sup>1</sup>Institute for Medicinal Plants Research Dr Josif Pancic

<sup>2</sup>Institute of Chemistry, Technology and Metallurgy

gzdunic@mocbilja.rs

Keywords: NMR; DOSY; bergamot; essential oil

High resolution diffusion-ordered NMR spectroscopy allows the separation of signals from different compounds based on their diffusion coefficients. The diffusion coefficients of the components should be significantly different and without overlapping signals in <sup>1</sup>H NMR spectrum. Modification of the solvent matrix in which a sample is dissolved can change the diffusion coefficients and <sup>1</sup>H NMR chemical shifts of the components. It has been demonstrated that diffusion behavior in DOSY experiments can be manipulated by adding micellar surfactant in the solution, allowing better resolution ("matrix-assisted DOSY") [1,2].

In this study, a possible application of DOSY for the structural characterization of commercial available bergamot essential oil (Frey + Lau, Germany) has been considered. The main components of the oil were limonene, linalool acetate, and linalool, as determined previously by GC-MS. In order to obtain better resolution in DOSY spectrum, various NMR solvents were used. Additionally, normal and reverse micelles were created by adding surfactants such as Triton X-100 or cetyltrimethylammonium bromide (CTAB) in order to further vary diffusion behavior of the essential oil components. The matrices used were dimethyl sulfoxide-*d*<sub>6</sub>/Triton X-100, dimethyl sulfoxide-*d*<sub>6</sub>/D<sub>2</sub>O, CDCl<sub>3</sub>, CDCl<sub>3</sub>/D<sub>2</sub>O/CTAB, D<sub>2</sub>O/Triton X-100, acetone-*d*<sub>6</sub>, hexane-*d*<sub>14</sub>, cyclohexane-*d*<sub>12</sub>, cyclohexane-*d*<sub>12</sub>/Triton X-100, toluene-*d*<sub>8</sub>, and toluene-*d*<sub>8</sub>/Triton X-100.

The DOSY spectra recorded in this study revealed that the best results can be achieved if CDCl<sub>3</sub> or toluene-*d*<sub>8</sub> are used as solvents. Addition of the surfactants had an impact on diffusion coefficients; however, the resolution in DOSY axis has not been improved.

Acknowledgments: The authors acknowledge their gratitude to the Ministry of Education and Science of Serbia for the financial support (Project No. 46013).

1. RW Adams et al. (2011) *Org. Biomol. Chem.* 9: 7062-7064.
2. CF Tormena et al. (2010) *Magn. Reson. Chem.* 48: 550-553.

**P 82. Chemical variability of the leaf essential oil of *Xylopia aethiopica* (Dunal) A. Rich. from Côte d'Ivoire**

Yapi TA<sup>1</sup>; Boti JB<sup>2</sup>; Ahibo AC<sup>2</sup>; Bighelli A<sup>1</sup>; Castola V<sup>1</sup>; Casanova J<sup>1</sup>; Tomi F<sup>1</sup>

<sup>1</sup>Université de Corse-CNRS

<sup>2</sup>Université de Cocody-Abidjan

*felix.tomi@univ-corse.fr*

Keywords: *Xylopia aethiopica*; Côte d'Ivoire; chemical composition; variability

The aim of the present study was to evidence either homogeneity or a chemical variability within 48 oil samples of *Xylopia aethiopica* isolated from leaves harvested in six Ivoirian forests.

The chemical composition of 48 leaves oil samples of *Xylopia aethiopica* was investigated by GC(FID), in combination with retention indices and by <sup>13</sup>C-NMR. Statistical analysis was obtained by combination of hierarchical clustering dendrogram and principal components analysis.

Twenty-three components accounting for 82.5-96.1% of the oil composition were identified. The composition was dominated by monoterpene hydrocarbons,  $\beta$ -pinene (up to 61.1%),  $\alpha$ -pinene, (up to 18.6%), and sesquiterpene hydrocarbons (germacrene D, up to 28.7%). Statistical analysis allowed the distinction of two groups on the basis of  $\beta$ -pinene and germacrene D contents. The chemical composition of Group I (38 oil samples) was dominated by  $\beta$ -pinene, while the Group II (10 samples) was characterized by the association  $\beta$ -pinene and germacrene D. The monoterpenes, 4,4-dimethyl-2-vinylcyclohexene B (up to 5.1%) and 3,3-dimethyl-1-vinylcyclohexene A (up to 3.5%), previously found for the first time as natural components from *X. aethiopica* root oil [1] were identified in 28 out of 48 samples.

It could be pointed out that the four forests that produce  $\beta$ -pinene-rich *X. aethiopica* leaf oil are inland forests while oil samples containing appreciable amounts of  $\beta$ -pinene and germacrene D are located near the littoral.

1. TA Yapi, JB Boti, BK Attioua, AC Ahibo, A Bighelli, J Casanova, F Tomi (2012) Phytochem. Anal., in press.

### P 83. Comparison of different extraction methods for volatile compounds of *Thymus mastichina*

Tovar IM<sup>1</sup>; Sponza S<sup>2</sup>; S-Manzanera MCA<sup>1</sup>; Schmiderer C<sup>2</sup>; Novak J<sup>2</sup>

<sup>1</sup>Intituto Tecnológico Agrario de Castilla y León. Ctra. Burgos Km. 118 C.P. 47071 Finca Zamadueñas, Valladolid, Spain

<sup>2</sup>Institute for Applied Botany and Pharmacognosy, University of Veterinary Medicine Vienna, Veterinaerplatz 1, A-1210 Vienna, Austria

mentovin@itacyl.es

Keywords: *Thymus mastichina*; volatile compounds; extraction methods; GC-MS; GC-FID

Thyme is a genus conformed by 220 species distributed throughout most of Eurasia. *Thymus mastichina* is an endemic plant of the Iberia peninsula and is traditionally used in that region [1]. Nowadays the essential oil of *Thymus mastichina* is an important product for the perfumery industry [2].

The objective of the study is to compare three different methods for extracting volatile compounds: microdistillation, hydrodistillation and solvent extraction.

A representative sample of flowers and leaves of *Thymus mastichina*, collected in the region of Castilla and Leon (Spain), was extracted with dichloromethane, distilled in a standard Clevenger type apparatus or in the Eppendorf MicroDistiller. All extraction methods were performed in four replications. The quantitative analysis of the volatile compounds was carried out by GC-MS and GC-FID with biphenyl as internal standard.

The statistical evaluation by multivariate analysis of variance shows that the profile of essential oil compounds is significantly influenced by the extraction method with most of the compounds contributing to this effect but not reacting in the same way. Not only the solvent extract was different compared to the two distillation methods, but also the results between standard Clevenger type distillation and microdistillation differed in most of the compounds. The volatile fraction of *Thymus mastichina* presents a complex volatile composition, where the main compounds are 1,8-cineol and linalool. The average amounts of 1,8-cineol were 13.6, 22.3 and 14.1 mg/g plant material for hydrodistillation, microdistillation and solvent extraction respectively, the amounts of linalool 1.3, 1.5 and 0.9 mg/g, respectively.

The study shows that the extraction method influences the extracted amounts of the single volatile compounds. Although the microdistillation presents the best results, the hydrodistillation that is commonly used in practice shows satisfactory results.

1. Morales, R. (2010) Género *Thymus* L. Flora Iberica 12: 349-409. Real Jardín Botánico, CSIC, Madrid

2. Stahl-Biskup, E., & Saez, F. (2002) Thyme, The genus *Thymus*. Taylor and Francis

## P 84. Possibilities of use rugosa hybrid roses grown in Latvia as a source of volatile oil

Sparinska A<sup>1</sup>; Rostoks N<sup>2</sup>; Kruma Z<sup>3</sup>; Alsina Ina<sup>3</sup>

<sup>1</sup>University of Latvia, Botanical Garden

<sup>2</sup>University of Latvia, Faculty of Biology

<sup>3</sup>Latvia University of Agriculture

zanda.kruma@llu.lv

Keywords: *Rosa rugosa*; volatiles; DPPH

The diploid species *Rosa rugosa* Thunb. and its hybrids have a wide range of uses. When *R. rugosa* species became known in Europe in the middle of 19<sup>th</sup> century, it was heralded as the forbear of a new race of garden roses [1]. In China the petals of full flowering *Rosa rugosa* and its hybrids are used for a wide range of medicinal products and as a source of rose oil [2]. In contrast, on coastlines of Europe and North America *Rosa rugosa* is considered as invasive species, which rapidly spreads and destroys native plant communities [3]. The aim of this work is to understand possibilities of using Latvia grown Rugosa hybrids for production of volatile oil. The composition of volatile compounds and free radical scavenging activity (1,1-diphenyl-2-picryl-hydrazil (DPPH) % inhibition) was measured in the petals of the varieties 'Ritausma', 'Sniedze', 'Zaiga', 'Liga', 'Frau Dagmar Hastrup' [4], *R. rugosa* and *R. rugosa* 'Plena'. Volatiles from roses were extracted using solid phase microextraction with subsequent separation by gas chromatography and identification by comparison of their mass spectra with mass spectral libraries. The hydrogen atom or electron donation abilities of the corresponding extracts were measured from the bleaching of the purple-colored methanol solution of DPPH [5]. The absorption was read against a blank at 517 nm. More than forty volatile aroma compounds were identified in petals. Among those, phenylethylalcohol, betacitronelol and nerol were predominant, but the composition of compounds varied among varieties. The variety 'Sniedze' scavenged ~ 90% of free radicals, which was the best result for Rugosa hybrids. *R. rugosa* 'Plena' had the highest content of volatiles, but the pedigree of all varieties, *R. rugosa*, exhibited the highest free radical scavenging activity (91.26 % of total amount). Thus, in addition to its ornamental value in Latvia *Rosa rugosa* appeared to be preferable for use as a source of volatile oil.

Acknowledgements: European Social Fund within the project «Support for Doctoral Studies at University of Latvia», National Botanic garden, PhD Dzidra Rieksta.

1. H. Bruun et al. (2005). *Journal of Ecology*, (1975), 441-470.
2. M. Isermann (2008). *Flora*, 2008.
3. Y Hashidoko et al.. (2001). *Bioscience, biotechnology, and biochemistry*, 65(9), 2037-43.
4. Dz. Rieksta (1988).. LPSR ZA BD, Riga. p.2.
5. Z. Youwei et al (2010). *Society*, (August 2011), 397-401.

**P 85. Volatiles stability of field propagated *Thymus caespititius* chemotypes**

Lima AS<sup>1</sup>; Bahcevandziev K<sup>2</sup>; Figueiredo AC<sup>1</sup>; Pedro LG<sup>1</sup>; Barroso JG<sup>1</sup>; Trindade H<sup>1</sup>

<sup>1</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, DBV, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal, C2, Campo Grande, 1749-016 Lisboa, Portugal

<sup>2</sup>Escola Superior Agrária de Coimbra, CERNAS, Bencanta, 3040-316 Coimbra, Portugal

aslima@fc.ul.pt

Keywords: *Thymus caespititius*; Chemotypes

Endemic of the NW Iberian Peninsula, including Azores and Madeira archipelagos, *Thymus caespititius* Brot. (Lamiaceae) is an aromatic plant with well characterized essential oils and possessing several chemotypes, namely thymol, carvacrol, sabinene and  $\alpha$ -terpineol [1-3]. Although genetically controlled, the biosynthesis of essential oils is strongly affected by environmental and agronomic conditions of the growing place and by the developmental stage of the plant material, among other physiological factors [4]. The present work reports a two-year study on volatiles stability of field propagated *T. caespititius* chemotypes.

A total of ten chemically different individual plant samples, originally collected at the Azores archipelago (Terceira, S. Jorge and Flores islands) and mainland north of Portugal, were grown at Escola Superior Agrária de Coimbra since 2008. Throughout 2010 and 2011, aerial part samples were collected during the vegetative and the flowering periods. Volatiles were isolated by distillation-extraction and chemical analysis performed by GC and GC-MS, as previously reported [2].

Although fluctuations were recorded in the relative amount of some oil components over the two-year study, eight of the ten individuals maintained their original chemotype, while two changed the volatile profile. Plant D1, characterized by having high amounts of sabinene (85%) in the original collection site (S. Jorge, Azores) revealed a chemical profile with sabinene (6-40%) and high amounts of carvacrol (24-47%) following field establishment at Coimbra. Chemotype F in its original collection site (S. Jorge, Azores) showed no carvacrol in the essential oil, while the field plant established at Coimbra was rich in carvacrol (43-57%) and thymol (21-26 %).

Both biotic and abiotic factors can contribute to these chemical modifications. To understand these changes, the isolation of genes involved in scent production on different chemotypes is in progress.

Acknowledgments: Partially supported by FCT - PTDC/AGR-GPL/101334/2008.

1. Lima AS et al. (2010) Acta Hort. 860: 81-86.
2. Trindade H et al. (2008) Biochem. Sys. Ecol. 36: 790-797.
3. Figueiredo AC et al. (2008) Curr. Pharm. Des. 14: 3120-3140.
4. Figueiredo AC et al. (2008) Flavour Fragr. J. 23: 213-226.

**P 86. Chemical composition and insecticidal activity of *Thuja occidentalis* and *Tanacetum vulgare* essential oils against *Alphitobius diaperinus* Panzer.**

Zawitowska B<sup>1</sup>; Szumny A<sup>2</sup>; Szczepanik M<sup>1</sup>

<sup>1</sup>Nicolaus Copernicus University, Toruń, Poland

<sup>2</sup>Wrocław University of Environmental and Life Sciences, Wrocław, Poland

beatamalina87@wp.pl

Keywords: *Alphitobius diaperinus*; essential oils

Plant extracts, including essential oils have a great potential for pest management, but their insecticidal activity varies[1]. The essential oils (EOs) from the leaves of *Thuja occidentalis* and flowers of *Tanacetum vulgare* were obtained by steam distillation. The isolation, identification and quantification of the volatile compounds was performed using a gas chromatograph (GC) coupled to a mass spectrometer (MS), a Saturn 2000 MS Varian Chrompack [2] The major components of *T. occidentalis* EO were  $\alpha$ -thujone (69,8%) followed by  $\beta$ -thujone (9.47 %) and 1-terpinen-4-ol (2.67%); the main components of *T. vulgare* were  $\beta$ -thujone (61%) and camphor (12.99%).

The insecticidal activity of the EOs and their major components,  $\alpha$ - and  $\beta$ -thujone against different larval stages of the lesser mealworm, *Alphitobius diaperinus* Panzer, a cosmopolitan pest inhabiting chicken and broiler houses in vast numbers, was evaluated under laboratory conditions. The earlier (10 days old) and later (30 days old) larval stages were reared on diets containing 1 % acetone solutions of tested compounds. Body weight gain, mortality, number of pupae and adults, and their body mass were recorded.

Insecticidal activity of both EOs and pure monoterpenes depended on the age of larvae. The highest activity against young larvae showed *T. occidentalis* EO, followed by  $\alpha$ -thujone, tansy EO and  $\beta$ -thujone. The total mortality caused by these compounds was 65.0, 59.83, 46.88, 45.63%, respectively. The differences between the activity of EO and its major component against the tested insects were not significant. The growth and developmental cycle of younger larvae was affected in opposition to the older larvae, where it was less disturbed and their development was similar to the control.

1. C Regnault-Roger (1997) Integrated Pest Manag. Rev. 2: 25-34.

2. A Szumny et al. (2010) J. Food Eng. 97: 253-260.

**P 87. The influence of packaging materials on the composition of the dried dill (*Anethum graveolens* L.) volatiles during storage**

Kruma Z<sup>1</sup>; Galoburda R<sup>1</sup>; Muizniece-Brasava S<sup>1</sup>

<sup>1</sup>Latvia University of Agriculture

zanda.kruma@llu.lv

Keywords: dill; volatiles; packaging material; SPME

Dill (*Anethum graveolens* L.) is a popular herb used in many regions, including Baltic countries. Dill is widely used for flavoring foods and beverages due to its pleasant spicy aroma. The aim of this work was to determine the best packaging material for storage of dried dill based on volatile compounds composition.

Dill leaves were cut in slices  $0.4 \pm 0.1$  cm and dried using a microwave-vacuum drier „Musson-1” (OOO „Ingredient”, St. Petersburg, Russia). Characteristic parameters of drying program were as follows: number of magnetrons was decreased along the drying process (starting at 4, and then followed by 3, and finally, 2); pressure 12.00-14.63 kPa; drum rotation speed – 6 rpm; product mass per load – 1 kg; drying time – 17 minutes.

Dried dill samples were packed in hermetically sealed monolayer polyethylene, multilayer Multibarrier 60 (MB60) and multilayer aluminium/polyethylene pouches both in vacuum and atmospheric ambience. Samples were stored at  $+20 \pm 1$  C temperature. The quality parameters were analysed before packaging and during the storage for two years. Volatiles from dried dill were extracted using solid phase microextraction (SPME) Divinylbenzene/Carboxen/Polydimethylsiloxane (DVB/Car/PDMS) fiber and analysed by gas chromatography-mass spectrometry. Compounds were identified by comparison of their mass spectra with mass spectral libraries (Nist98), and by calculation of linear retention indexes and comparison with literature data.

The main aroma compounds of dill leaves are  $\alpha$ -phellandrene and dill ether. Composition of dried dill differs from fresh dill, and significant increase in content of  $\alpha$ -phellandrene,  $\beta$ - phellandrene was detected. Results showed that more volatiles were preserved in dill packed in Multibarrier 60 material pouches, whereas, comparing packaging ambiances, more volatiles in vacuum packed samples were determined. Control sample packed in PE pouches showed the lowest results.

The composition of volatile compounds of dried dill is affected both by the packaging material and ambience in pouches, and results showed that it is very important to choose the most appropriate material for dried dill storage.

Acknowledgment: This research has been prepared within the framework of the ESF Project, Formation of the Research Group in Food Science”, Contract Nr. 2009/0232/1DP/1.1.1.2.0/09/APIA/VIAA/122.



**P 88. Aromatic compounds of traditional and novel components of chicory coffee**

Zawirska-Wojtasiak R<sup>1</sup>; Wojtowicz E<sup>2</sup>

<sup>1</sup>Faculty of Food and Nutrition Science, Poznan University of Life Sciences, Poland, 60-637 Poznan, ul. Wojska Polskiego 28

<sup>2</sup>Department of Food Concentrates in Poznan, Institute of Agricultural and Food Biotechnology, Poland, 61-361 Poznan, ul. Starolecka 40

renazaw@up.poznan.pl

Keywords: chicory coffee, aroma, traditional and novel components

Chicory coffee is identified with the category of *wellness* beverages. Aroma attributes of chicory coffee result not only from the raw materials used, but most of all from roasting process which leads to considerable changes in chemical composition of the product. The aim of the study was aroma identification of traditional (roasted chicory, barley, sugar beets) and novel (artichoke, hawthorn, lovage) raw materials of chicory coffee. Novel components have high level of chlorogenic acid which may be an aroma precursor of phenols which have spice and smoke flavor [1].

Methods: roasting process in a device by Probat Werke BRZ-2 (Emmericham Rhein), volatiles isolation by SPME/DVB/CAR/PDMS, the identification of volatiles using GC/MS 5975C VL MSD equipped with a HP-5MS capillary column (Agilent Technologies) and NIST05 library.

Results: In the aroma of roasted chicory, barley, ray, sugar beet there were identified 36, 30, 25, 15 compounds respectively. The dominated were pyrazines and phenols which are characteristic for roasted coffee. It can be concluded, that the most aromatic among the traditional materials for coffee substitute is chicory, while the sugar beet is the lowest one. The novel proposed materials are also the source of interesting odorants, particularly roasted artichoke. Among 23 identified volatiles there were: acetaldehyde, furfural, 5-methyl-2-furfural, 2-methyl-furan, 5-methyl-2-furancarboxaldehyde, 2-methoxy-4-(1-propenyl)-phenol, 2-methyl-phenol, 2,5-dimethyl-pyrazine, 3-ethyl-2,5-dimethylpyrazine, 3,5-diethyl-2-methyl-pyrazine, 3,5-diethyl-2-propyl-pyrazine, p-vinylbenzohydrazine. It should be pointed out the occurrence of several pyrazines, and phenols characteristic for coffee aroma. Some identified odorants in roasted hawthorn were following: acetaldehyde, furfural, 2-methyl-5-(1-methylethenyl)-1-one, hexadecanoic acid methyl ester and also 4-heptyl-phenol. In roasted lovage the occurrence of 2-methoxy-4-vinyl-phenol was stated, according to many authors the important odorant in natural coffee as well as other component: furfural, 5-methyl-2-furancarboxaldehyde, 4-mercaptophenol, 6-methyl-2-pyridinamine, 2-methyl-6-(2-propenyl)-phenol. According to the obtained data the novel materials may be a source of some characteristic coffee aroma components.

Acknowledgments: This study was financially supported by the National Centre of Science, Poland, Grant No 2011/01/B/NZ9/01060.

1. M N Clifford (2000) J. Sci. Food Agr. 80: 1033-1043

**P 89. Dry and wet seasons they determine the phytochemical profile of essential oils from *Copaifera langsdorffii* Desf. in different vegetation types of the Brazilian Cerrado**

de Almeida LFR<sup>1</sup>; Portella RO<sup>1</sup>; Frei F<sup>2</sup>; Marques MOM<sup>3</sup>

<sup>1</sup>Department of Botany, Institute of Biosciences (IB), UNESP - Univ. Estadual Paulista, Botucatu, Sao Paulo, Brazil

<sup>2</sup>Department of Biological Sciences, FCL, Univ. Estadual Paulista – UNESP, Assis/SP, Brazil.

<sup>3</sup>Agronomic Institute of Campinas, IAC, Campinas, Sao Paulo, Brazil

luizfernando@ibb.unesp.br

Keywords: terpenes; Cerrado; sazonalidade; climate changes

The climate change in the atmosphere provides increases of temperature and redistribution of averages of rainfall (1). We can observe a decrease in soil moisture, could lead to increased periods of drought, especial in tropical and savanna biomes (2). The dry and wet seasons from different configurations of the Cerrado physiognomical domain were able to promote changes in the essential oil of *Copaifera langsdorffii* Desf. Different ecophysiological conditions of regions and climate change may be important factors for changes the chemical profile of terpenes in the essential oil. The study carried out tries to understand the effect of dry periods (August to September) and rainy (October to February) in the chemical profile of essential oil of *C. langsdorffii*. To this end, the research objectives are: (i) make comparisons in relation to production, yield and essential oil composition between the dry and rainy season, (ii) evaluate the phytochemical profile in different formations of the Cerrado. Analysis by GC-MS oil *Copaifera langsdorffii* 24 showed the presence of volatile compounds being monoterpenes, sesquiterpenes and oxygenated sesquiterpenes. The cluster analysis by the dissimilarity matrix calculated by the Euclidean distance which method of Square agglomeration was UPGMA gathering areas according to the chemical profile of the essential oil. Germacrene-D, gamma-muuroleno, delta-elemene, alpha-humulene and trans-caryophyllene are substances which varied concentrations. The seasonality effect is most evident at sites of semideciduous forest than in the cerrado sensu. Nevertheless, the production of essential oils correspond an effect of adjustment to the climate, with increases protection against stress (3).

Acknowledgments: Funding body: Fundação de Amparo a Pesquisa do Estado de São Paulo (FAPESP) financial support to the project nº 2010/15585-6.

1. IPCC: *Climate Change: Synthesis Report*. Geneva, Switzerland, 104 p (2007).

2. J. Sheffield and E.F. Wood. *Climate Dynamics*, 31, 79-105 (2008).

3. E. Grøndahl and B.K. Ehlers. *J. Ecol.*, 96, 981-992 (2008).

**P 90. Antimicrobial activity of essential oils against *Paenibacillus larvae***

Gende LB<sup>1</sup>; Afonso Pires SM<sup>2</sup>; Fernandez NJ<sup>1</sup>; Damiani N<sup>1</sup>; Churio MS<sup>3</sup>; Fritz R<sup>4</sup>; Eguaras MJ<sup>1</sup>

<sup>1</sup>Laboratorio de Artrópodos-FCEyN, UNMDP, CONICET, Argentina

<sup>2</sup>Centro de Investigação de Montanha (CIMO)/Escola Superior Agrária, Instituto Politécnico de Bragança, Portugal

<sup>3</sup>Departamento de Química- FCEyN, UNMDP, CONICET, Argentina

<sup>4</sup>Departamento de Química- FCEyN, UNMDP, Argentina

spires@ipb.pt

Keywords: essential oils; antimicrobial activity; *Paenibacillus larvae*

American foulbrood is a serious bacterial disease that affects *Apis mellifera* colonies; the causative agent is *Paenibacillus larvae* [1]. The aim of the study was to evaluate *in vitro* the antimicrobial activity of 32 essential oils against *P. larvae*. Oils from 21 botanical species were analyzed by gas chromatography (CG and CG/EM). All essential oils were classified according to the composition of their main components in two groups: benzene ring compounds (BRC) and terpene compounds (TC). Minimal inhibitory concentration (MIC) in MYT broth [2] was assessed by the microdilution method. Final serial dilution concentrations of the essential oils ranged between 2,000-12.5 mg/L. The bacterial isolates were collected from different Argentina's region. The chromatographic analysis showed that a 67% of the essential oils contained predominately terpene compounds, while the remaining 33% included mainly compounds with benzene rings. From the TC group, *Cymbopogon citratus* essential oil showed the better antimicrobial activity against *P. larvae* with MIC values between 150 and 250 mg/L. The essential oils from *Aloysia polystachya* and *Mentha spp.* had the lowest inhibitory activity. Among the oils from the BRC group, one of the lowest MIC values was found with cinnamon essential oil (*Cinnamomun zeylanicum*) being between 25 and 50 mg/L; *Origanum vulgare* showed the highest MIC values (350-400 mg/L). Essential oils, especially those with BRC in their composition, presented inhibitory capacity against *P. larvae* strains.

Acknowledgments: This work was supported by UNMDP and CONICET.

1. E Genersch (2010) J Invertebr Pathol. 103: 10–19.

2. LB Gende, MJ Eguaras, R Fritz (2008) Rev Argent Microbiol. 40:147-150.

### P 91. Chemical composition of leaf and stem oils of *Cupressus tonkinensis* silba from Vietnam

Bazzali O<sup>1</sup>; Tran HT<sup>2</sup>; Nguyen TH<sup>2</sup>; Pham VT<sup>2</sup>; Phan KL<sup>2</sup>; Tran MH<sup>2</sup>; Tomi F<sup>1</sup>; Casanova J<sup>1</sup>; Bighelli A<sup>1</sup>

<sup>1</sup>Université de Corse-CNRS

<sup>2</sup>Institut of Ecology and Biological Resources, Hanoi

ange.bighelli@univ-corse.fr

Keywords: *Cupressus tonkinensis*; essential oil composition;  $\alpha$ -pinene; sabinene; cedrol; <sup>13</sup>C-NMR

*Cupressus tonkinensis* was treated for a long time as a synonym of *C. torulosa* D. Don. and it was proposed as new species in 1994.[1] The two species differed by morphological characters and they were distinct in their RAPDs.[2] The aim of this work was to investigate the chemical composition of *C. tonkinensis* leaf and stem oils from Vietnam.

Leaves and stems of *C. tonkinensis* produce monoterpene-rich oils whose composition is investigated for the first time by combination of chromatographic and spectroscopic techniques. The two samples were analyzed by GC(RI), GC/MS and <sup>13</sup>C-NMR. Moreover, the leaf oil sample was also fractionated over silica gel column and all the fractions of chromatography analyzed by GC(RI) and <sup>13</sup>C-NMR.

The combination of analytical methods allowed the identification of 47 compounds, accounting for 95.6 and 87.9% of volatile constituents of the essential oils of leaves and stems respectively. Alpha-pinene (25.7%), sabinene (23.3%) and terpinen-4-ol (12.6%) were the major components of the leaf oil that contained also beta-elemol (3.8%). In contrast, alpha-pinene (48.3%) was by far the major component of the stem oil beside myrcene (11.6%) and cedrol (8.7%).

The composition of the leaf oil of *C. tonkinensis* from Vietnam is close to that of the leaf oil from *C. torulosa* cultivated in the Argentinean Patagonia [3] but stem oil differs by its high amount of  $\alpha$ -pinene, myrcene and cedrol. Leaf and stem oils of *C. tonkinensis* differ also from the essential oil isolated from the foliage of *C. torulosa* var. *cashmeriana* trees cultivated in Yunnan Province, China. [4]

Acknowledgements: The authors thank the NAFOSTED for financial support (Project 10605.21.09) and Pr. J. Costa (CPN-Lab, University of Corsica) who placed the GC/MS equipment at our disposal.

1. J Silba (1994) J. Int. Conifer Preserv. Soc. 1: 23.
2. K Rushforth et al. (2003) Biochem. Syst. Ecol. 31: 17–24.
3. RA Malizia et al. (2000) J. Essent. Oil Res. 12: 59-63.
4. LG Cool et al. (1998) Biochem. Syst. Ecol. 26: 899-913.

**P 92. Resolution of diastereoisomeric mixture of secondary bicyclic alcohol obtained from Carenko®**

Kuriata-Adamusiak R<sup>1</sup>; Strub D<sup>1</sup>; Turowska-Tyrk I<sup>2</sup>; Lochynski S<sup>3</sup>

<sup>1</sup>Department of Bioorganic Chemistry, Faculty of Chemistry, Wrocław University of Technology, Wrocław, Poland

<sup>2</sup>Institute of Physical and Theoretical Chemistry, Faculty of Chemistry, Wrocław University of Technology Faculty, Wrocław, Poland

<sup>3</sup>Institute of Cosmetology, Wrocław University College of Physiotherapy, Wrocław, Poland

renata.kuriata@pwr.wroc.pl

Keywords: terpenoids; stereochemistry; olfactory compounds

Terpenoids are broad class of compounds which can be used as intermediates for the fragrance industry and chiral synthons in organic synthesis. Their structures in some cases tend to force stereo- and enantioselective course of reactions [1].

Our studies are focused on chemical transformations of (+)-3-carene leading to compounds with potential industrial as olfactory compounds. (+)-3-Carene is a bicyclic hydrocarbon derived from common in Poland Scotch Pine (*Pinus sylvestris* L.). We used it as a starting material in two-step synthesis of bicyclo[4.1.0]heptene derivative in form of pure diastereoisomers and their mixture.

The first step to obtain pure diastereoisomers of designed compound was the chemical acetylation of 1 in acetic anhydride with addition of zinc bromide [2]. Ketone 2, commercially called Carenko® was reduced with LiAlH<sub>4</sub> which led to the desired alcohol 3). The separation of diastereoisomers were performed by several recrystallizations in mixture of methanol:water (1:1). The absolute configuration of obtained one diastereoisomer was determined by X-Ray crystallography and Mosher's esters but the second only by Mosher's esters. The olfactory properties of diastereoisomeric mixture and pure diastereoisomers were also evaluated.

Synthetic and stereochemical details of the applied procedures and olfactory properties of derivatives will be presented.

Acknowledgements: Presented work was supported by Ministry of Science and Higher Education, Grant No N N204 136538 and project „Enterprising PhD student - an investment in the innovative development of the region” part-financed by the European Union within European Social Foundation.

1. M Miyazawa et al. (2003) J. Chem. Technol. Biotechnol. 78: 620-625.
2. M Muhlstadt, P Richter (1967) Chem. Ber. 100: 1892-1897.

### P 93. Insecticidal and fungicidal activities of herb isolates obtained by different methods

Sajfrtova M<sup>1</sup>; Sovova H<sup>1</sup>; Karban J<sup>1</sup>; Pavela R<sup>2</sup>

<sup>1</sup>Institute of Chemical Process Fundamentals of the ASCR

<sup>2</sup>Crop Research Institute

sajfrtova@icpf.cas.cz

Keywords: plant protection, supercritical fluid extraction, hydrodistillation, insecticidal activity, fungicidal activity

Many green plants produce biologically active secondary metabolites and can provide valuable sources of natural drugs, natural pesticides and biofertilizers [1, 2]. In the search for alternatives to conventional pesticides, essential oils from aromatic plants have been widely investigated. In particular, terpenes and terpenoids belong to active components of essential oils [3, 4].

The aim of this study was a comparison of pesticidal activity and chemical composition of the isolates from *Pelargonium graveolens*, *Lippia javanica*, *Artemisia afra* and *Eucalyptus grandis* obtained using supercritical fluid extraction by carbon dioxide (SFE) and hydrodistillation. Extracts rich in non-polar components were obtained using SFE at pressure of 28 MPa and temperature of 50 °C. The residual plant material after SFE was subjected to maceration by methanol for 48 hours to isolate polar components. Product of hydrodistillation was pure essential oil. The yields of these isolates were evaluated for each plant material in dependence on used separation methods.

The composition of volatile compounds in the isolates was determined by gas chromatography (GC-MS and GC-FID). The insecticidal activity of isolates was measured on caterpillars of *Spodoptera littoralis* by in terms of contact toxicity (LD50, LD90) and antifeedancy. For antifungal bioassay, standard dilution method for evaluation of inhibition effect on growth of model pathogenic and toxinogenic fungi (*Fusarium oxysporum*, *Penicillium expansum* and *Aspergillus fumigatus*) was used.

Strong pesticidal effects were observed for all isolates but significant differences among the particular isolates and among the plants were found. The efficiency of CO<sub>2</sub> extracts was comparable or lower than that of hydrodistillates and higher than the efficiency of macerates. The essential oils showed the strongest insecticidal and fungicidal activities, however, their yields were 2-4 times lower than the yield of CO<sub>2</sub> extraction.

Acknowledgments: The financial support from the Technology Agency of the Czech Republic via grant TA01010578 is gratefully acknowledged.

1. PJ Landolt et al. (1999) Environ. Entomol. 28: 954-960.
2. S Abd-Ellatif et al. (2011) J. Agric. And Biol. Sci. 6: 25-32.
3. EA Klein Gebbinck et al. (2002) Phytochem. 61: 737-770.
4. SK Srivastava et al. (2003) Indian J. Chem. B. 42: 3155-3158.

**P 94. Association of *Thymbra capitata* essential oil and chitosan (TCCH hydrogel): a putative therapeutic tool for the treatment of vulvovaginal candidosis**

Palmeira-de-Oliveira R<sup>1</sup>; Gaspar C<sup>1</sup>; Salgueiro L<sup>2</sup>; Cavaleiro C<sup>2</sup>; Martinez-de-Oliveira J<sup>1</sup>; Queiroz JA<sup>1</sup>; Rodrigues AG<sup>3</sup>

<sup>1</sup>CICS-UBI - Health Sciences Research Center, Faculty of Health Sciences, University of Beira Interior, Covilhã, Portugal

<sup>2</sup>Centre of Pharmaceutical Studies, Faculty of Pharmacy, University of Coimbra, Portugal

<sup>3</sup>Cardiovascular Research & Development Unit, Faculty of Medicine, University of Porto, Porto, Portugal

ligia@ff.uc.pt

Keywords: Therapeutic tool, vulvovaginal candidosis, chitosan, *Thymbra capitata* essential oil

In this study we tested the anti-*Candida* effect of *Thymbra capitata* essential oil plus chitosan and developed a new therapeutic tool.

The classical checkboard methodology was used to determine the MIC resulting from products association [1]. The incorporation of *T. capitata* essential oil in a chitosan hydrogel using lactic acid as the solvent resulted in the TCCH hydrogel. Its anti-*Candida* activity was studied upon eighteen *Candida* cells, according to the CLSI M27-A3 micromethod [2].

The association of both natural products revealed, in the checkboard technique, an additive effect upon *Candida* and TCCH hydrogel showed to be active upon *Candida* planktonic at concentrations lower than individual compounds. TCCH hydrogel presents an acidic nature (4.3), compatible with the vaginal pH.

Being a new product with an acidic nature compatible with the vaginal environment and presenting a potent effect upon *Candida* cells, TCCH hydrogel could represent a valuable tool for the treatment of vulvovaginal candidosis.

1. Vitale SC, Afeltra J, and Dannaoui E (2005) in Ernst E.J. and Rogers P.D. (Eds) Antifungal Agents: Methods and Protocols. Humana Press, Memphis, pp. 118.
2. Clinical Laboratory Standards Institute CLSI (2008). Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts, M27-A3, Wayne, Pennsylvania.

**P 95. Essential oil and extracts from *Callitris intratropica* heartwood, sapwood and bark obtained by hydrodistillation, solvent extraction and headspace solid phase microextraction.**

Wanner J<sup>1</sup>, Schmidt E<sup>2</sup>, Jirovetz L<sup>2</sup>, Buchbauer G<sup>2</sup>

<sup>1</sup>Kurt Kitzing Co., Hintert Alten Schloss 21, 86757 Wallerstein, Germany

<sup>2</sup>Department of Clinical Pharmacy and Diagnostics, University of Vienna, Althanstr. 14 UZAII, A-1090 Vienna, Austria

[juergen.wanner@kurtkitzing.de](mailto:juergen.wanner@kurtkitzing.de)

Keywords: *Callitris intratropica*, essential oil, extract, HS-SPME

The wood and bark of *Callitris intratropica* has been used for a long time in folk medicine by the Australian aborigines [1], as an anti-termite timber [2] and since some years as a source of a now commercially available essential oil with a blue color due to the presence of azulenes [3]. The wood itself has a pleasant warm balsamic odor and the oil has a composition with substances unique to the callitris genus.

In the presented study commercial raw and refined oil, lab distilled oil, solvent extracts from different parts of the wood and headspace solid phase microextracts were analysed by GC-MS-FID. Quantitative (by FID detection) and qualitative analyses (by MS identification) were carried in one instrument with the use of a simple home-made MS-FID splitter. No azulenes were found in the wood itself so that the assumption suggests that the formation of azulenes takes place during the distillation process.

1. IA Ogunwande, VS Saroglou, E Skaltsa, AO Ogunbinu and D Kubmarawa (2009) JEOR 21:61-66
2. L Doimo, RJ Fletcher and BR D'Arcy (1999) JEOR 11:415-422
3. L Doimo (2001) JEOR 13:25-29



**P 96. Assessment of the antioxidant activity of *Lippia gracilis* Schauer aerial parts essential oil by different methodologies**

Teixeira ML<sup>1</sup>; Cardoso MG<sup>2</sup>; Souza JA<sup>2</sup>; Aazza S<sup>3</sup>; Miguel MG<sup>3</sup>; Gago G<sup>3</sup>; Barroso JG<sup>4</sup>; Pedro LG<sup>4</sup>; Figueiredo AC<sup>4</sup>

<sup>1</sup>Universidade Federal de Lavras, Departamento de Química, Caixa Postal 3037, Campus Universitário, Lavras, 37200-000, Minas Gerais, Brazil

<sup>2</sup>Universidade Federal de Lavras, Departamento de Química, Caixa Postal 3037, Campus Universitário, Lavras, 37200-000, Minas Gerais, Brasil

<sup>3</sup>Universidade do Algarve, Faculdade de Ciências e Tecnologia, Departamento de Química e Farmácia, Centro de Biotecnologia Vegetal, IBB, Faro, 8005-139 Faro, Portugal

<sup>4</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portugal

josefina@dqi.ufla.br

Keywords: *Lippia gracilis*; Verbenaceae; antioxidant activity; essential oil

*Lippia gracilis* Schauer is a Verbenaceae species native to northeastern Brazil. An ethnobotanical survey of some Brazilian medicinal plants showed that this plant, popularly known as *cidreira da serra*, is used in the treatment of several ailments, namely gastrointestinal infections, as well as respiratory and skin problems [1]. In the present study the chemical composition and antioxidant activity of *Lippia gracilis* essential oil was assessed by different methodologies.

The essential oil was isolated by hydrodistillation from *L. gracilis* aerial parts, and analysed by Gas Chromatography and Gas Chromatography-Mass Spectrometry as in [2]. The antioxidant activity was evaluated using three different methods as in [2,3]: free radical scavenging activity using 2,2-diphenyl-1-picrylhydrazyl (DPPH), Trolox equivalent antioxidant capacity (TEAC) or ABTS [2,2'azinobis(3-ethylbenzothiazoline-6-sulfonic acid)] method and the  $\beta$ -carotene bleaching method (coupled oxidation of  $\beta$ -carotene and linoleic acid).

Oxygen-containing monoterpenes (50%) and monoterpene hydrocarbons (46%) dominated *L. gracilis* essential oil. Carvacrol (42%) *p*-cymene (18%),  $\gamma$ -terpinene (17%) and thymol (5%) were the main ( $\geq 5\%$ ) essential oil components.

*L. gracilis* essential oil antioxidant ability showed a linear variation with increasing oil concentration for all the assessed methods (DPPH, TEAC and  $\beta$ -carotene bleaching method), showing a half maximal inhibitory concentration (IC<sub>50</sub>) of 434 $\mu$ g/mL, 625 $\mu$ g/mL, and 388 $\mu$ g/mL, respectively.

Acknowledgments: Partially supported by CNPq, CAPES, FAPEMIG.

1. Marinho MJM et al. (2011) *Rev. Bras. Plantas Med.* 13: 246-252.

2. Miguel G et al. (2011) *Natural Product Res.* 25: 526-541.

3. Lopes-Lutz et al. (2008) *Phytochemistry* 69: 1732–1738.

**P 97. Antiphytoviral activity of essential oil of two *Eryngium* species**Dunkić V<sup>1</sup>; Vuko E<sup>1</sup>; Bezić N<sup>1</sup>; Kremer D<sup>2</sup>; Ruščić M<sup>1</sup><sup>1</sup>Faculty of Science, University of Split, Croatia<sup>2</sup>Faculty of Pharmacy and Biochemistry, University of Zagreb, Croatia

dunkic@pmfst.hr

Keywords:  $\beta$ -caryophyllene; caryophyllene oxide; *Eryngium alpinum*; *E. amethystinum*; satCMV

In this study antiphytoviral activity of essential oils of *Eryngium alpinum* L. and *E. amethystinum* L. from Croatia against satellite associated cucumber mosaic virus (satCMV) was investigated. Water distilled essential oils from aerial parts of plants have been analysed by GC and GC/MS using VF-5ms capillary column. Thirty-two components representing 92.4% of the total oil were identified in the essential oil of *E. alpinum* and thirty-five components representing 93.1% of the total oil of *E. amethystinum*. The oil of *E. alpinum* was characterized by a high concentration of caryophyllene oxide (21.6%), bicyclogermacrene (11.8%) and germacrene D (10.3%) while the main components of *E. amethystinum* oil were  $\beta$ -caryophyllene (19.7%) and  $\alpha$ -pinene (12.3%). Germacrene D, caryophyllene oxide and bicyclogermacrene were also identified as the major compounds in the oil of *E. serbicum*, *E. palmatum* and *E. rosulatum* [1,2]. A comparison (t-test) of the mean number of lesions on the oil-treated *Chenopodium quinoa* plants with the corresponding control showed that both *E. alpinum* and *E. amethystinum* oils significantly reduced infection with satCMV with antiviral activity rate of 77.8% and 80.5%, respectively. Our previous investigation confirmed that sesquiterpenes-rich essential oils as well as oil components  $\beta$ -caryophyllene and caryophyllene oxide inhibit development of local lesions on plants infected with CMV [3-5]. Compared with the oils previously tested and with individual application of  $\beta$ -caryophyllene and caryophyllene oxide, both *Eryngium* oils showed stronger inhibition of local lesions development. We can conclude that combination of  $\beta$ -caryophyllene/ caryophyllene oxide with some other oil components resulted with significant antiviral efficiency. Our investigation showed that essential oils isolated from *E. alpinum* and *E. amethystinum* possess strong antiviral activity against satCMV infection by inducing resistance response in the host.

Acknowledgements: This work was supported by the Ministry of Science, Education and Sports of the Republic of Croatia, projects no. 177-1191192-0830 and 006-0000000-3178.

1. C. Capetanos et al. (2007) J. Serb. Chem. Soc. 72: 961-965.
2. J. Palá-Paúl et al. (2006) Biochem. Syst. Ecol. 34: 796-801.
3. V. Dunkić et al. (2010) Molecules. 15: 6713-6721.
4. V. Dunkić et al. (2011) Nat. Prod. Comm. 6: 1385-1388.
5. N. Bezić et al. (2011) Molecules. 16: 8119-8129.

**P 98. Chemical composition and antibacterial activity of *Lychnophora pinaster* Mart. essential oil**

Resende JMV<sup>1</sup>; Cardoso MG<sup>2</sup>; Batista LR<sup>1</sup>; Souza JA<sup>2</sup>; Rodrigues LMA<sup>2</sup>; Andrade MA<sup>2</sup>; Barroso JG<sup>3</sup>; Pedro LG<sup>3</sup>; Figueiredo AC<sup>3</sup>

<sup>1</sup>Universidade Federal de Lavras, Departamento de Ciência dos Alimentos, Caixa Postal 3037, Campus Universitário, Lavras, 37200 000, Minas Gerais, Brasil

<sup>2</sup>Universidade Federal de Lavras, Departamento de Química, Caixa Postal 3037, Campus Universitário, Lavras, 37200-000, Minas Gerais, Brasil

<sup>3</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portugal

mcardoso@dqi.ufla.br

Keywords: *Lychnophora pinaster*; Asteraceae; arnica; antibacterial; essential oil

Several studies have shown the antimicrobial efficacy of essential oils and in the last decades intensive studies were carried out indicating the possibility of their use in the control of foodborne pathogens, contributing to a future reduction in the employment of antibiotics [1,2].

*Lychnophora pinaster* Mart., is an Asteraceae listed as vulnerable to extinction in Brazil, and locally known as *arnica*. This medicinal plant is commonly used for its analgesic, antiinflammatory, antiprotozoal, antibacterial and antifungal properties. The hydro-alcoholic extract is widely used for the antiinflammatory and anesthetic values [3]. The goal of the present work was to characterize the chemical composition and antibacterial activity of *L. pinaster* essential oil.

The essential oil was isolated by hydrodistillation from *L. pinaster* aerial parts, and analysed by Gas Chromatography and Gas Chromatography-Mass Spectrometry as in [4]. Antibacterial ability of *L. pinaster* essential oil was tested by the agar well method [2] against *Staphylococcus aureus* ATCC 6538, *Listeria monocytogenes* ATCC 19117, *Escherichia coli* ATCC 11229, *Salmonella choleraesius* ATCC 6539 and *Pseudomonas aeruginosa* ATCC 15442. Several concentrations of the essential oil diluted with dimethylsulfoxide (DMSO), to a final volume of 10µL, were assessed (500, 250, 125, 63, 31, 16, 8 and 4µg/mL). The adopted statistical design was in randomized blocks, with 3 repetitions. Data was analysed using Tukey test ( $\alpha = 0.05$ ).

*trans*-Methyl cinnamate (62%),  $\beta$ -caryophyllene (21%),  $\alpha$ -humulene (6%) and  $\beta$ -pinene (5%) were the main ( $\geq 5\%$ ) essential oil components. Antimicrobial activity was detected only against *Salmonella choleraesius* at 31µg/mL.

Acknowledgments: Partially supported by CNPq, CAPES, FAPEMIG.

1. S. Bounatirou et al. (2007) *Food Chem* 105: 146-155.
2. EA Ostrosky, et al. (2008) *Rev. Bras. Farmacogn.* 18: 301-307.
3. de Souza et al. (2007) *Hort Science* 42: 1665-1669.
4. G. Miguel et al. (2011) *Nat. Prod. Res.* 25: 526-541.

**P 99. Chemical composition and repellence activity of *Philodendron imbe* Schott (Cipó imbé) essential oil against *Tetranychus ludeni***

Andrade J<sup>1</sup>; Cardoso MG<sup>1</sup>; Maluf WR<sup>2</sup>; Teixeira ML<sup>1</sup>; Oliveira CM<sup>1</sup>; Silva LF<sup>1</sup>; Barroso JG<sup>3</sup>; Pedro LG<sup>3</sup>; Figueiredo AC<sup>3</sup>

<sup>1</sup>Universidade Federal de Lavras, Departamento de Química, Caixa Postal 3037, Campus Universitário, Lavras, 37200-000, Minas Gerais, Brasil

<sup>2</sup>Universidade Federal de Lavras, Departamento de Agricultura, Caixa Postal 3037, Campus Universitário, Lavras, 37200-000, Minas Gerais, Brasil

<sup>3</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portugal

mcardoso@dqi.ufla.br

Keywords: *Philodendron imbe*, Cipó imbé; Araceae; *Tetranychus ludeni*; spider mite; essential oil

Spider mites, such as *Tetranychus ludeni*, are considered significant constraints to vegetable production [1]. Control of these spider mites using synthetic pesticides is not only hazardous to Man but has also the potential for induction of resistance in mites.

The neotropical genus *Philodendron* Schott (Araceae), presents significant landscape interest due to the large number of ornamental species. The bactericidal, insecticidal and antiprotozoal activities against *Trypanosoma cruzi* and *Trichomonas vaginalis* has been encouraged the phytochemical study of species of this genus. The leaf tea is locally used in coastal regions of Brazil as a vermifuge [2]. In the present study the chemical composition and repellence activity of *Philodendron imbe* Schott (*Cipó imbé*) essential oil against *T. ludeni* was assessed.

The essential oil was isolated by hydrodistillation from *Philodendron imbe* roots, and analysed by Gas Chromatography and Gas Chromatography-Mass Spectrometry as in [3]. The level of repellence to spider mites was assessed on tomato leaflets as in [4]. In addition to control experiments (T1: water plus Tween), different concentrations of the essential were evaluated (T2: 0.25%, T3: 0.50%, T4: 0.75%, T5: 1.0% and T6: 1.5%). The distances the mites moved after 20, 40 and 60min were measured. The randomized blocks statistical design was adopted, with 3 repetitions. Data was analysed using Tukey test ( $\alpha = 0.05$ ).

Sesquiterpene hydrocarbons (92%) dominated *P. imbe* essential oil,  $\beta$ -bisabolene (65%), *trans*- $\alpha$ -bergamotene (10%),  $\alpha$ -copaene (3%),  $\delta$ -cadinene (3%), *trans*- $\beta$ -farnesene (2%) and  $\beta$ -selinene (2%), being the main components.

The distance the mites moved increased with lowest essential oil concentration used, T3: 6.1mm < T2: 8.3mm < T1: 16.5mm. The shorter distance the mites moved, that is, the repellence activity, was observed with T4: 5.1mm, T5: 4.8mm and T6: 4.2mm essential oil concentrations

Acknowledgments: Partially supported by CNPq, CAPES, FAPEMIG.

1. GJ Moraes, HW Flechtmann (2008) In: Manual de acarologia: acarologia básica e ácaros de plantas cultivadas no Brasil., Ribeirão Preto, Brasil.
2. SJ Mayo (1988) *Acta Bot. Bras.* 1: 27-40.
3. MD Mendes et al. (2011) *Ind. Crop. Prod.* 33: 710-719.
4. CA Aragão et al. (2002) *Acta Bot. Bras.* 16: 83-88.

**P 100. Essential oil composition and variability of *Salvia nemorosa* L. (Lamiaceae) in the area of Vienna, Austria**

Chizzola R

University of Veterinary Medicine, Department for Farm Animals and Veterinary Public Health,  
Institute of Applied Botany and Pharmacognosy, Vienna, Austria

Remigius.Chizzola@vetmeduni.ac.at

Keywords: essential oil; *Salvia nemorosa*; sabinene; sesquiterpenes; hexadecanoic acid

Studying the aromatic flora of Eastern Austria we report here the composition of the essential oil from leaves, flowers and stems of *Salvia nemorosa*. This plant is an element of the pannonic flora from Eastern Austria. It is growing in semi-dry meadows and on embankments. Plants were collected from five sites in the northern parts of Vienna and along the Danube during the flowering stage and divided in leaves, stems and flowers. These samples were dried in the ambient air and hydrodistilled in a Clevenger type apparatus or extracted by microdistillation. The obtained oils were diluted with hexane and analysed by GC and GC/MS using an apolar Rtx-5 column for the separation. The compounds were identified according to their mass spectra and retention indices (1).

The oil content of all plant parts was below 0.05%. The oil from the flowers was dominated by sabinene (37-44%) followed by germacrene D (9-14%),  $\beta$ -caryophyllene (8-12%),  $\alpha$ -thujene (5-8%) and  $\gamma$ -terpinene (2-8%). The leaf oils were rich in  $\beta$ -caryophyllene (14-41%), germacrene D (14-38%) and caryophyllene oxide (5-20%). Hexadecanoic acid (56-60%) was the main compound in the stem volatile fractions.

In sum the plants of all five locations represent the same chemotype but differed from plants collected in Iran with  $\beta$ -caryophyllene, germacrene B and caryophyllene oxide as main oil compounds (2) and plants from Serbia having an oil were hexadecanoic acid, spathulenol and caryophyllene oxide prevailed (3).

Acknowledgements: The research was supported by "Hochschuljubiläums-Stiftung der Stadt Wien", Project H-02328/2007.

1. R. P. Adams, Identification of Essential Oil Components by Gas chromatography / Mass Spectrometry, 4<sup>th</sup> ed. Allured Pub. Corp., Carol Stream, IL (2007).
- 2 M Mirza, F Sefidkon (1999) Flavour Fragrance J. 14, 230-232
- 3 D Malencic et al. (2002) Flavour Fragrance J. 19, 225-228

**P 101. Effect of systemic administration of essential oils and main components on honeybee survival**

Porrini MP<sup>1</sup>; Afonso Pires SM<sup>2</sup>; Rossini C<sup>3</sup>; Garrido PM<sup>1</sup>; Hermida L<sup>4</sup>; Gende LB<sup>1</sup>; Eguaras MJ<sup>1</sup>

<sup>1</sup>Laboratorio de Artrópodos-FCEyN, UNMDP, CONICET, Argentina

<sup>2</sup>Centro de Investigação de Montanha (CIMO)/Escola Superior Agrária, Instituto Politécnico de Bragança, Portugal

<sup>3</sup>Laboratorio de Ecología Química, Facultad de Química, UdelaR, Uruguay

<sup>4</sup>Centro de Investigación y Desarrollo en Química, Instituto Nacional de Tecnología Industrial, Buenos Aires, Argentina

spires@ipb.pt

Keywords: essential oils; survival; *Apis mellifera*

Controlling bee diseases with non contaminant products is a challenge in apicultural research. Essential oils and their main components have been widely studied as alternative treatments for honeybee pathologies [1, 2, 3]. However, there is little information about prolonged systemic administration. The aim of this study was to evaluate, in laboratory assays, the effect of long term consumption of essential oils and main components. Oils were obtained by hydrodistillation from *Laurus nobilis*, *Cinnamomun zeylanicum*, *Origanum vulgare*, *Rosmarinus officinalis* and *Eucalyptus* spp. and were analyzed by gas chromatography. The main components administered were 1,8-cineol,  $\beta$ -myrcene, cinnamic aldehyde, carvacrol and  $\alpha$ -phellandrene. Substances were administered *ad libitum* to newly emerged bees at concentrations of 0; 333; 3,333 and 6,666 ppm, on sucrose syrup, throughout 11-18 days. Mortality and substances consumption were measured daily. Survival analysis was performed using Gehan-Breslow test and pairwise multiple comparisons between survival curves ( $\alpha= 0, 05$ ). Substances consumption was analyzed using one way ANOVA. Bees that received cinnamon oil showed a lower survival than control at concentrations higher than 333 ppm ( $p$ -values $<0.001$ ). Consumption of cinnamic aldehyde, the main component of this oil (79.3%), also caused lower survival at the same concentrations ( $p$ -values $<0.001$ ). Eucalyptus oil caused a lower survival rate when it was administered at 6,666 ppm, although 1,8 cineol, its main component (63.5%), was not toxic for bees at any concentration. Carvacrol, a main component of many oregano essential oils, showed toxic effects at 3,333 and 6,666 ppm. Essential oils did not cause differences in consumption rate ( $p= 0.275$ ) while main components solutions, except for carvacrol, were less consumed than control at the three concentrations. Treatments did not cause dysentery to bees. Our results contribute to understanding the effect of repeated systemic doses of these substances, which is important to design long term pharmacological studies and treatments development.

Acknowledgments The authors wish to thank the collaboration made by María Laura Umpiérrez and the contribution made by Enrique Tkacik who provided *L. nobilis* essential oil.

1. C Costa, M Lodesani, L Maistrello (2010). Apidol. 41 (2):141-150.

2. LB Gende, I Floris et al. (2008). Bull. Insectol. 61(1):1-4.

3. ME Umpiérrez, E Santos (2011), A. González, Rossini. Phytochem. Rev. 10(2): 227-244.

**P 102. The use of essential oils constituents' effect on shelf life of fresh-cut *Salicornia ramosissima* and *Sarcocornia perennis***

Antunes MDC<sup>1</sup>; Gago C<sup>1</sup>; Branquinho AR<sup>1</sup>; Julião M<sup>1</sup>; Faleiro L<sup>2</sup>; Panagopoulos T<sup>3</sup>; Miguel MG<sup>1</sup>

<sup>1</sup>Faculdade de Ciências e Tecnologia, IBB-CBV, Edf. 8, Campus de Gambelas, 8005-139 Faro, Portugal

<sup>2</sup>Faculdade de Ciências e Tecnologia, IBB-CBME, Edf.8, Campus de Gambelas, 8005-139 Faro, Portugal

<sup>3</sup>Faculdade de Ciências e Tecnologia, CIEO, Edf. 8, Campus de Gambelas, 8005-139 Faro, Portugal

[mantunes@ualg.pt](mailto:mantunes@ualg.pt)

Keywords: Fresh-cut salads, halophytes, quality, antioxidants, eugenol, citral.

The halophyte plants *Salicornia ramosissima* and *Sarcocornia perennis* have been exploited in gourmet cuisine combined with fresh salads as substitute of salt due to their nutritional value. The objective of this work was to evaluate the ability of the essential oil constituent's citral and eugenol on the preservation of the quality of these halophyte plants through shelf life at two temperatures.

The fresh tips of the plants (7-8 cm long) were washed in tap water and, after removal of excess water, immersed in solutions of 0.025% citral (w/v), 0.025% (w/v) eugenol, or just distilled water (control), during 2 minutes. The samples were prepared for storing in polyethylene foam packages, covered with low density polyethylene film of 15 µm thickness and put in refrigeration rooms at 1, and 6 °C and relative humidity of ≈90%. At 0, 7 and 14 days samples were removed and quality measurements were taken: color, weight loss, electrolyte leakage, antioxidant activity, phenolic compounds, bacteria, yeasts and molds. A taste panel was also performed.

Results showed that temperature was the main factor affecting storage ability of fresh-cut salicornia and sarcocornia, being 1°C better temperature. General quality parameters were slightly better preserved through 14 days at 1 °C when the fresh-cut halophytes were treated with 0.0025% citral or eugenol. Taste panel performed well through shelf life, being some parameters scored higher in citral and eugenol treated fresh-cut halophyte plants than control.

Acknowledgements: This research was partially supported by the Project: PTDC/AGR-ALI/121085/2010-INNCOAT - Innovative edible coatings to improve storage of small fruit and fresh-cut.

**P 103. Volatiles components from the leaves of *Centaurea vlachorum* Hartvig (Asteraceae), growing wild in Albania**

Hodaj E<sup>1,2</sup>, Lazari D<sup>1</sup>, Abazi S<sup>3</sup>

<sup>1</sup>Laboratory of Pharmacognosy, School of Pharmacy, Aristotle University of Thessaloniki, GR-54124 Thessaloniki, Greece

<sup>2</sup>Agri-Food Technology Department, Faculty of Biotechnology and Food, Agricultural University of Tirana, Kamz, Albania

<sup>3</sup>Chemistry Department, Faculty of Natural Sciences, University of Tirana, Tirana, Albania

entelahodaj@hotmail.com

Keywords: *Centaurea vlachorum*, Asteraceae, essential oil, GC-MS.

The genus *Centaurea* is a polymorphous genus of the family Asteraceae and includes grassy plants, from annual to perennial, rarely suffruticose. The aerial parts of several species of *Centaurea* are used in popular medicines in many countries as antibacterial, antimicrobial, hypoglycemic, cytotoxic and phytotoxic agents.

*Centaurea vlachorum* Hartvig (Asteraceae) is a new species discovered lately in Albania. It was found at the altitudes between 1600-2000m above sea level, in the Lura Lakes region. There are no reports in the literature about the chemical analyses of this plant.

For the present study, the leaves of the plant were collected during its flowering season, early July, dried in darkness till a constant weight, grinded and used as such for hydro distillation using a Clevenger type apparatus according to standard procedures. The volatile components were analyzed by Gas Chromatography-Mass Spectrometry. The major compounds found in essential oil were: benzaldehyde (21.0%), spathulenol (7.9%) and caryophyllene oxide (10.0%),



**P 104. Chemical characterization and biological activity of *Margotia gummifera* essential oil**

Valente J<sup>1</sup>; Zuzarte M<sup>2</sup>; Gonçalves MJ<sup>2</sup>; Lopes MC<sup>3</sup>; Cavaleiro C<sup>2</sup>; Cruz MT<sup>3</sup>; Salgueiro L<sup>2</sup>

<sup>1</sup>Center of Pharmaceutical Studies, Faculty of Pharmacy, Health Science Campus, University of Coimbra, Portugal e Centro de Neurociências e Biologia Celular da Universidade de Coimbra, 3000-295 Coimbra, Portugal

<sup>2</sup>Center of Pharmaceutical Studies, Faculty of Pharmacy, Health Science Campus, University of Coimbra, Azinhaga de S. Comba 3000-354, Coimbra, Portugal

<sup>3</sup>Centro de Neurociências e Biologia Celular e Faculdade de Farmácia, Universidade de Coimbra, 3000-295 Coimbra, Portugal

jurita8@yahoo.com.br

Keywords: *Margotia gummifera*; essential oils; antifungal activity; anti-inflammatory activity; cytotoxicity

This study was designed to elucidate the composition, cytotoxicity, antifungal and anti-inflammatory activities of the essential oil of *Margotia gummifera* (Desf.) Lange. Ripe umbels with mature seeds were submitted to water distillation in a Clevenger-type apparatus and the oils were analysed by GC and GC-MS, as previously reported [1]. The oil was characterized by high contents of monoterpene hydrocarbons, being the main compounds myrcene (20.4%) and sabinene (22.5%). The antifungal activity (MIC and MLC) was evaluated against dermatophyte strains, using the macrodilution method according to Clinical and Laboratory Standards Institute. *Microsporum canis* FF1 and *Trichophyton rubrum* CECT 2794 are the more sensitive strains. Assessment of macrophages viability was performed using the MTT assay. The results showed that the oil is not cytotoxic at concentrations ranging from 0.32 to 1.25 µL/mL. This study also evaluated the anti-inflammatory properties of *M. gummifera* oil, through the measurement of the pro-inflammatory mediator. After cells activation with lipopolysaccharide (LPS) for 24 h, nitrite production increased to 162.8% above control, while in the presence of the oil concentrations 0.32 µl/ml, 0.64 µl/ml and 1.25 µl/ml, nitrite production was reduced to 140.7%, 139.9% and 128.8% of the control, respectively. Furthermore, we also addressed the NO scavenging activity of *M. gummifera* using SNAP (S-nitroso-N-acetylpenicillamine) as NO donor and we observed that the oil exhibits a slight scavenging activity at the concentration 0.64 µl/ml.

These results indicate that the biological activity of *Margotia gummifera* essential oil should be explored and suggest its potential as a natural source of new anti-inflammatory drugs.

Acknowledgements: Funding: FCT (Fundação para a Ciência e a Tecnologia) project CEF/POCI2010/FEDER

1. C.Cavaleiro, L. Salgueiro, M.G.Miguel, A.Proença da Cunha (2004) *J Chromatogr A*. 1033:187-190.

**P 105. Effect of citral in rat mesenteric artery: evidences of the involvement of calcium channels**

Santos MRV<sup>1</sup>; Moreira FV<sup>1</sup>; Fraga BP<sup>1</sup>; Cunha PS<sup>1</sup>; Araujo AAS<sup>1</sup>; Quintans-Júnior LJ<sup>2</sup>; Bonjardim LR<sup>1</sup>

<sup>1</sup>Federal University of Sergipe

<sup>2</sup>Federal University of Sergipe - Brazil

marcio@infonet.com.br

Keywords: calcium channels, citral, vasorelaxant effect, rats

Citral is a monoterpene found in essential oils of various herbs with activity on the cardiovascular system. This study aimed to evaluate the vasorelaxant effect of the citral in rings of rat superior mesenteric artery.

Male wistar rats (200 – 300 g) were euthanized by exsanguination under anesthesia and superior mesenteric artery was removed and cut in rings (1-2 mm). These rings were then mounted in organ baths containing Tyrode's solution at 37 °C and gassed with carbogen. For isometric tension recordings, each ring was fixed in a force transducer connected to an acquisition system. All procedures described in this work were approved by the Ethical Committee on Animal Experiments from UFS under protocol number 37/2009.

In rings pre contracted with phenylephrine (10 µM), citral ( $10^{-7}$  -  $10^{-2}$  M) was able to induce relaxations ( $pD_2 = 2.52 \pm 0.10$ ;  $E_{max} = 103.4 \pm 10.2\%$ ;  $n = 6$ ) that was not affected after removal of the endothelium ( $pD_2 = 2.34 \pm 0.15$ ;  $E_{max} = 107.2 \pm 4.3\%$ ;  $n = 6$ ) or in rings without endothelium pre-contacted with KCl 80 mM ( $pD_2 = 2.04 \pm 0.12$ ;  $E_{max} = 101.3 \pm 7.1\%$ ;  $n = 6$ ). In a calcium-free solution, citral ( $3 \times 10^{-4}$  and  $10^{-3}$  M) was able significantly ( $p < 0.05$ ) to inhibit the  $CaCl_2$ -induced contractions ( $10^{-5}$  –  $10^{-2}$  M) up to  $88.6 \pm 3.1\%$ . Furthermore, citral ( $3 \times 10^{-4}$  M) was not capable to induce an additional effect on maximal relaxation of nifedipine (10 µM), a L type voltage-operated calcium channel blocker.

These results demonstrate that citral induce vasorelaxation of mesenteric artery possibly caused by the inhibition of the  $Ca^{2+}$  influx through voltage-operated  $Ca^{2+}$  channels.

Acknowledgments: CNPq/FAPITEC, Brazil.

**P 106. *In vitro* antimicrobial activity and cytotoxic potential of North-West Romanian propolis**

Niculae M<sup>1</sup>; Stan L<sup>2</sup>; Spinu M<sup>1</sup>; Marghitas LAI<sup>3</sup>; Sandru CD<sup>1</sup>; Pall E<sup>1</sup>

<sup>1</sup>*Discipline of Infectious Diseases, University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca Romania*

<sup>2</sup>*Department of Food Engineering, University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca Romania*

<sup>3</sup>*Department of Beekeeping and Sericulture, University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca Romania*

nicmisi23@yahoo.com

Keywords: antimicrobial; cytotoxic; propolis phenolics; flavonoids

Propolis is a honeybee product demonstrated with complex therapeutic effects [1,2,3]. Objective: the aim of the study was to evaluate antimicrobial activity of ethanolic extract of North-West Romanian propolis samples against pathogens of veterinary significance (*Escherichia coli*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Malassezia pachydermatis*) and to investigate the cytotoxic potential. Methods:

Propolis samples phenolic profile was characterized by spectrophotometric methods. The antimicrobial efficacy was assessed by agar diffusion and broth microdilution method, while the cytotoxic potential was estimated considering propolis biocompatibility on human fibroblasts cell culture (cell line HFL-1) using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay and evaluating the cell morphology and attachment level.

Propolis samples, with total phenolics of 38,02%  $\pm$ 2,34% as determined by Folin Ciocalteu method, presented high amounts of total flavonoids (9 $\pm$ 0,3%), where 1,74-9,22% belong to the group of flavones/flavonols and 1,96-4,01% were flavanones/dihydroflavonols. Based on the bacterial growth inhibition diameter zone (25-26mm) and values determined as MICs and MBCs (0,125-2%v/v) an important antimicrobial activity was suggested for all tested propolis samples, with the most intense effect against Gram positive strains. MTT test data indicated concentration dependence of propolis-induced effect, some dilutions stimulated cell viability, while higher concentrations had moderate expressed cytotoxicity.

The propolis samples taken in this study presented the typical poplar composition profile with flavonoids and phenolic acids as main biological active compounds. Chromatographic analysis confirmed the data obtained spectrophotometrically. Propolis phenolic profile and biological activity evaluation facilitated the selection of samples with antimicrobial activity and moderate or absent cytotoxic potential.

Acknowledgments: We wish to thank Dr. Olga Soritau and Dr. Virag Piroska from the Laboratory of the Oncology Institute "Ion Chiricuța" Cluj-Napoca, Romania for technical support during cytotoxicity screening.

1. D Teerasripreecha et al. (2012) BMC Complement Altern Med 12: 27.
2. ACHF Sawaya et al. (2004) Braz J Microbiol. 35(1-2): 104-109.
3. S Mohammadzadeh et al. (2007) Food Chem. 103: 1097-1103.

**P 107. Vasorelaxant effect induced by geraniol in rat mesenteric artery**

Santos MRV<sup>1</sup>; Fraga BP<sup>1</sup>; Cunha PS<sup>1</sup>; Moreira FV<sup>1</sup>; Barreto AS<sup>2</sup>; Araujo AAS<sup>1</sup>; Nunes RS<sup>1</sup>; Ribeiro EAN<sup>3</sup>; de Medeiros IA<sup>4</sup>; Marchioro M<sup>1</sup>

<sup>1</sup>Federal University of Sergipe

<sup>2</sup>Federal University of Sergipe – Brazil

<sup>3</sup>Federal University of Alagoas

<sup>4</sup>Federal University of Paraiba

marcio@infonet.com.br

Keywords: geraniol, essential oils, vasorelaxant effect, rat mesenteric artery

Geraniol is a monoterpene found in essential oils of various herbs with activity on the cardiovascular system [1]. The objective of the present study was to evaluate the vasorelaxant effect induced by geraniol in rat mesenteric artery.

Male Wistar rats (200 – 300g) were euthanized by exsanguination under anesthesia and superior mesenteric artery was removed and cut in rings (1-2 mm), which were mounted in organ baths containing 10 mL of Tyrode's solution at 37°C and gassed with carbogen. For isometric tension recordings, each ring was fixed in a force transducer connected to an acquisition system.

In mesenteric artery rings with functional endothelium pre-contracted with 10 µM of phenylephrine (control), geraniol ( $10^{-8}$  –  $10^{-2}$  M) was able to induce relaxation in a concentration-dependent manner ( $E_{max} = 110 \pm 5\%$ ;  $n = 6$ ) which was not attenuated after removal of endothelium ( $E_{max} = 108 \pm 6\%$ ;  $n = 4$ ) or after 1 mM of tetraethylammonium, a non-selective blocker of  $K^+$  channels ( $E_{max} = 109,2 \pm 3\%$ ;  $n = 6$ ). In endothelium-denuded rings pre-contracted with KCl 80 mM, geraniol produced relaxation that was significantly ( $p < 0.05$ ) higher than those obtained in rings with functional endothelium pre-contracted with phenylephrine ( $E_{max} = 142 \pm 14\%$ ;  $n = 6$ ). Furthermore, the incubation with  $10^{-4}$ ,  $3 \times 10^{-4}$  or  $10^{-3}$  of geraniol was able to antagonize the tonic contractions induced by  $CaCl_2$  ( $10^{-7}$  –  $3 \times 10^{-2}$ ) and phasic contractions induced by phenylephrine in without calcium solution.

CONCLU

Taken together, these results suggest that geraniol produces a vasorelaxant effect by an endothelium-independent mechanism in rat mesenteric artery. This effect appears to be due to an inhibition of the  $Ca^{2+}$  influx through voltage-operated  $Ca^{2+}$  channels associated to an inhibition of the  $Ca^{2+}$  release through phenylephrine sensitive intracellular  $Ca^{2+}$  stores.

Acknowledgments: CNPq/FAPITEC, Brazil.

**P 108. Counter-current chromatography as a tool to identify minor oxygenated sesquiterpenes in *Baccharis dracunculifolia* volatile oil.**

Queiroga CL<sup>1</sup>, Ferraz PC<sup>1</sup>, Coser RN<sup>1</sup>, Cavalcante MQ<sup>1</sup>, Sartoratto A<sup>2</sup>, Magalhães PM<sup>3</sup>.

<sup>1</sup>University of Campinas (UNICAMP), Chemical Biological and Agricultural Pluridisciplinary Research Center (CPQBA), Division of Phytochemistry, 13083-970, Box 6171, Campinas, São Paulo, Brazil

<sup>2</sup>CPQBA, Division of Organic and Pharmaceutical Chemistry

<sup>3</sup>CPQBA, Division of Agrotecnology

cqueiroga.fito@gmail.com

Keywords: *Baccharis dracunculifolia*, CCC, GC/MS, oxygenated sesquiterpenes.

*Baccharis dracunculifolia* (Asteraceae) is an aromatic native plant of Brazil, Paraguay, Uruguay and Argentina. The essential oil of *B. dracunculifolia* leaves has a special odor that attracts bees [1] and also is used in the cosmetic and perfumery industries. Many of the minor compounds are co-responsible by the bouquet. Their presence can add value to the product. The goal of our investigation was to use of counter current chromatography (CCC) as a tool to obtain fractions enriched of oxygenated sesquiterpene.

Essential oil of *B. dracunculifolia* leaves was extracted by hydrodistillation in a modified Clevenger system. A CCC, model Quattro HSCCC, was used. The biphasic solvent system was hexan-methanol-water (5:4:1)[2]. Isocratic elution was conducted in a tail-to-head manner at 850 rpm and flow rate 2.0 mL/min. 500mg of volatile oil sample was dissolved in upper phase/lower phase. Fractions of 1.5mL was collected and analyzed by TLC and GC/MS.

The TLC analysis of the fractions showed that the major sesquiterpene, nerolidol, eluted after fraction 12. The applied CCC conditions afforded nerolidol in high purity (92.5 %). Spathulenol eluted after nerolidol and was obtained fraction with 50.2% purity. The minor oxygenated sesquiterpenes co-eluted with spathulenol and was identified as a group of three cadinols.

CCC technique has many advantages over traditional solid-liquid chromatography, for example: isolation in semi-preparative scale, good mass recovery and compounds are obtained in high purity. This purification was used in order to isolate fractions enriched in cadinols, trace alcohols present in *B. dracunculifolia* essential oil.

Acknowledgments: PETROBRAS, UNICAMP.

1. MR Maróstica Junior et al. (2008) Ciênc. Tecnol. Aliment. 28: 178-181.

2. J Xie et al. (2009) Food Chem. 117: 375-380.

**P 109. Evaluation of chemical composition and *in vitro* antiproliferative activity of essential oils of different parts of *Aldama linearifolia* Chodat**

Rehder VLG<sup>1</sup>; de Oliveira ASS<sup>2</sup>; Bombo AB<sup>3</sup>; Ruiz ALTG<sup>1</sup>; de Carvalho JE<sup>4</sup>; da Glória BA<sup>3</sup>

<sup>1</sup>Research Center for Chemical, Biological and Agricultural - CPQBA-UNICAMP

<sup>2</sup>Research Center for Chemical, Biological and Agricultural -CPQBA-UNICAMP

<sup>3</sup>ESALQ- São Paulo University

<sup>4</sup>Research Center for Chemical, Biological and Agricultural - CPQBA-UNICAMP

rehder@cpqba.unicamp.br

Keywords: *Aldama linearifolia*; *Viguiera linearifolia*; antiproliferative activity; essential oil

*Aldama linearifolia* synonym of the *Viguiera linearifolia* (Asteracea) was perennial herbs provided with thickened underground system and potentially aromatic resins. The objective of this study was to analyze the chemical composition and evaluate *in vitro* the antiproliferative activity of the essential oils (EO's) from the different parts of *A. linearifolia*. Plants were collected at flowering stage in Ponta Porã - MS, Brazil and separated into roots, xylopodia, stems and leaves. The essential oils (EO's) were obtained by hydrodistillation in Clevenger system using fresh plant and analyzed by GC-MS. The activity was performed according to protocols recommended by the *National Cancer Institute* (NCI) for nine human tumor cell lines: U251 (CNS), MCF-7 (breast), NCI-ADR/RES (multi-drug resistant ovarian), 786-0 (kidney), NCI-H460 (lung), PC-03 (prostate), OVCAR-03 (ovarian), HT-29 (colon), K-562 (leukemia), and HaCat (strain normal cells). The higher yield of EO was observed in roots and xylopodia (0.23 and 0.14%) whereas the stem and leaves had low yields (0.07 and 0.05%). EO's of roots and xilopodia showed only monoterpenes: a-pinene (77 and 66%), b- pinene (7.4 and 8.5%) and 3-carene (15.6 and 17%). The leaves presented several monoterpenes (90%) and the main were D-limonene (17%) and b-pinene (10.3%). The stems OE presented large amount of sesquiterpenes (88%), and the main were bicyclogermacrene (30%), germacrene D (12%) and spathulenol (11%). The antiproliferative activity *in vitro* showed that the four oils are active. From the curves of cell growth as a function of concentration of the samples were calculated TGI (*Total Growth Inhibition*). The EO's of the stems and leaves were selectivity for the strains of the resistant ovary (27.84  $\mu\text{g.mL}^{-1}$ ) and prostate (34.36  $\mu\text{g.mL}^{-1}$ ) and the OE's of the roots and xilopodia showed cytotoxicity with TGI < 100  $\mu\text{g.mL}^{-1}$ . The toxicity the EO's strain against normal cells showed that the roots EO was less cytocide (TGI > 250 $\mu\text{g.mL}^{-1}$ ).

**P 110. Composition and biological activities of *Seseli rigidum* Waldst. & Kit. fruit essential oil**

Ilić M<sup>1</sup>; Jovanović VS<sup>1</sup>; Jovanović O<sup>1</sup>; Mitić V<sup>1</sup>; Jovanović S<sup>1</sup>; Petrović G<sup>1</sup>; Stojanović G<sup>1</sup>

<sup>1</sup>University of Niš, Faculty of Science and Mathematics, Višegradska 33, Nis, Serbia

*marija.fertico@gmail.com*

Keywords: *Seseli rigidum* Waldst. & Kit.; essential oil; antioxidant activity; antimicrobial activity; cholinesterase inhibition

*Seseli rigidum* Waldst. & Kit. is an endemic Serbian plant species known as “devesilje”, indicating its ability to cure nine diseases. Composition of *S. rigidum* fruit essential oil was determined by GC and GC-MS [1]. Antioxidant activity was estimated as free radical scavenging capacity towards DPPH and ABTS, and total reducing power assay [2]. The antimicrobial activity was investigated using micro well-dilution method against Gram (+) bacteria: (*Staphylococcus aureus*, *Bacillus cereus*), Gram (-) bacteria (*Escherichia coli*, *Pseudomonas aeruginosa*) and fungal strains (*Candida albicans*, *Aspergillus niger*) [3]. For anticholinesterase activity, modified Ellman’s assay was utilized [4].

A total of 36 components were identified representing 94.4% of *S. rigidum* fruit oil. The most abundant compounds in the oil were  $\alpha$ -pinene (37.8%) and sabinene (13.5%). All three antioxidant assays revealed weak antioxidant capacity of *S. rigidum* fruit essential oil in contrast with significant activity against all examined bacterial and fungal strains, especially against Gram (+) bacteria *S. aureus*. Inhibition of pooled human serum cholinesterase and horse serum cholinesterase was significant, referring to neostigmine bromide as standard, mainly due to dominance of  $\alpha$ -pinene, as an evaluated cholinesterase inhibitor in the same experiments.

Composition of *S. rigidum* fruit essential oil corresponds to the data reported for *Seseli* species from Serbia, regarding the main component, but there are significant differences in the presence and distribution of other components. Remarkable anti-cholinesterase activity of essential oil, opens promising prospective in prevention of neurodegenerative diseases.

Acknowledgements: The research was supported by the Serbian Ministry of Education and Science (Grant no. 172047).

1. V Mitić et al. (2011) *J. Essent. Oil Res.* 23: 70-74.
2. G Stojanović et al (2010) *Cent. Eur. J. Biol.* 5: 808-813.
3. T Mihajilov-Krstev et al (2009) *Cent. Eur. J. Biol.* 4(3): 411–416.
4. V Stankov-Jovanović et al (2007) *J. Clin. Lab. Anal.* 21: 124-131.

**P 111. Composition of the volatile oils of *E. dilatatum* Lam. and *E. pandanifolium* Cham. & Schldl.**

Fernandes ESS; Salgueiro L; Cavaleiro C

Centro de Estudos Farmacêuticos / Faculdade de Farmácia da Universidade de Coimbra, Portugal

cavaleir@ff.uc.pt

Keywords: Essential oil, *Eryngium dilatatum*, *Eryngium pandanifolium*

*Eryngium* L. is probably the most extensive and taxonomically complex genus of the Apiaceae family, including about 250 species distributed all around the world. We report now on the composition of the essential oil of two species growing wild in Portugal, *E. dilatatum* and *E. pandanifolium*. The first grows in dry and stony fields of the Iberian Peninsula and North-Africa, the latter, with a broader distribution, prefers well-drained soils and sunny places, being frequent in the region of “Baixo Mondego”, in the center of Portugal.

The compositions of the oils isolated by water distillation were established by GC and GC-MS according a methodology previously reported <sup>[1]</sup>. The oil of *E. dilatatum* is mainly composed by sesquiterpene hydrocarbons (35.6%) and oxygen containing sesquiterpenes (23.6%), being Z- $\alpha$ -chrysanthenyl acetate (11.1%), germacrene D (10.3%),  $\alpha$ -pinene (9.2%), bicyclogermacrene (8.1%), spathulenol (5.9%) and E-caryophyllene (4.5%) the major components. Differently, the oil of *E. pandanifolium* is predominantly composed by monoterpene hydrocarbons (41.8%);  $\alpha$ -Pinene (20.8%),  $\beta$ -elemene (10.6%) and limonene (5.8%) are the most representative components.

Acknowledgements: Authors are grateful to Fundação para a Ciência e Tecnologia (FCT), POCI 2010/FEDER for financial support.

1. Cavaleiro, C.; Gonçalves, M.J.; Serra, D.; Santoro, G.; Tomi, F.; Bighelli, A.; Salgueiro, L.; Casanova, J. Composition of a volatile extract of *Eryngium duriaei* subsp. *juresianum* (M. Laínz) M. Laínz, signalised by the antifungal activity. *Journal of Pharmaceutical and Biomedical Analysis* 2011, 54, 619–622



**P 112. Chemical composition, biosynthesis sites and evaluation of *in vitro* antiproliferative activity of essential oils of *Aldama filifolia* (Asteraceae)**

da Glória BA<sup>1</sup>; Bombo AB<sup>1</sup>; de Oliveira ASS<sup>2</sup>; Ruiz ALTG<sup>2</sup>; de Carvalho JE<sup>2</sup>; Rehder VLG<sup>2</sup>

<sup>1</sup>ESALQ - São Paulo University

<sup>2</sup>Research Center for Chemical, Biological and Agricultural - CPQBA-UNICAMP

bagloria@usp.br

Keywords: anatomy; human tumor cell lines; monoterpenes; secretory canals; vegetative organs

Phytochemical studies have shown the pharmacological potential of *Viguiera (lato senso)* species [1,2]. This study aimed to analyze the chemical composition, to indicate the sites of biosynthesis and to evaluate the antiproliferative activity of the essential oils (EO's) of *Aldama filifolia* (= *Viguiera filifolia*). Plants were collected in Alto Paraíso de Goiás, GO, Brazil and separated into roots, xylopodia, stems and leaves. The extraction was carried out by hydrodistillation in Clevenger using fresh plant material, for three hours. The analyses were performed by gas chromatography with mass selective detector (GC-MS). Anatomical analyses were performed using usual histological techniques. Tests for *in vitro* antiproliferative activity were performed according to protocols recommended by the *National Cancer Institute* (NCI), for nine human tumor cell lines: U251 (CNS), MCF-7 (breast), NCI-ADR/RES (multi-drug resistant ovarian), 786-0 (kidney), NCI-H460 (lung), PC-03 (prostate), OVCAR-03 (ovarian), HT-29 (colon), K-562 (leukemia), and a strain normal cells (HaCat). From the curves of cell growth as a function of concentration of the samples were calculated TGI (Total Growth Inhibition - effective concentration for total inhibition of cell growth). The EO's yield (% w/w) was 1.29 in leaves, 0.22 in roots, 0.12 in xylopodia and 0.09 in aerial stems. The major compounds were Myrcene in leaves (86.0%) and aerial stems (73.1%) and  $\alpha$ -Pinene in roots (70.01%) and xylopodia (69.18%). EO's were found into secretory canals consisted of simple or multiseriate epithelium and variable diameter, in all analyzed organs. The results of studies on antiproliferative activity *in vitro* showed that only underground EO's are active. The EO's of the roots were more selective for CNS and ovarian with TGI's values of 94.62 and 55.64  $\mu\text{g}/\text{mL}^{-1}$ , respectively. The EO's of the xylopodia was active for other lines, especially leukemia and prostate, with TGI's values of 0.27 and 34.74  $\mu\text{g}/\text{mL}^{-1}$ .

Acknowledgments: We thank to The National Council for Scientific and Technological Development (CNPq) for research grants awarded to B.A.G. and V.L.G.R. and São Paulo Council for Research (FAPESP) – (Thematic Project Proc. n° 2010/51454-3 and Proc. n° 2010/01931-0), for the financial support and grant for the second author.

1. FB Da Costa et al. (1996) *Biochem. Syst. Ecol.* 24: 585-587.

2. TC Carvalho et al. (2011) *Molecules* 16: 543-551.

**P 113. Chemical composition and pharmacological activities of essential oils of *Lavandula* spp.**

Martins MR<sup>1</sup>; Arantes S<sup>1</sup>; Candeias MF<sup>1</sup>; Pereira M<sup>2</sup>; Cruz-Morais J<sup>1</sup>

<sup>1</sup>ICAAM & Departamento de Química, Escola de Ciências e Tecnologia, Universidade de Évora, Évora, Portugal

<sup>2</sup>ICAAM & Departamento de Planeamento, Ambiente e Ordenamento, Universidade de Évora, Évora, Portugal

mrm@uevora.pt

Keywords: *L. stoechas* subsp. *luisieri*; *L. viridis*; Acute toxicity; Analgesic effect; Anti-inflammatory activity

*Lavandula* spp. belong to the family *Lamiatae* and some species are often used in popular medicine and have been used for centuries in a large number of medical applications and in aromatherapy. Although similar ethnobotanical properties of *Lavandula* spp., its essential oils, general chemical composition and therapeutic applications differ from different species.

*Lavandula stoechas* L. subsps. *luisieri* (Rozeira) Rozeira and *L. viridis* L'Hér are endemic to the Iberian Peninsula, widespread in the South of Portugal, namely in Southern Alentejo and Algarve. The aim of our study was evaluate the chemical composition and toxicological and pharmacological activities of leaves essential oils of spontaneous plants of *L. stoechas* L. subsps. *luisieri* (Alentejo) and *L. viridis* (Algarve). The essential oils of these wild plants, collected in spring, were obtained by hydrodistillation in a *Clevenger*-type apparatus and its chemical composition was evaluated by GC/FID. The acute toxicity of essential oils was evaluated "*in vitro*" using brine shrimp (LC<sub>50</sub>) and "*in vivo*" using *Swiss* mice (DL<sub>50</sub>). The analgesic and anti-inflammatory pharmacological properties of *L. stoechas* subsp. *luisieri* essential oil were evaluated in mouse or rats by the *Amour-Smith* and carrageen-induced paw edema tests, respectively.

Results showed important differences in chemical composition of essential oils from two species analyzed either to diversity and proportion of its constituents. The essentials oils showed citotoxicity against *Artemia salina* and a DL<sub>50</sub> higher than 2000 mg/kg for mice. The analgesic and anti-inflammatory activities of essential oils were exhibit for the doses of 100 and 200 mg/kg.

These essential oils from *Lavandula* spp. showed important biological properties and studies will continue in order to clarify its hepatotoxicity and nephrotoxicity and to evaluate its potential use for pharmacological and nutritional applications.

**P 114. Chemical composition and antibacterial activity of five chemotypes of *Chamomilla recutita* (L.) Rauschert cultivated in Europe, India and South Africa**

Schmidt E<sup>1</sup>, Höferl M<sup>1</sup>, Wanner J<sup>2</sup>, Jirovetz L<sup>1</sup>, Buchbauer G<sup>1</sup>, Gochev V<sup>3</sup>, Geissler M<sup>4</sup>

<sup>1</sup>Department of Clinical Pharmacy and Diagnostics, University of Vienna, Althanstrasse 14, A-1090 Vienna, Austria

<sup>2</sup>Kurt Kitzing Co., Hinterm Alten Schloss 21, D-86757 Wallerstein, Germany

<sup>3</sup>Department of essential oils, University of Food Technologies, 26 Maritza Boulevard, 4002 Plovdiv, Bulgaria

<sup>4</sup>Shimadzu Germany, Department of GC and GC-MS, Albert-Hahn-Strasse 6-10, D-47269 Duisburg, Germany

info@artandfragrance.de

Keywords: *Chamomilla recutita*; chemotype; antibacterial activity; chemical composition

*Chamomilla recutita* (L.) Rauschert exists according to the literature from 4 to 6 chemotypes [1,2]. As the chamazulene type is not mentioned there, it could be concluded that there are probably 7 chemotypes.[3] All chemotypes are characterized by their different chemical composition. In continuation of our work on chemical composition and antimicrobial activities of essential oils we selected the following chamomile ct oils: Type A from Hungary (24.1%  $\alpha$ -bisabolol oxide A ), Type B from South Africa (12.1%  $\alpha$ -bisabolol oxide B), Type C from Hungary (11.5%  $\alpha$ -bisabolol), Type D from India (5.7%  $\alpha$ -bisabolol , 6.0%  $\alpha$ -bisabolol oxide A, 5.9%  $\alpha$ -bisabolol oxide B) Type F from Hungary (11.5%  $\alpha$ -bisabolol), 23.2% chamazulene). The essential oils were analysed using GC/FID and GC/MS on polar and apolar columns.

Antimicrobial activity of the essential oils were tested against one Gram-positive (*Staphylococcus aureus*, ATCC 6538) and three Gram-negative bacteria (*Escherichia coli*, ATCC 25922; *Salmonella abony*, ATCC 6017; *Pseudomonas aeruginosa*, ATCC 27853) as well as one yeast (*Candida albicans*, ATCC 10231) by using serial broth dilution [3] method. It could be verified, that the essential oils possess medium antimicrobial activity against Gram-positive *Staphylococcus aureus*, and Gram-negative *Escherichia coli*, *Salmonella abony* but less against *Pseudomonas aeruginosa*. The highest activity was achieved by the yeast *Candida albicans*. This shows clearly the high value of chamomile oils in medicine as well as in cosmetics.

1. H Schilcher, Die Kamille, (1987), Wiss. Verlagsges. Stuttgart

2. P Rubiolo, F Belliardo, C. Cordero, E Liberto, B Sgorbini, C Bicchi (2006) Phytochem Anal. 217-225

3. C Wagner, W Friedt, RA Marquard, F Ordon, (2005) Plant Science, 169: 917-927

4. NCCLS, Approved Standard, (1990) NCCLS Publication M7-A2, Villanova(PA) , USA,

**P 115. Antioxidant, antimicrobial and toxicological properties of *Schinus molle* L. essential oils**

Martins MR<sup>1</sup>; Alcobia D<sup>2</sup>; Arantes S<sup>1</sup>; Candeias F<sup>1</sup>; Tinoco MT<sup>1</sup>; Cruz-Morais J<sup>1</sup>

<sup>1</sup>ICAAM & Departamento de Química, Escola de Ciências e Tecnologia, Universidade de Évora, Évora, Portugal

<sup>2</sup>ICAAM & Departamento de Química, Escola de Ciências e Tecnologia, Universidade de Évora, Évora, Portugal

mrm@uevora.pt

Keywords: *S. molle*; essential oils; antioxidant activity; antimicrobial activity; brine shrimp mortality test

*Schinus molle* L. is commonly known as pink pepper or American pepper, of *Anacardiaceae* family, from subtropical regions of South America, introduced and naturalized in South Europe, including Portugal. In folk medicine, plant extracts and essential oil has related as having antibacterial, antiviral, antifungal, anti-inflammatory, antitumoral, antispasmodic, analgesic and antidepressive properties.

The aim of present study was to evaluate the chemical composition and biological activities of essential oil extracted from leaves and fruits of *S. molle*. For this purpose, the essential oils were analyzed by gas chromatography (GC/FID) and antioxidant properties were evaluated by the free radical DPPH and by system  $\alpha$ -carotene/linoleic acid methods. The antimicrobial activities were screened against pathogenic bacteria and fungi and food spoiling fungi by the disc diffusion assay and minimal inhibitory concentration (MIC) was determined for sensitive strains. Toxicity of essential oils were carried out by the brine shrimp mortality test (EC<sub>50</sub>) and acute lethal dose (DL<sub>50</sub>) determination by oral administration in Swiss mice. The major components in leaf essential oil were  $\alpha$ -phellandrene, beta-phellandrene and limonene, while myrcene,  $\alpha$ -phellandrene and 1,8-cineole are the main components in the fruit essential oil. The essential oils of leaf and fruit of *S. molle* showed antioxidant activity through the two mechanisms: the ability to capture free radicals and protection of lipid peroxidation. These oils exhibited also a broad microbial activity spectrum, against pathogenic bacteria Gram-positive and Gram-negative and *Candida* spp. The fruit essential oil showed high cytotoxicity against *Artemia salina*.

Essential oils of leaves and fruits of *S. molle* showed significant antioxidant and microbial properties, so the studies continue to clarify more in deep its toxicity, including hepatotoxicity and nephrotoxicity, and to evaluate its medicinal or nutraceutical potential.

**P 116. Chemical composition and phytotoxic effects of *Origanum vulgare* L. essential oil against *Portulaca oleracea* L. and *Conyza canadensis* (L.) Cronq.**

Sancho MV<sup>1</sup>; Salamone A<sup>2</sup>; Rellán DG<sup>1</sup>; Ferrer MAB<sup>3</sup>; Tortajada HB<sup>1</sup>

<sup>1</sup>*instituto Agroforestal Mediterráneo, Universitat Politècnica De València*

<sup>2</sup>*CRA-Unità di ricerca per il recupero e la valorizzazione delle specie floricole mediterranee*

<sup>3</sup>*Departament de Farmacologia, Facultat de Farmàcia, Universitat de València*

merversa@doctor.upv.es

Keywords: *Origanum vulgare*, essential oil, herbicidal activity, GC, GC/MS, seed germination

The chemical composition of the essential oil from a commercial sample of *Origanum vulgare* L. was determined by GC and GC/MS. A total of 23 compounds accounting for the 99.25% of the total oil were identified. The monoterpene hydrocarbons fraction constituted 50.54% of the oil, while the oxygenated monoterpenes reached 48.65%. No sesquiterpene compounds were found. The main compounds were carvacrol (29.16%), *p*-cymene (27.58%), limonene (13.63%) and thymol (12.06%).

Different chemotypes of *O. vulgare*, as carvacrol [1, 2 y 3], thymol, caryophyllene, sabinene,  $\gamma$ -terpinene and  $\beta$ -cubenene [4 y 5] have been described. Terpinen-4-ol and  $\gamma$ -terpinene were the main compounds of *O. vulgare* essential oil from Chile [6]. The essential oil here analyzed is carvacrol chemotype. This compound is usually found in samples of *O. vulgare* from Brasil [3]. Great variances have been detected in the composition of *O. vulgare* from different origins.

The herbicidal potential of the essential oil was tested *in vitro* against *Portulaca oleracea* L. and *Conyza canadensis* (L.) Cronq., two important weeds in Mediterranean crops. The higher concentrations applied (0.5 y 1  $\mu$ l/ml) were very effective on both weeds, but *C. canadensis* showed more sensibility. Its germination was completely inhibited, while *P. oleracea* germination was blocked completely only at the concentration of 1  $\mu$ l/ml, being reduced 71.6% at 0.5  $\mu$ l/ml. The lower doses (0.125 y 0.25  $\mu$ l/ml) showed no effect on *P. oleracea* germination but inhibited *C. canadensis* germination 64.2 and 92.6% respectively.

1. N Fischer et al. (1988) J. Agr. Food Chem. 36: 996-1003.

2. C Franz, J Novak (1997) In: S Padulosi (Ed.) Oregano: Proceedings of the IPGRI International Workshop, Italy, Rome, pp. 49-56.

3. MRA Rodrigues et al. (2004) J. Agr. Food Chem. 52: 3042-3047.

4. H Schulz et al. (2003) J. Agr. Food Chem. 51: 2475-2481.

5. M Baranska et al. (2005) Anal. Bioanal. Chem. 381: 1241-1247.

6. Busatta et al. (2007) Braz. J. Microbiol. 38: 610-616.

**P 117. Classification of oregano (*Origanum vulgare* L.) genotypes grown in Latvia**

Galoburda R; Kruma Z; Paura L; Alsina I

*Latvia University of Agriculture*

*ruta.galoburda@llu.lv*

Keywords: oregano, volatiles, genotype, morphological parameters

Oregano (*Origanum vulgare* L.) is a perennial herb, belonging to plant family *Lamiaceae*, which grow in Latvia both wild and cultivated. A number of studies showed that oregano is very variable in chemical composition of essential oils depending on geographical origin, vegetative stage and used treatment. The aim of current research was to classify oregano genotypes based on the volatile composition and morphological parameters.

Ten genotypes of oregano were analysed in the current study. Flowers and leaves were collected separately; air dried and stored in sealed bags until analysis. Morphological parameters were determined according to standard methods. Volatiles from dried oregano were extracted using headspace autosampler and analyzed by gas chromatography-mass spectrometry. Compounds were identified by comparison of their mass spectra with mass spectral libraries (Nist98), and by calculation of linear retention indexes and comparison with literature data. The data obtained from the analysis of the oregano were analyzed by means of multivariate analysis; employing hierarchical cluster analysis. The method used was linkage between-groups. The distances between samples were calculated using square Euclidean distances. As a pre-treatment of data was carried transform values of variables (average zero and standard deviation 1) called Z scores. The dendrogram similarity scales that are generated by the SPSS program range from zero (greater similarity) to 25 (lower similarity). A linear correlation analysis was performed in order to determine relationship between morphological parameters and volatile compounds.

The studied oregano genotypes differ both in quantitative and qualitative content of aroma compounds. Also the dendrograms obtained by hierarchical cluster analysis showed that the genotypes of oregano are quite heterogeneous. Totally 35 volatile compounds were identified in Latvian oregano samples. Sabinene, caryophyllene, germacrene D and Z- $\beta$ -ocimene are the major aroma forming compounds. Mainly weak correlations between volatile compounds and morphological parameters were detected. Moderate correlation ( $r = -0.51$ ) between oregano flower color and total peak area of identified aroma compounds were observed, meaning that more volatiles are in plants with more pink flowers instead of dark violet.

**P 118. Chemical composition, leishmanicidal, cytotoxic and hemolytic activity of the essential oil from leaves of *Cymbopogon winterianus* Jowitt ex Bor.**

Dias CN<sup>1</sup>, Rodrigues KAF<sup>2</sup>; Lima DS<sup>2</sup>; Carneiro SMP<sup>2</sup>; Carvalho FAA<sup>2</sup>; Alves LPL<sup>1</sup>; Maia JGS<sup>3</sup>; Moraes DFC<sup>1</sup>

<sup>1</sup>Programa de Pós-Graduação em Saúde e Ambiente, Universidade Federal do Maranhão, 65085-580 São Luís, MA, Brazil;

<sup>2</sup>Núcleo de Pesquisas em Plantas Medicinais, Programa de Pós-Graduação em Farmacologia, Universidade Federal do Piauí, 64049-550 Teresina, PI, Brazil;

<sup>3</sup>Programa de Pós-Graduação em Química, Universidade Federal do Pará, 66075-900 Belém, PA, Brazil

cla\_noleto@yahoo.com.br

Keywords: *Cymbopogon winterianus*; cytotoxic; hemolytic; leishmanicidal

*Cymbopogon winterianus* (Poaceae) is usually known as citronella grass and is used in folk medicine as analgesic, anxiolytic and anticonvulsant. This study aimed to evaluate the toxic activity of essential oil (EO) from leaves of *C. winterianus* against promastigotes forms of *Leishmania amazonensis* as well as its cytotoxic activity to macrophages and its hemolytic potential on human erythrocytes. The EO was obtained by hydrodistillation and then analyzed by GC and GC/MS. All bioassays were carried out at concentrations of 400; 200; 100; 50; 25; 12,5; 6,25 e 3,12 mg/L of the essential oil. The leishmanicidal activity was evaluated *in vitro* using  $1 \times 10^6$  promastigotes per well in 96-well plates with the EO and then counted in a Neubauer haemocytometer at 24, 48 and 72 h [1]. In the cytotoxicity assay, murine peritoneal macrophages were incubated for 48 h at the tested concentrations, then MTT was added and the results were analyzed in a spectrophotometer [2]. The hemolytic activity was investigated using erythrocytes (human O<sup>+</sup>) and cell lysis was determined spectrophotometrically [3]. The leaves of *C. winterianus* provided colorless oil, with yield of 1.3%. Among the identified constituents, monoterpenes were 75.8% and sesquiterpenes 19.0%. The major constituents were citronellal (26.5%), geraniol (16.2%) and elemol (14.5%). In the leishmanicidal test, the oil causes 100% of inhibitory effect at the concentration of 400 mg/L and the IC<sub>50</sub> value was 89 mg/L after 72h of exposure. The EO caused low toxicity against the macrophages with a CC<sub>50</sub> value of 376 mg/L. The toxic action of oil on erythrocytes was also low, with 14.2%, 11% and 6.7% of hemolysis at concentrations of 400, 200 and 100 mg/L respectively and 0% of hemolysis in other concentrations. The specie exhibited promising activity as a leishmanicide, with high selectivity for promastigote forms of *L. amazonensis* and with low toxicity to erythrocytes and macrophages.

Acknowledgments: The authors are grateful to CNPq/BIONORTE, FAPEMA and CAPES for their financial support.

1. Oliveira-Silva et al. (2008) Am. J. Trop. Med. Hyg. 78: 745-749.

2. Medeiros et al. (2011) Paras. Inter. 60: 237-241.

3. LÖFGREN et al. (2008) Exp. Paras. 118: 197-202.

**P 119. Phytochemical screening and antinociceptive activity of *Lippia microphylla* CHAM. (Verbenaceae) leaf essential oil in rodents**

André Sales Barreto<sup>1</sup>, Pollyana de Souza Siqueira Lima<sup>1</sup>; Angélica Maria Lucchese<sup>2</sup>; Pericles Barreto Alves<sup>1</sup>; Darlisson de Alexandria Santos<sup>1</sup>; Jullyana de Souza Siqueira Quintans<sup>1</sup>; Douglas S. Prado<sup>1</sup>; Roberta S.D. Cardoso<sup>1</sup>; Natalia Barreto Rocha Souto<sup>1</sup>; Lucindo José Quintans Júnior<sup>1</sup>

<sup>1</sup>Federal University of Sergipe;

<sup>2</sup>Universidade Estadual de Feira de Santana

asbfisio@hotmail.com

Keywords: *Lippia microphylla*; Monoterpenes; Camphor; (*E*)-Caryophyllene; Pain

*Lippia microphylla* Cham. (Verbenaceae) is a medicinal plant known in Brazilian Northeast as 'alecrim-de-tabuleiro' and it is a plant used popularly to treat heart disease, inflammatory and painful disorders. However, there is little information about *L. microphylla*'s biological properties. Thus, we investigated the antinociceptive effect of the essential oil obtained from the leaves of *L. microphylla* (LEO) in mice. LEO was obtained by hydrodistillation in a Clevenger-type apparatus using 1200 g of dried leaves. The identification of the components was made through comparison of substance mass spectrum with the database of the GC-MS, literature and retention index. Male mice (26-30g) were pretreated with LEO (25, 50 or 100 mg/kg, orally route, *per os*, p.o.), morphine (5 mg/kg, i.p.) or vehicle (distilled water + Tween 80 0.2%), before acetic acid- (0.85%) (writhing reflex) or formalin (20 µl of 1%)-induced nociception tests. The motor coordination was also evaluated using Rota rod (8 rpm, 180 s). Experimental protocols and procedures were approved by the Universidade Federal de Sergipe Animal Care and Use Committee (CEPA/UFS # 26/09). The obtained data were evaluated by one-way analysis of variance (ANOVA) followed by Tukey's test. The phytochemical analysis of LEO demonstrated the presence of camphor (28.698%), (*E*)-caryophyllene (11.47%), bicyclogermacrene (10.22%), camphene (9.45%) and borneol (7.64) and as the main compounds. LEO pre-treatments inhibited writhing ( $p < 0.001$ ) in the acetic acid test, 25 ( $17.0 \pm 1.5$ ), 50 ( $15.5 \pm 1.9$ ) or 100 ( $10.1 \pm 1.8$ ) mg/Kg, respectively, when compared to control group ( $28.5 \pm 2.9$ ). In the formalin test first phase only the doses of 25 ( $42.2 \pm 5.9$ ), 50 ( $41.3 \pm 4.9$ ) or 100 ( $35.2 \pm 3.4$ ) mg/Kg had promoted a reduction in the time spend licking the paw with  $p < 0.01$  or  $p < 0.001$ , when compared to control group ( $64.7 \pm 4.3$ ). In addition, in the second phase all doses reduce the time ( $p < 0.001$ ) at doses 25 ( $72.6 \pm 8.6$ ), 50 ( $41.8 \pm 5.0$ ) or 100 ( $27.2 \pm 4.1$ ) mg/Kg, respectively, when compared to control group ( $136.5 \pm 8.9$ ). Such results were unlikely to be provoked by motor abnormality (data not shown). Our results suggest that *L. microphylla* leaf essential oil modulates inflammatory and central nociception and might represent important tool for management and/or treatment of painful conditions.

Acknowledgments: Financial Support by FAPITEC/SE/Brazil, CNPQ/Brazil, CAPES/Brazil.



**P 120. Evaluation of the cytotoxicity of an essential oil obtained from leaf of *Xylopi*a *laevigata***

Grace Anne Azevedo Doria<sup>1</sup>, Rebeca O Souza<sup>2</sup>; Emmanoel Vilaça Costa<sup>1</sup>; Lucindo José Quintans Júnior<sup>1</sup>; Maria José V Fonseca<sup>2</sup>; Adriano Antunes de Souza Araújo<sup>1</sup>; Thanany B. da Silva<sup>1</sup>; Jullyana de Souza Siqueira Quintans<sup>1</sup>

<sup>1</sup>Federal University of Sergipe;

<sup>2</sup>University of São Paulo - RP

gracedoria@hotmail.com

Keywords: *Xylopi*a *laevigata*, citotoxicity, phytochemical analysis

Investigations have explored that natural products have been source of most active ingredients of medicines. For that reason, it is necessary to evaluate the toxicity of natural products, to guarantee the secure use. *Xylopi*a *laevigata* unknown to scientists have been use for years by Brazilian people to cure diseases, such as kidney problems and inflammations. In this behalf, the proposal of this study was to investigate the citotoxicity of an essential oil obtained from leaf *Xylopi*a *laevigata* (EOX).

EOX was extracted by hydrodistillation of the fresh leaves of *X. laevigata*. The components of the EOX were identified by comparison of substance mass spectrum with the database of the GC-MS, literature and retention index. For the cytotoxicity assay, L929 fibroblast cell lines were plated in 96-well plates and the essential oil was added in serial dilutions for 24 hours. After that, the viability of fibroblast cells was quantified by neutral red (NR) assay. The values were obtained of three independents tests.

The phytochemical analysis of EOX showed the presence of g-muurolene (17.78%),  $\delta$ -cadinene (12.23%), bicyclogermacrene (7.77%),  $\alpha$ -copaene (7.17%), germacrene D (6.54%) and (E)-caryophyllene (5.87%) as the main compounds. It was observed that EOX at 100  $\mu\text{g/mL}$  and 200  $\mu\text{g/mL}$  decreased 40% and 80% of the cell viability, respectively. Therefore, the value of IC<sub>50</sub> was 125.0  $\mu\text{g/mL}$  to fibroblast cell line.

Research with essential oils of plants whose similar components described in EOX showed cytotoxic activity against tumor cells, such as germacrene D and bicyclogermacrene [1]. It is suggested that the reduction of the cell viability observed can be associated the presence of these compounds in EOX. New studies are required for elucidation of the mechanisms of EOX cytotoxic activity.

Acknowledgments: Financial support by CNPq, FAPITEC/SE.

1. MC Palazzo et al. (2009). *Rec. Nat. Prod.* 3: 32-37

**P 121. Residues of fungicides in citrus essential oils from Brazil**Rodrigues AS<sup>1</sup>, Alves AAR<sup>1</sup>; Rezende CM<sup>1</sup>; Bizzo HR<sup>2</sup><sup>1</sup>Instituto de Química, Universidade Federal do Rio de Janeiro;<sup>2</sup>Embrapa Food Technology

bizzo@ctaa.embrapa.br

Keywords: fungicides residues; citrus essential oils; essential oils from Brazil; carbaryl

Brazil is one of the largest exporters of essential oils (EO), and *circa* 95% of the exported volume is due to citrus oils [1]. The country is also one of the largest consumer of fungicides. Residues of organochlorides and organophosphorides pesticides have been found in Brazilian citrus oils in amounts well above those allowed by *Codex Alimentarius* standards [2]. Continuing a systematic investigation on contaminants from Brazilian citrus oils, the objective of this work was to develop a method for the detection and quantification of fungicide residues, namely carbaryl, delta-cyhalothrin, dithianon, prochloraz, tebuconazole and thiabendazole in citrus oils. The oils were passed through solid phase extraction column (C18) and after elution and concentration, the samples were analyzed by gas chromatography coupled to mass spectrometry using a DB-5 (30m X 0.25mm X 0.25 µm) capillary column. Helium was used as carrier gas at 1.0mL/min. Mass detector was operated in selective ion monitoring (SIM) in electronic ionization mode at 70eV. Calibration curves were built for each fungicide with standards. The limits of detection (LOD) and quantification (LOQ) were, respectively, 6.25 mg/L and 15,6 mg/L for carbaryl, 0.61 mg/L and 1.53 mg/L for delta-cyhalothrin, 1.33 mg/L and 3.32 mg/L for dithianon, 0.94 mg/L and 2.36 mg/L for prochloraz, 0.60 mg/L and 1.51 mg/L for tebuconazole and 2.21 mg/L and 5.53 mg/L for thiabendazole. Fifteen samples of commercial oils from different producers were analyzed. In one sample of lime oil (OE1) were found residues of carbaryl (18.0 mg/L), dithianon (7.9 mg/L) and tebuconazole (6.81 mg/L). A second sample of lime oil (OE10) was contaminated with 16,1 mg/L of carbaryl. Delta-cyhalothrin was detected in sample OE2 (orange oil) in 3.87 mg/L. In these oils, the residues found are well above those stated in *Codex Alimentarius*.

Acknowledgments: Conselho Nacional de Desenvolvimento Tecnológico (CNPq).

1. Bizzo HR et al. (2009) Quim. Nova 32: 588-594.

2. Alves AAR et al. (2012) J. Braz. Chem. Soc. 23: 306-314.

**P 122. Hydrodistillation vs steam distillation coupled to Clavenger vs Dering essential oil separators in antioxidant molecules recovery from aromatic plants**

Zhao<sup>1</sup>, Menut<sup>2</sup>; Venskutonis<sup>3</sup>; Talou<sup>1</sup>

<sup>1</sup>Universite de Toulouse, INP-ENSIACET;

<sup>2</sup>Universite de Montpellier, Institut des biomolécules Max Mousseron;

<sup>3</sup>Kaunas Technology University

talou@cict.fr

Keywords: Aromatic plants; natural antioxidants; steam distillation; hydrodistillation; Dering apparatus; Clavenger apparatus

Essential oils were extracts obtained by hydrodistillation or steam distillation of aromatic plants combined to an essential oil separator, either the Clavenger apparatus, very popular in West Europe, nor the Dering apparatus, widely used in Eastern Europa countries. In otherhand, apart the odorous fraction, aromatic plants appeared as a valuable source for natural antioxidant molecules. If such components could be obtained from essential oils, extraction cake which represented up to 95% of the raw distilled material was reported as a potential complementary source for natural antioxidants. In the framework of the AROMATIC Program, we have applied to Medieval Aromatic Plants (aka forgotten ones) the agrorefinery concept based on a valorization of the entire plant by sequential extractions of molecules of interest while not penalizing the subsequent valorization of residual by-products as biosourced molecules [1]. This strategic approach was recently applied to coltsfoot (*Tussilago farfara*) originated from Europe (France, Lithuania) or Asia (China) [2]. In the present paper, we reported the cross-evaluation of two extraction process (HydroDistillation (HD) and Steam Distillation (SD)) combined with two essential oil separators (Clavenger Apparatus (CA) and Dering Apparatus (DA) on both essential oils composition and antioxidant molecules recovery from extraction cake. Results are expressed as comparison of HD-CA, HD-DA, SD-CA and SD-DA for i) volatiles chromatographic profiles of essential oil by GC-FID and GC-MS; ii) antioxidant activities of methanolic extracts of corresponding extraction cakes evaluated by measuring radical scavenging activity with DPPH assay. If GC profiles are similar in the four cases, extraction yields of essential oils were higher for HD than SD while for antioxidant fractions opposite was reported. Regarding DA vs CA, if the first one allow to minimize the essential oils extraction time while the second make easier their collection, their impact on the antioxidant fraction recovery were limited [3].

[1] C El Kalamouni, T Talou, C Raynaud, R Venskutonis (2009), *Chemine Technologija*, 3 (52), 69-73

[2] T M Zhao, D Dobravalskyte, C Menut, R Venskutonis, T Talou (2012), Study of antioxidant activity of various coltsfoot extracts originated from France, China and Lithuania, 7th Baltic Conference on Food Science and Technology, Kaunas (Lithuania)

[3] T M Zhao, INP Doctorate thesis (2013, in progress)

**P 123. Crushing finger device and flash aroma dispenser: two innovative tools for native oils analysis**Dobravalskyte<sup>1</sup>, Zhao<sup>1</sup>; Talou<sup>1</sup><sup>1</sup>Universite de Toulouse, INP-ENSIACET

talou@cict.fr

Keywords: Aromatic plants; native oils; artificial crushing finger device; flash aroma dispenser

Essential oil plants are aromatic crops used in perfumery, cosmetic and food industry which extracts were obtained by hydrodistillation or steam distillation. But such degrading methods generated artefacts due mainly to in situ chemical reactions (hydrolysis) and consequentatively the obtained extract's odor could differ from initial plant ones. Access to the naturally emitted odorants was possible by crushing between fingers a couple of leaves or flowers followed by immediate sniffing of the generated volatiles. But the poor reproducibility of such manual method did not allow to apply it for an instrumental analysis of emitted volatiles. In the framework of the AROMATIC Program, we developed/adapted devices in order to analyze native volatile compounds emitted just after instrumental crushing of leaves, i.e. Artificial Crushing Finger device (ACF) and Flash Aroma Dispenser (FAD) respectively based on a mechanical and a gas flash pressure crushing. By exploding storage vesicles located on leaves, ACF and FAD, coupled with either a Tenax trap or SPME system, allowed to perform a rapid evaluation of native oil content with only a couple of leaves. First developed and based on a modified artificial mouth, AFD allowed to instrumentally mimic the total crushing of leaves between fingers but appeared for fragile leaves to be too destructive [1]. FAD is based on a whipped cream siphon in which a gas (NO<sub>2</sub> or CO<sub>2</sub>) flash pressure increase, allowed to reach to a partial explosion of storage vesicles and consequentatively appeared to be a more respective method. In the present paper, we reported the comparison of volatiles chromatographic profiles of essential oil from our model plant, balsamite (*Tanacetum balsamita*), obtained by hydrodistillation with the ones obtained with crushed entire fresh leaves processed by ACF or FAD for which emitted volatiles were concentrated by SPME. Microscopic observations of processed leaves allowed to compare ACF and FAD efficiency regarding explosion of vesicles or secretory hairs and destruction of leaf's structure [2]

[1] D Dobravalskyte, C El Kalamouni, JL Berdague, P Tournayre, C Raynaud, T Talou (2009), ISEO 2009, Savigliano (Italy)

[2] Dobravalskyte D, co-tutelle Doctorate Thesis INP-KTU, (2013, in progress)

**P 124. Phytotoxicity and antimicrobial activity of essential oils in plant products treatment**

Smid J<sup>1</sup>, Kloucek P<sup>1</sup>; Frankova A<sup>1</sup>; Kuchtova P<sup>1</sup>

<sup>1</sup>Czech University of Life Sciences

smidj@af.czu.cz

Keywords: Essential oil; Seed treatment; Shelf-life; Organic farming; Model system; Pathogens

Essential oils (EOs) cause disturbance of cell membranes leading to cell death. This effect is used in antimicrobial treating, but plant products treatment causes tissue damage and irreversible degradation of the treated material. One of the main problems during treatment with EOs is keeping vitality and structure of fresh products (such as fruits, vegetables and seeds). Products should not be exposed to EO in high concentration for long time.

EOs of oregano (*Origanum vulgare*), cinnamon (*Cinnamomum zeylanicum*), clove (*Syzygium aromaticum*) and lemon grass (*Cymbopogon citratus*) were tested for antimicrobial activity on poppy and wheat seeds, potato, carrot, cucumber, strawberry, mung beans, apple and lemon. Products were treated by different methods using EOs vapours. Every treatment shows decrease of antimicrobial contamination and, in higher or equal concentrations of EOs, phytotoxicity detectable by browning of fruits and vegetables and by reduction in seeds germination.

Generally, the EOs from oregano and cinnamon have strongest antimicrobial effect, but they have strongest negative effect on plant material also. Clove EO applied on seeds still has required antimicrobial effect with less damage on plant materials in comparison with both previous EO. Lowest effect in antimicrobial activity and phytotoxicity shows lemon grass EO. For example, oregano EO inhibits growth of *Alternaria alternata* on carrot roots in 16  $\mu\text{l.l}^{-1}$  and cause tissue damage in 32  $\mu\text{l.l}^{-1}$ .

Antimicrobial activity of EOs is well-known, but has been compared with phytotoxicity in literature only rarely [1]. As our results demonstrate phytotoxicity of EOs has to be taken into account in model systems.

Acknowledgements: This presentation is supported by University internal grant agency - CIGA 20122008 – Plant products treatment with essential oil vapours under different physical conditions.

1. MA Ibrahim et al. (2001) Agric. Food. Sci. Fin. 10/3: 243-259.

**P 125. Poppy volatiles and their role in attracting poppy root weevil**

Pavel Kloucek<sup>1</sup>, Petr Vlazny<sup>1</sup>; Blanka Kalinova<sup>2</sup>; Michal Hoskovec<sup>2</sup>; Jakub Smid<sup>1</sup>; Adela Frankova<sup>1</sup>

<sup>1</sup>University of Life Sciences Prague;

<sup>2</sup>Institute of Organic Chemistry and Biochemistry

kloucek@af.czu.cz

Keywords: *Papaver rhoeas*; *Papaver somniferum*; *Stenocarus ruficornis*; distillation

The Czech Republic is the world's largest producer of low morphine food grade poppy (*Papaver somniferum*) with yearly average of 30 000 ha of harvested area. Poppy root weevil (*Stenocarus ruficornis*) is one of its major pests causing large losses of young plants. The adult beetles overwinter in the soil or plant debris and become active in the spring. They search for poppy plants, mate, feed and place eggs almost exclusively on different *Papaver* species [1]. The heaviest damage is done by larvae feeding on the roots. Therefore, we have decided to investigate, if some of the volatile compounds emitted by *Papaver* spp. are able to attract the beetle.

Fresh aerial plant material of autumn- and spring-sown low morphine poppy and corn poppy plants were distilled in clevenger apparatus and the volatiles were captured into layer of hexane due to low yield. GCxGC-MS and GC-EAD analyses revealed that the main components were phytol, fatty acids and green leaf volatiles. Among them, 3Z-hexen-1-ol, 3Z-hexenyl acetate and other compounds were identified as active by EAD when using *S. ruficornis* antenna.

The results could be used for development of signalization traps and hence for effective application of insecticides.

**P 126. Chemotypes of *Eugenia uniflora* (Myrtaceae) from Rio de Janeiro, Brazil**André AC, Alves FC; Santos-Junior HM; Rezende CM

Universidade Federal do Rio de Janeiro

crezende@iq.ufrj.br

Keywords: *Eugenia uniflora*; essential oil; sesquiterpene chemotypes; antinociceptive activity

*Eugenia uniflora* L. (Myrtaceae), native of Brazil and known as Surinam cherry, is of great interest to food and cosmetic Brazilian industries. In this study, we evaluated the existence of *E. uniflora* chemotypes from the essential oils obtained from the leaves of some previously investigated material, related to genetic variability [1] and collected at Grumari, Rio de Janeiro, RJ. GC-MS data were analyzed using Principal Component Analysis (PCA) and resulted in four different groups (A-D). Group A, consisting of nine specimens, showed a major peak with IRL equal to 1516. After NMR analysis, two co-eluting isomers were identified as 3-atractylone and furanoeudesmene [2]. The group B (eight specimens) showed a major peak with IRL 1774. Analysis by NMR resulted in the structure of the sesquiterpene 6-beta-acetoxy-5alpha-H-guaian-1(10),3-diene. The four representatives of the group C showed a chromatographic profile with two major components, IRL 1674 and 1787, which were identified by NMR as selina-1,3,7 (11)-trien-8-one and its oxide, respectively. In group D (nine specimens) two major peaks were observed. The first IRL is in accordance with group A. The second (IRL 1870) is an epicurzerenone isomer. Analysis of variance (One-way ANOVA) complemented with Dunnett's test showed that all treatments, at the dose tested (200 mg / kg), were effective in antinociceptive activity using albino mice (Swiss) 2, reducing the number of writhings compared with the control group (Tween 20/EtOH/H<sub>2</sub>O 1:1:10, \* p<0.05). Due to popular usage, the National Health Surveillance Brazilian Agency - ANVISA - regulated the use of cherry tree leaves in teas in the 5th edition of the Brazilian Pharmacopoeia (2010). Through its essential oil, *E. uniflora* is regulated by the presence of curzerene isomers (IRL 1845). However, it is demonstrated in this work that this species has different sesquiterpenes chemotypes and that other chemical markers should be used for its characterization, that are possibly related to its biological activity.

Acknowledgements: CNPq, CAPES, FAPERJ.

1. Margis R et al. (2002) Biodivers. Conserv. 11: 149-163.
2. Amorim AC L et al. (2009) Phytomedicine 16: 923-928.

**P 127. Biotransformation of geraniol by *Penicillium oxalicum* and *Aspergillus niger***

Alves PB<sup>1</sup>, Jesus HCR<sup>1</sup>; Franco CRP<sup>2</sup>; Silva GH<sup>3</sup>; Porto ALM<sup>4</sup>

<sup>1</sup>Federal University of Sergipe, Department of Chemistry, 49100-000, São Cristóvão, SE, Brazil;

<sup>2</sup>Federal University of Sergipe, Department of Biology, 49100-000, São Cristóvão, SE, Brazil;

<sup>3</sup>Federal University of Sergipe, Department of Chemistry, 49500-000, Itabaiana, SE, Brazil;


<sup>4</sup>University of São Paulo, Institute of Chemistry of São Carlos, 13560-970, São Carlos, SP, Brazil

periclesbalves@gmail.com

Keywords: biotransformation; geraniol; *Penicillium oxalicum*; *Aspergillus niger*

This study sought to evaluate the bioconversion of geraniol by the fungi *Penicillium oxalicum*, isolated from the marine alga *Caulerpa* sp, and *Aspergillus niger* obtained from *Hancornia speciosa* Gomes leaves. Geraniol (50 mg dissolved in 400 µL of DMSO) was added in 250 mL Erlenmeyer flasks containing mycelium of the fungus previously grown for three days in 100 mL of malt extract (2%). The mixtures were incubated in a shaker (150 rpm) at 32 °C, and the reaction progress was monitored periodically by removing aliquots for GC-MS (Shimadzu, QP-5050A model) analysis. Identification of reaction products was carried out by comparison of the observed *spectra* with spectra from the literature<sup>1</sup> and confirmed by co-injection of authentic standards. An oxidative process was observed in the biotransformation of geraniol by *P. oxalicum*, producing pure sulcatone (Figure 1). In the reaction, all the substrate has been converted into sulcatone in just 40 hours, showing a high catalytic activity of the fungal enzymes. Demeyttenare & Pooter (1996) reported a similar reaction using *Penicillium italicum*, but with a lower yield of sulcatone (69-76%) and a longer reaction time (4 days). In the biotransformation of geraniol by *A. niger* there was an isomerization of the geraniol to form linalool. The conversion rate was 17% in 120 hours. After this period, there were no significant changes in the chromatographic profile. This study has demonstrated that the fungus *P. oxalicum*, through its associated enzymes, is an efficient biocatalyst for conversion of geraniol to sulcatone with a high purity (99% as checked with GC) and a molar yield of >99% during a short period of 40 hours. Biotransformations with *A. niger* converted geraniol into (+/-)-linalool (40% after 120 h).

Figure 1. Bioconversion of geraniol by *P. oxalicum* and *A. niger*



1. RP Adams. (2007) *In: Identification of essential oil components by gas chromatography/quadrupole mass spectroscopy*, 4<sup>th</sup> Ed. Illinois USA: Allured Publishing Corporation, Carol Stream.
2. JCR Demyttenaere, HL De Pooter (1996) *Phytochemistry*. 41:1079-1082.



**P 128. Biological activity of essential oil from leaves of *Tetradenia riparia* Hochstetter. Codd. (Lamiaceae) grown in western Amazon, Brazil**

Torquilha HS<sup>1</sup>, Godoy RLO<sup>2</sup>, Bobeda CRR<sup>1</sup>, Freitas-Silva O<sup>2</sup>, Sobrinho AFS<sup>2</sup>, Rosas EC<sup>3</sup>, Henriques MCMO<sup>3</sup>, Leon LL<sup>3</sup>, Bozza M<sup>3</sup>

<sup>1</sup>IFRJ Campus Nilópolis, Brazil

<sup>2</sup>EMBRAPA, Brazil

<sup>3</sup>FIOCRUZ, Brazil

hpal@ig.com.br

Keywords: Myrrh; essential oil; biological activity; antileishmanial activity; *Trypanosoma cruzi* activity; neoplastic activity

*T. riparia* Hochst.Codd. is a species from Lamiaceae family that have been used in traditional medicine in several places around the world [1,2,3,4]. In Brazil, this exotic species was the introduced firstly as an ornamental plant [5]. Its volatile oil was already reported [6]. Although its current use as a medicinal herb there is a need to know all about its pharmacological properties as well the search for new botanicals inhibitors to provide a competitive alternative to current drugs. The aim of this study was to evaluate some biological activities of essential oil obtained from fresh leaves of *T. riparia*. The results showed the oil had no cytotoxicity property, without inflammatory activity based on carrageenan-induced pleurisy on live animals, and protein leakage. However the anti-inflammatory activity in mouse paw edema showed 56.69%, and 95.98% at a dose of 1 mg/well and 99.22% at a dose of 10 mg/well, presenting the production of nitric oxide *in vitro*. *T. riparia* oil also presented a significant antileishmania activity to *Leishmania amazonensis* promastigotes. *In vitro* this oil was assayed at concentrations 5.0, 10.0, 20.0, 40.0, 80.0, 120.0, 160.0 and 320.0 µg/mL for 24h. Promastigotes viability was analyzed and the IC<sub>50</sub> was 0.47 mg/mL. The evaluation of the antineoplastic activity of essential oil of myrrh was performed *in vitro* using seven tumor cell lines. The results demonstrated that the essential oil of myrrh was able to inhibit cell growth in five of the seven strains studied (SP2 / 0 with IC<sub>50</sub> =86.60%; Neuro-2a with CI<sub>50</sub> = 86.60%; P3653 with IC<sub>50</sub> = 74.60%; BW with IC<sub>50</sub> = 82.00% and Ehrlich carcinoma with CI<sub>50</sub>=68.48%). Leishmanicidal activity was demonstrated by MTT colorimetric assay. This oil showed potent toxic effects on epimastigote forms of *Trypanosoma cruzi*, with values of IC<sub>50</sub> = 100% at 0.6 mg/mL. The biological activity potential of the oil of myrrh showed a wide spectrum and therefore studies in this line of research should be continued.

**P 129. Influence of storage time and temperature on the chemical composition of *Hyptis pectinata* (L.) Poit essential oil**

Alves PB<sup>1</sup>, Jesus AS<sup>2</sup>; Blank A<sup>2</sup>; Alves MF<sup>2</sup>; Arrigoni-Blank MF<sup>2</sup>; Lima RN<sup>1</sup>

<sup>1</sup>Departamento de Química, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil.;

<sup>2</sup>Departamento de Engenharia Agrônômica, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil.

periclesbalves@gmail.com

Keywords: *Hyptis pectinata*; medicinal and aromatic plant; volatile oil; postharvest; chemical constituents

The aim of this work was to evaluate the influence of storage and temperature on the chemical composition of *Hyptis pectinata* L. Poit essential oil. The effect of storage time of the essential oil of *H. pectinata* was evaluated (0, 15, 30, 60, 90, 120, 150, 180, 240, 300 and 360 days) at two temperatures [room ( $\pm 27^{\circ}\text{C}$ ) and freezer ( $-20^{\circ}\text{C}$ ) temperature]. The essential oil was obtained from dry leaves by hydrodistillation using a Clevenger apparatus. Qualitative analysis of the chemical composition of the essential oils was performed using a gas chromatograph coupled to a mass spectrometer (GC-MS). A quantitative analysis of the essential oil components was conducted in a gas chromatograph equipped with a flame ionization detector (FID). Comparing the compounds in the essential oil of *H. pectinata*,  $\beta$ -caryophyllene was notable due to its high concentration, decreasing from 44.88% to 42.91% when stored at room temperature for 360 days and increasing from 44.88% to 46.15% when stored in freezer for 300 days. In addition, caryophyllene oxide was also found to be a significant compound, increasing from 15.54% to 21.82% when stored at room temperature for 360 days and from 15.54% to 18.15% when stores in freezer for 300 days. The storage of essential oil in freezer resulted in minor instability of the major constituents. The essential oil of *H. pectinata* may be stored at  $-20^{\circ}\text{C}$  for 10 months.

Acknowledgments: We thank FAPITEC/SE, CNPq and CAPES for their financial support.

**P 130. Essential oils from *Hypericum undulatum* Schousboe ex Willd**Guedes AP<sup>1</sup>, Sousa RMOFS<sup>1</sup>; Fernandes-Ferreira M<sup>2</sup>

<sup>1</sup>Centre for the Research and Technology of Agro-Environmental and Biological Sciences, Quinta de Prados, Apartado 1013, 5001-801 Portugal; Department of Biology, University of Minho, Campus Gualtar, 4710-057 Braga, Portugal;

<sup>2</sup>Centre for the Research and Technology of Agro-Environmental and Biological Sciences, Quinta de Prados, Apartado 1013, 5001-801 Portugal; Department of Biology, Faculty of Science, University of Porto, 4169-007 Porto, Portugal; Mapprod Lda, Braga, Portugal

rosemarie@bio.uminho.pt

Keywords: essential oils; leaves; flowers; ripened seed capsules; *n*-nonane; sesquiterpenes.

The constituents of the essential oils (EO) of fresh leaves, stems, ripened seed capsules and flowers of *Hypericum undulatum* (Clusiaceae) obtained by hydro-distillation were identified by GC-MS and quantified by GC. A different EO composition profile was recorded for each one of the different organs. The highest yield of total EO expressed in terms of percentage by biomass dry weight was obtained from leaves (0.9%), followed by ripened seed capsules (0.8%), flowers (0.6%) and stems (0.3%). The most complex composition profile was that of flowers (97 compounds) followed by ripened seed capsules, leaves and stems with 85, 83 and 47 compounds, respectively. The *n*-alkanes group, from which *n*-nonane was the dominant compound corresponded to 84%, 42%, 37% and 24% of the EO of stems, flowers, ripened seed capsules, and leaves, respectively, being the major group of EO constituents in the *H. undulatum* organs with the exception for leaves where the sesquiterpene hydrocarbons group predominated (59%). A complete series of *n*-alkanes from C<sub>22</sub> to C<sub>29</sub> was identified in leaves whereas in stems the only long-chain *n*-alkanes detected were *n*-heptacosane and *n*-nonacosane. Although the EO from leaves contained a broad range of sesquiterpene hydrocarbons, the dominant compound was the *n*-nonane (21.6%). The absolute content of this *n*-alkane corresponded to around 0.2% of the dried biomass of leaves and around 0.3% of the dried biomass of stems, flowers and ripened seed capsules of *H. undulatum*. The second major group of compounds in the EO of leaves was that of alkanes while in EO of stems, flowers and ripened seed capsules was that of sesquiterpene hydrocarbons which represented around 8%, 34% and 23%, respectively.

Acknowledgements: This work was sponsored by EU (FSE/FEDER) and Portuguese Government (FCT) through the Project PTDC/AGR-AAM/70418/2006 (*Hypericum* Biotech).

**P 131. Identification of volatile constituents in freeze-dried powder bark of mature fruit *Syzygium malaccense* Merr. & Perry using selective techniques SPME and GC-MS.**

Bobeda CRR<sup>1</sup>, Torquillo HS<sup>1</sup>; Machado AMR<sup>1</sup>; Godoy RLO<sup>2</sup>; Pacheco S<sup>2</sup>; Andressa MS<sup>2</sup>; Gouvea ACSM<sup>2</sup>

<sup>1</sup>IFRJ-Campus Nilópolis-Brazil;

<sup>2</sup>Embrapa-Brazil

claudio.bobeda@ifrj.edu.br

Keywords: Myrtaceae; *Syzygium malaccense*; SPME; GC-MS; essential oil.

The family Myrtaceae is represented by approximately 140 genera, which together has more than 3,000 species that are distributed in tropical and subtropical regions of the Midwest and Southeast regions of Brazil, reaching beyond the limits of the country to reach the land of Uruguay, Argentina and Paraguay. The fruits of *S. malaccense* (jambo) have economic importance, being consumed raw or as candies, ice cream and refreshments, and they are used as flavoring in alcoholic distillates. The literature reports that studies of the volatile components carried out with whole ripe fruit of the genus *Syzygium*, using the hydrodistillation showed the presence of monoterpenes, sesquiterpenes and other volatiles compounds like alcohols, aldehydes and esters [1,2]. The objective of this study was to identify the volatile chemical components in freeze-dried powder of the bark of mature fruits of *S. malaccense* using polydimethylsiloxane fiber SPME and GC-MS techniques. The mature fruits were purchased at supermarkets in the metropolitan region of Rio de Janeiro (Brazil) and their peels were removed, frozen and then lyophilized before SPME-GC-MS analysis. The mass spectra obtained were compared with the NIST mass spectra library [3] and data from literature [4], showing a profile of monoterpene, sesquiterpenes, esters, aldehydes and alcohols which the major constituents are (*E*)-3-hexen-1-ol acetate, phenyl-ethyl alcohol, *D,L*-limonene (racemic),  $\beta$ -caryophyllene,  $\alpha$ -humulene and  $\beta$ -selinene. Thus, it is concluded that the use of SPME technique on lyophilized powder of the jambo peel is suitable for the identification of its volatile constituents and provides similar results those described in the literature.

Acknowledgments: To IFRJ by financial and institutional support.

1. J A Pino et al (2004). Flavour Frag. J. 19: 32-35.

2. K C Wong and F Y Lai (1996). Flavour Frag. J. 11: 61-66.

3. Library of Mass Spectra NIST.

4. R P Adams (1995). Identification of Essential Oil Components by Gas Chromatography and Mass Spectrometry. Allure Publishing Corporation. Illinois.

**P 132. Scents from Brazilian Cerrado: The essential oil from *Porophyllum angustissimum***

Ferreira R<sup>1</sup>, Rezende CM<sup>1</sup>; Santana HCD<sup>2</sup>; Vieira RF<sup>2</sup>; Alves RBN<sup>2</sup>; Silva MA<sup>3</sup>; Bizzo HR<sup>4</sup>

<sup>1</sup>Instituto de Química, Universidade Federal do Rio de Janeiro;

<sup>2</sup>Embrapa Genetic Resources and Biotechnology; <sup>3</sup>Instituto Brasileiro de Geografia e Estatística;

<sup>4</sup>Embrapa Food Technology

bizzo@ctaa.embrapa.br

Keywords: *Porophyllum angustissimum*; essential oil; Cerrado; Brazilian flora; myrcene; Astereaceae

Cerrado (savannah) is the second largest Brazilian bioma, but the first in number of endemic species and the most threaten by anthropic pressure. A huge amount of non investigated aromatic species make the Cerrado a very promising source for flavour and fragrance applications. Considering this, Embrapa has started a research project to study the aromatic species from this bioma in order to propose sustainable alternatives for their commercial use. *Porophyllum angustissimum* Gardner (Asteraceae) is a herbaceous plant endemic to the Brazilian Cerrado flora, presenting a strong scent [1]. It has brown inflorescences, leaves slightly blue, and occurs on a disperse and small population. Plant material was collected from six individuals in the Ecological Reserve of the Brazilian Institute of Geography and Statistics (IBGE) in Brasilia, on April 2012. Vouchers were deposited on Embrapa Genetic Resources and Biotechnology herbarium (CEN 2419). The dried plant (aerial parts) was extracted in a Clevenger type apparatus for 2 hours. The oil was analyzed by gas chromatography and mass spectrometry using an Agilent 6890GC equipped with a FID and a 5973N MSD. For separation of the oil components a DB-5 (30m X 0.25mm X 0.25 µm) capillary column was used, with either helium (for MS) or hydrogen (for FID) as carrier gas (at 1.0mL/min). Column temperature rised from 60°C to 240°C at 3°C/min. Mass detector was operated in electronic ionization mode at 70eV. For identification, both mass spectra and retention indices were used [2]. The essential oil was obtained in 0.2% yield. The major compounds present were myrcene (40.6%), (*E*)-2-dodecenal (37.5%) and limonene (3.4%).

Acknowledgment: Conselho Nacional de Desenvolvimento Tecnológico (CNPq).

1. Nakajima J (2012) Lista de Espécies da Flora do Brasil. Jardim Botânico do Rio de Janeiro, Rio de Janeiro.
2. Adams RP (2007) Indentification of Essential Oil Components by Gas Chromatography / Mass Spectrometry. 4th Allured, Illinois.

**P 133. A preliminary study of limonene concentration from essential orange oil (*Citrus sinensis*) by using a new distiller prototype.**

Torres-Martínez LG<sup>1</sup>, García-Fajardo JA<sup>1</sup>; Arriola-Guevara E<sup>2</sup>; Suárez-Jacobo Á<sup>1</sup>; Gutiérrez-González P<sup>2</sup>

<sup>1</sup>Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de Jalisco, AC. Unidad Noreste-PIIT. Vía de Innovación 404, Apodaca, Nuevo León 66629, México;

<sup>2</sup>Universidad de Guadalajara, Avenida Juárez No. 976, Guadalajara, Jalisco 44100, México.

*lilia\_gtm@hotmail.com*

Keywords: Prototype, Limonene, wiped film reboiler, surface response

Nowadays, the production of natural essential oils is being intensified due to the wide industrial interest. Orange oil is a complex mixture composed of hydrocarbon terpenes, oxygenated compounds, pigments, waxes, resins and flavonoids. One compound of commercial interest has been Limonene obtained from citrus oil. Limonene is widely used in cosmetics, pharmaceuticals, food products, it is also used as a botanical insecticide. Limonene is oxidized easily and can be distilled without decomposition, however at elevated temperatures it can be degraded. In this sense, it is important to study the beneficial techniques of separation and concentration that allow obtaining more valuable products such as Limonene by using low temperatures. An ICL-04WR laboratory system with wiped film reboiler and reflux column (InCon Processing LLC, Batavia IL, USA) for sensitive materials was used in order to concentrate Limonene. In order to evaluate the temperature conditions, the evaporator temperature was evaluated ranged from 50 to 120°C, the pressure system was ranged from 1.7-10 torr and feed rate was 10 mL/min. Looking at the response surface of the factor evaporator temperature and pressure it is observed that Limonene content increase when temperature increase and the pressure decrease. The equation for Limonene content from the two processing variables was: %Limonene = 89.038-1.87982\*P<sup>2</sup>+0.0947551\*P\*T+0.132182\*P<sup>3</sup>-5.24555\*10<sup>-6</sup>\*T<sup>3</sup> (R<sup>2</sup>=80.11). These preliminary results demonstrate the potential of this new distiller prototype as an alternative distillation process for thermally sensitive compounds.

**P 134. Antifungal activity of essential oils and isolated compounds**

Rebello RA<sup>1</sup>, Begnini IM<sup>1</sup>; Meneghetti EK<sup>1</sup>; Dognini J<sup>1</sup>; Bella Cruz A<sup>2</sup>; Cruz RCB<sup>2</sup>; Zimmermann AC<sup>2</sup>

<sup>1</sup>Department of Chemistry, Universidade Regional de Blumenau - FURB, Blumenau, SC, 89012-900, Brazil;

<sup>2</sup>Health Sciences Center, Universidade do Vale do Itajaí - UNIVALI, Itajaí, SC, 88302-202, Brazil

ricardorebello@furb.br

Keywords: *Drimys angustifolia*; essential oils; myristicin; cyclocolorenone; antifungal assay.

Natural products are fundamental source of chemicals with diverse structures and biological properties, thus providing an important alternative for new antimicrobial agents [1]. This work aim the isolation, characterization and antifungal evaluation of essential oils (EOs) from *Drimys angustifolia* [2] and isolated compounds.

Leaf and branch EOs from *D. angustifolia* were obtained by hydrodistillation under nitrogen atmosphere using a Clevenger type apparatus. The oils were characterized by means of gas chromatography (GC-FID) and gas chromatography coupled to mass spectrometry (GC-MS). The sesquiterpenes cyclocolorenone and bicyclogermacrene, and the arylpropanoid myristicin were isolated from the leaf essential oil by column chromatography on silicagel. Pure hexane and the binary mixture (hexane:dichloromethane) were used as eluents. Drimenol was obtained from the branch essential oil using crystallization in cold hexane. EOs and isolated compounds were assayed against 10 different types of filamentous fungi and yeast with their Minimum Inhibitory Concentration (MIC) being expressed in ppm.

The EOs were active against all the fungi tested, with CIM ranging from 31.25 to 1,000 ppm. The most active myristicin and cyclocolorenone exhibited MIC values of 15.6-500 ppm and 62.5-1,000 ppm, respectively, being the activity of the former consistent to the literature [3]. Bicyclogermacrene was inactive against 6 of the assayed fungi. *E. floccosum*, *C. neoformans* and *M. gypseum* were the most sensitive microorganisms.

The EOs from *D. angustifolia* and isolated compounds exhibited promising antifungal activity. Myristicin and cyclocolorenone embody structural features enabling to access semi-synthetic derivatives for further antifungal screening.

Acknowledgments: FURB, UNIVALI, FAPESC, INCT catalysis.

1. M Saleem et al. (2010) Nat. Prod. Rep. 27: 238-254.

2. RC Forzza et al. (2010) Flora do Brasil. Jardim Botânico RJ, Rio de Janeiro.

3. A Maxis et al. (2012) Mycopathologia (Epub).

**P 135. Antidermatophytic and antileishmanial activities of essential oil from *Lippia gracilis* genotypes**

Melo JO<sup>1</sup>, Farias Junior PA<sup>2</sup>; Bitencourt TA<sup>3</sup>; Blank AF<sup>1</sup>; Fernandes RPM<sup>4</sup>; Fachin AL<sup>3</sup>; Scher R<sup>5</sup>; Cruz EMO<sup>6</sup>; Jesus HCR<sup>7</sup>; Alves PB<sup>7</sup>; Arrigoni-Blank MF<sup>6</sup>; Franca SC<sup>3</sup>; Belebony RO<sup>3</sup>

<sup>1</sup>Departamento de Engenharia Agrônômica, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil.;

<sup>2</sup>Departamento de Fisiologia, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil.;

<sup>3</sup>Unidade de Biotecnologia, Universidade de Ribeirão Preto, Ribeirão Preto, São Paulo, Brazil;

<sup>4</sup>Departamento de Fisiologia, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil.;

<sup>5</sup>Departamento de Morfologia, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil.;

<sup>6</sup>Departamento de Engenharia Agrônômica, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil.;

<sup>7</sup>Departamento de Química, Universidade Federal de Sergipe, Cidade Universitária, CEP 49100-000, São Cristóvão, Sergipe, Brasil

[scher@ufs.br](mailto:scher@ufs.br)

Keywords: *Lippia sidoides*; *Trichophyton rubrum*; *Leishmania chagasi*; thymol; carvacrol

The aim of this work was to evaluate the antidermatophytic and antileishmanial activities of essential oil from *Lippia gracilis* genotypes. The leaves of three *L. gracilis* genotypes, including LGRA-106, LGRA-109, and LGRA-110 were collected from the Active Germplasm Bank located in the "Campus Rural da UFS" Research Farm at the São Cristovao County, Sergipe State, Brazil. The essential oils were obtained by hydrodistillation and their constituents were characterized by gas chromatography coupled to mass spectrometry (GC/MS). The susceptibility of *Trichophyton. rubrum* strains, MYA3108 and *TruMDR2* to essential oils from LGRA-106 and LGRA-109 was determined by serial microdilution method. The leishmanicidal activity of essential oil from LGRA-106 and LGRA-110 was assayed on promastigotes of *Leishmania chagasi* and cell viability was determined by MTT method. The oxygenated monoterpene thymol was the main component of the essential oil from LGRA-106 (59.26%) while carvacrol was more abundant in LGRA-109 and LGRA-110 (54.56 and 48.92%, respectively). The *L. gracilis* essential oils evaluated in this study presented antifungal and leishmanicidal activities. However, LGRA-106 essential oil exhibited the best fungicidal activity at a concentration of 46.87 µg/mL. Both, LGRA-106 and 110 essential oils had similar antileishmanial effect, showing IC<sub>50</sub> of 96.34 nL/mL and 86.23 nL/mL, respectively. The results obtained in this study suggest that essential oil of *L. gracilis* is a potential source of thymol and carvacrol and a promising antimicrobial agent, mainly fungicidal agent, since the concentrations that completely eliminate the fungi tested, were comparable to those observed for the common antifungal drug fluconazole.

Acknowledgments: The authors thank FAPITEC/SE, CNPq and CAPES for their financial support of this work.



**P 136. Chemical analysis of *Hypochaeris maculata* ssp. *pelivanovicii* (Velen.) Hayek essential oil**

Jelena Lazarevic<sup>1</sup>, Aleksandra Djordjevic<sup>2</sup>; Bojan Zlatkovic<sup>3</sup>; Gordana Stojanovic<sup>2</sup>

<sup>1</sup>Department of Chemistry, Faculty of Medical Sciences, University of Niš, Bul. dr Zorana Đinđića 81, 18000 Niš (Serbia);

<sup>2</sup>Department of Chemistry, Faculty of Science and Mathematics, University of Niš, Višegradska 33, 18000 Niš (Serbia);

<sup>3</sup>Department of Biology and Ecology, Faculty of Science and Mathematics, University of Niš, Višegradska 33, 18000 Niš (Serbia)

jelena217@yahoo.com

Keywords: *Hypochaeris maculata* ssp. *pelivanovicii*, Asteraceae, essential oil

Cat ear, the common name for a number of annual and perennial herbs from the genus *Hypochaeris* (Asteraceae, Cichorieae), are in many areas considered a weed. Phytochemical data related to the genus are scarce, including a small scale investigation on sesquiterpen lactones (germacranolides, guaianolides and eudesmanolides) [1] and on flavonoids [2].

The above ground parts of Balkan endemics *Hypochaeris maculata* ssp. *pelivanovicii* were collected in July 2007, from highland meadows (natural populations), on Mountain Stara Planina (East Serbia). The composition of hydrodistilled essential oil was investigated by means of GC and a GC-MS analysis.

The analysis resulted in the identification of one hundred twenty five components, accounting for 89.1% of the total oil. Main identified constituents were hexadecanoic acid (14.0%), heneicosane (10.9%), nonacosane (10.4%) and tricosane (7.2%). Fatty acid derived compounds were predominant compound class in the oil (70%), while terpenoids comprised only 8.9% of the compounds detected.

To the best of authors' knowledge, this would be the first report on essential oil chemistry of genus *Hypochaeris* in general.

C. Zidorn (2006) *Biochem. Syst. Ecol.* 34: 144-159.

V. Sareedenchai, C. Zidorn (2010) *Biochem. Syst. Ecol.* 38: 935-957.

**P 137. Chemical composition and biological activities of ten different species of the Asteraceae family from Nepal**

Prabodh Satyal<sup>1</sup>, Prajwal Paudel<sup>1</sup>; Ambika Poudel<sup>2</sup>; Katherine E. Woods<sup>1</sup>; Dr. William N. Setzer<sup>1</sup>

<sup>1</sup>Department of Chemistry, University of Alabama in Huntsville, AL-35899, USA;

<sup>2</sup>Department of Chemistry, Tribhuvan Univ. Biratnagar, Nepal

ps0013@uah.edu

Keywords: Asteraceae, bioactivities, chemodiversity, essential oil

There are more than 22750 species of plants belonging to the Asteraceae family world wide. Within the flavor and fragrance industry, Asteraceae family is considered to be a major volatile oil source for commercial usage. In our study ten species from Nepal were analyzed for leaf essential oil. The species included: *Artemisia dubia*, *Artemisia vulgaris*, *Matricaria chamomilla*, *Blumea lacera*, *Eupatorium adenophorum*, *Chrysanthemum cinerifolium*, *Parthenium hysterophorus*, *Xanthium strumarium*, *Ageratum conyzoides*, and *Artemisia indica*. Essential oil compositions were determined using an Agilent 6890 GC coupled with Agilent 5973 mass selective detector. The chemical composition for each essential oil were compared within the genera and family. Further analysis of biological activity was accomplished via various bio-assays including: cytotoxicity, anti-microbial, larvicidal, brine shrimp lethality, nematocidal activity, and insecticidal activity. Chemical compositional analysis showed taxonomic variation as well as geographical and climatic influences on the essential oil chemotype among the species. Biological testing showed wide variation in activity. Through our research, deeper understanding of chemodiversity and species differentiation within the Asteraceae family has been developed. In addition, bio-assay results have shown potential medicinal value of the species. Out of those oils *Matricaria chamomilla* and *Artemisia indica* has shown strong evidence to be therapeutically important essential oils for parasites and microbiological organisms. *Artemisia vulgaris* and *Parthenium hysterophorus* has shown strong evidence for allelopathy and insecticidal activities. Overall, this study has been a strong evidence for Nepal as a promising source of natural products.

Acknowledgments: We would like to express our gratitude to Tilak Gautam for identification of plants and to Dr. Bernhard Vogler for technical assistance.

1. P Satyal et al. (2012) J. Chem. Pharmaceut. Res. 4:437-439.
2. P Satyal et al. (2012) J. Essen. Oil Bearing Plants 15: (Accepted).
3. WN Setzer (2009) Nat. Prod. Commun. 4:1305-1316.
4. P Satyal et al. (2012) Phamacog. Res. 4:(Accepted).

**P 138. Analysis of sandalwood essential oil by GC-MS with ionic liquid stationary phases**

Cagliero C<sup>1</sup>, Bicchi C<sup>2</sup>; Cordero C<sup>2</sup>; Liberto E<sup>2</sup>; Sgorbini B<sup>2</sup>; Rubiolo P<sup>2</sup>

<sup>1</sup>Laboratory of Phytochemical Analysis, Dipartimento di Scienza e Tecnologia del Farmaco University of Torino;

<sup>2</sup>Laboratory of Phytochemical Analysis, Dipartimento di Scienza e Tecnologia del Farmaco University of Torino,

patrizia.rubiolo@unito.it

Keywords: *Santalum* spp.; Santalols; Farnesols; GC-MS; Ionic liquid stationary phases

Commercial sandalwood essential oils are obtained by hydrodistillation of the heartwood of 20-year-old trees mainly from three *Santalum* species (family Santalaceae): a) *Santalum album* L., also known as 'East Indian', (Indonesia, Sri Lanka, Northern Western Australia), b) *S. austrocaledonicum* Vieill., (New Caledonian and Vanuatu archipelagos), c) *S. spicatum* (R. Br.) A. DC, (Southwestern Australia).

The Indian type is the most considered in perfumery because of its organoleptic properties. Quality control (QC) of *S. album* and *S. spicatum* essential oils are nowadays regulated with international norms [1,2] requiring a primary free alcohol content, expressed as santalol, not lower than 90%, with (*Z*)- $\alpha$ -santalol ranging from 41 to 55% and (*Z*)- $\beta$ -santalol from 16 to 24%. Analysis of sandalwood essential oil has recently been reviewed by Baldovini et al [3].

Regulatory rules require that (*E,E*)- $\alpha$ -farnesol, a minor sandalwood component suspected to be a fragrance allergen, is quantified; unfortunately, this alcohols cannot be base-line separated in a single GC run, since it co-elutes with (*Z*)- $\beta$ -santalol on non-polar columns, and with (*Z*)- $\alpha$ -santalol on polar columns. Multidimensional GC methods including GCxGC and MDGC have therefore successfully been developed [4 – 6].

New perspectives have recently been opened up in GC by the introduction of ionic liquids as stationary phases because of their selectivity towards specific chemical classes that are completely different from those of the conventional phases.

This communication reports the results of a study dealing with the simultaneous separation of farnesol and santalol isomers in a single GC run and (*E,E*)- $\alpha$ -farnesol quantitation in a set of sandalwood essential oils from different species by GC-MS with an ionic liquid stationary phase column (Supelco – IL 60)

1. ISO-FDIS 3518 (2001)

2. ISO 22759(2009)

3. N Baldovini, C Delasalle, et al. (2011) Flavour Fragr. J., 26: 7–26

4. R Shellie, P Marriott, et al., (2004) J. Chromatogr. Sci. 42: 417.

5. A Klamecki, H Brévard, et al. (2006) Proc. of 37th International Symposium on Essential Oils, Grasse, Sept. 10–13, p.168.

6. D Sciarone, R Costa, et al. (2011) J. Chromatogr. A, 1218 137–142

**P 139. Control of the pinewood nematode *Bursaphelenchus xylophilus* by essential oils and extracts obtained from plants: a review.**

Barbosa P<sup>1</sup>, Vieira P<sup>1</sup>; Dias LS<sup>2</sup>; Pedro LG<sup>3</sup>; Barroso JG<sup>3</sup>; Figueiredo AC<sup>3</sup>; Mota M<sup>1</sup>

<sup>1</sup>NemaLab-ICAAM & Departamento de Biologia, Universidade de Évora, 7002-554 Évora, Portugal;

<sup>2</sup>Departamento de Biologia, Universidade de Évora, 7002-554 Évora, Portugal.;

<sup>3</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portugal

pmbarbosa@yahoo.com

Keywords: *Bursaphelenchus xylophilus*; essential oils; extracts; nematicides; plants

The pinewood nematode (PWN), *Bursaphelenchus xylophilus*, is a serious threat to forest ecosystems at a global scale. The nematode has become a major quarantine problem due to its capability to completely destroy *Pinus* spp. trees, with great damage to the wood industry. Controlling the nematode inside a living tree is quite difficult, the techniques used being often ineffective and quite expensive. In the coming years, most chemicals used to control nematodes will be banned and replaced by safer and environmentally friendly products. As so, chemicals naturally produced by plants will play an important role in controlling diseases such as pine wilt. Plants, particularly aromatic ones, are commonly used due to the chemical properties of their secondary metabolites. Among these, essential oils and/or extracts are highly employed and are being tested as possible control of some organisms, like nematodes. Recent publications have evaluated essential oils derived from different plant species as natural nematicides [1; 2], anti-bacterial [3], anti-fungal [4] as well as insecticidal [5]. Concerning control of the PWN, a significant amount of information on plants tested, results obtained and employed techniques, is available. Our revision has extensively gathered this information, making it easier to search, read and use. It may become useful information for future studies on the subject, since it will be possible to check the plants already tested. Although numbers aren't definitive, so far, tested plants are distributed amongst 148 families. The extracts or essential oils of plants belonging to the Asteraceae, Lamiaceae and Euphorbiaceae families show promising results on controlling the pinewood nematode.

Acknowledgments: Studies partially funded by FCT, under research contracts PTDC/AGR-CFL/117026/2010 and PTDC/AGR-CFL/120184/2010.

1. P Barbosa et al. (2010) *J Nematol.* 42: 8-16.
2. E Kim et al. (2011) *Nematology* 13(3): 377-380.
3. S Prabuseenivasan et al. (2006) *BMC Complementary Altern. Med.* 6: 39.
4. S Lee et al. (2007) *Plant Pathol. J.* 23(2): 97-102.
5. G Tarelli et al. (2009) *J Econ Entomol.* 102(3): 1383-1388.

**P 140. Volatiles from the edible mushroom *Lepista nuda*: comparison between wild and commercial samples**

Pinto S<sup>1</sup>, Sousa MJ<sup>1</sup>; Figueiredo AC<sup>2</sup>; Barroso JG<sup>2</sup>; Pedro LG<sup>2</sup>

<sup>1</sup>Instituto Politécnico de Bragança, Escola Superior Agrária, Departamento de Biologia e Biotecnologia; Cimo. IPB;

<sup>2</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal

sarapinto109@hotmail.com

Keywords: *Lepista nuda*; mushrooms; volatiles; linalool; pulegone

*Lepista nuda* (Bull. ex Fr.) Cooke, sometimes given the common name "blewit", is a *Basidiomycete*, from the *Tricholomataceae*, *Agaricales*. It is asaprotrophic species; growing alone, gregariously, or in clusters in organic debris, in woods or in urban settings. The fruity flavour makes it an excellent edible mushroom and it is commercially grown in Britain, France and the Netherlands [1].

Wild mushroom species showed to be less energetic than the commercial ones, with higher contents of protein and lower fat concentrations [2]. Mushrooms volatile compounds are less known although they play an important role for the flavour and organoleptic properties.

Wild *L. nuda* (Blewit) fruit bodies were collected from populations at two different habitats in Northeast Portugal: oak forest (OF) and pine forest (PF), and were also obtained from a commercial source (CS). Volatiles were isolated by distillation-extraction and analyzed by Gas Chromatography and Gas Chromatography-Mass Spectrometry as in [3].

In total, twenty two components were identified in each of the three samples, attaining from 84-94% of the volatile fraction. Differences between volatiles from wild growing and commercial samples were mainly quantitative. Linalool (17-26%), pulegone (12-14%) and limonene (10-11%) were the three major components in all samples. The main difference was observed in the percentage of 2-pentyl furan, present in small amount in the wild growing mushrooms (2%, OF and 5%, PF) and in considerable amount in the commercial sample (15%), where it was the second major component.

Eight-carbon (C8) volatiles are ubiquitous among fungi and characteristic of the fungal aroma, 1-octen-3-ol being the most abundant [4]. In the present study, 1-octen-3-ol was detected in small amounts in all samples (traces-2%) and the major components, linalool, limonene and pulegone, are not common in other mushroom volatiles.

Acknowledgments: Supported by FCT, COMPETE/QREN/UE project PTDC/AGR-ALI/110062/2009

1. IR Hall *et al* (2003) *Edible and Poisonous Mushrooms of the World*. Timber Press Inc., pp: 180-181.

2. FS Reis *et al* (2011) *Molecules*, 16, 4328-4338.

3. MD Mendes *et al*. (2011) *Ind. Crops Prod.* 33: 710-719.

4. E Combet *et al* (2006) *Mycoscience*, 47: 317-326.

**P 141. The way to ensure high quality aroma of decontaminated spices**

Elżbieta Wojtowicz<sup>1</sup>, Renata Zawirska-Wojtasiak<sup>2</sup>; Janusz Adamiec<sup>3</sup>

<sup>1</sup>Department of Food Concentrates in Poznan, Institute of Agricultural and Food Biotechnology, Poland, 61-361 Poznan, ul. Starolecka 40.

<sup>2</sup>Faculty of Food and Nutrition Science, Poznan University of Life Sciences, Poland, 60-637 Poznan, ul. Wojska Polskiego 2.

<sup>3</sup>Faculty of Process and Environmental Engineering, Technical University of Lodz, Poland, 90-924 Lodz, ul. Wolczanska 213.

renazaw@up.poznan.pl

Keywords: spices, aroma, key odorants, microencapsulated preparations

Spices become contaminated microbiologically during harvesting, drying and storage. The addition of contaminated spices to foodstuffs may be dangerous for the health of consumers. For this reason several methods of spices decontamination are used. All known and used methods of decontamination lead to a greater or lesser losses of aroma of leafy spices. The aim was provide high quality leafy aroma of decontaminated spices.

Materials: marjoram, thyme, microencapsulated powders of linalool, sabinene hydrate, thymol.

Methods: steam decontamination of spices at the Institute of Agricultural and Food Biotechnology in Poznan. Steam distillation: in a Deryng' apparatus. 1 g sample with 0.5 ml o-xylene. Internal standard was tetradecane. Distillation was run for 3 h. GC/MS: 5975C VL MSD with HP-5MS (30 m x 0.25 mm x 0.25 µm) (Agilent Technologies). Library NIST05. GC/O-AEDA: HP 5890 with an inlet splitter and a smelling port, a DB-5 (30 m x 0.53 mm x 0.25 µm). Separated fractions were smelled in successive double dilutions of analyzed spice distillates, until the last detectable aroma disappeared. SPME/PDMS: samples 100 mg, vials 10 ml, addition of 2.5 ml deionized water and 300 µl methanol. Extraction temperature 50°C, time 15 min. OAV: concentration of compounds / OT (odor threshold). Profile sensory analysis: 10 experts, 8 attributes for marjoram, 6 for thyme. Obtained data were interpreted by Principal Component Analysis. Preparing spices with microencapsulated powders of key odorants: mixer Mesko R-1000.

Results: Volatile aromatic compounds were identified in spices. Quantitative composition of these compounds was determined before and after spices decontamination. The GC/O analysis and the determination of odor activity values (OAV) showed that the primary compounds responsible for the aroma in analyzed spices were hydrate cis-sabinene and linalool in marjoram and thymol in thyme. It was found that the applied steam decontamination method resulted in relatively high losses of aroma. In order to supplement these losses of aroma compounds caused by decontamination, attempts were made to supplement their key odorants in the form of microencapsulated preparations. Results of chromatographic and sensory analyses of such prepared spices made it possible to state that an adequate addition of microencapsulated preparations of main odorants to decontaminated spices eliminated differences between decontaminated and control spices.

**P 142. Effect of the speed of the drying air on the quality of essential oil from *Aristolochia cymbifera* Mart. & Zucc.**

Montes AL<sup>1</sup>, Oliveira DEC<sup>1</sup>; Resende O<sup>2</sup>; Silva FG<sup>2</sup>; Guerreiro MC<sup>3</sup>; Sales J F<sup>2</sup>

<sup>1</sup>Student of the Graduate Program in Agricultural Sciences at IF Goiano;

<sup>2</sup>Professor at IF Goiano;

<sup>3</sup>Department of Chemistry, Federal University of Lavras

fabianocefetriv@yahoo.com.br

Keywords: Medicinal plants, Chemical composition, Content of essential oil, Extraction

The species *Aristolochia cymbifera* Mart. & Zucc., popularly known as “jarrinha”, “milhomem”, or “cassau”. This species is an herbaceous perennial vine native to Brazil that is characteristically vigorous and better adapted to hot environments.<sup>3</sup> According to these authors, *A. cymbifera* contains mono/diterpenes and sesquiterpenoids in the leaves, stems, and roots. In folk medicine, *A. cymbifera* is used for various problems and is considered to be a diuretic, antiseptic, and antispasmodic.<sup>3</sup>

The object of the present study was to evaluate the effect of three drying air speeds on the content and the chemical composition of essential oil from *A. cymbifera* Mart. & Zucc. The drying was conducted in a fixed-layer dryer manufactured from #16 metal sheets. The drying chamber measured 0.60 x 0.60 x 0.60 m, for a total volume of 0.216 m<sup>3</sup>, and contained a plate with 25 % perforation placed at a height of 0.33 m. The fan was of the centrifugal type, driven by a three-phase motor with a power of 1.5 HP and rotation at 1,720 rpm. Each dryer was composed of six swinging temperature sensors and four electrical resistors of 1,500 ohms, for a total of 6,000 ohms. The system also featured an automatic controller that managed the system and stored the data generated. The system was set to 34.7±1.5 °C with controlled air speeds of 0.5, 1.0, and 2.0 m·s<sup>-1</sup>. The essential oil was extracted by hydrodistillation. The chemical analyses were performed on a gas chromatography apparatus coupled to a Shimadzu QP5050A quadrupole mass spectrometer (GC-MS) (Kyoto, Japan).

Increasing the speed from 0.5 m·s<sup>-1</sup> to 2 m·s<sup>-1</sup> reduced the drying time from 22 to 16 h. The speed of the drying air did not influence the content of essential oil extracted from *A. cymbifera* leaves. The minor constituents germacrene, hex-2-enal, viridiflorol, and cedrol were influenced by the drying process. The major constituents, mainly spathulenol, were not influenced by the drying air speed.

1. Lorenzi, H.; Matos, F. J.A. (2002). Plantas medicinais no Brasil nativas e exóticas. Nova Odessa: Instituto Plantarum. 512 p. pg 77.

**P 143. Chemical synthesis and microbial transformation of biologically active derivatives of  $\beta$ -damascone**

Gliszczyńska Anna<sup>1</sup>, Gediga Jan<sup>1</sup>; Gładkowski Witold<sup>1</sup>; Wawrzeńczyk Czesław<sup>1</sup>

<sup>1</sup>Wroclaw University of Environmental and Life Sciences

*anna.gliszczynska@wp.pl*

Keywords: damascone; biotransformation; lactones; natural compounds; norisoprenoids

Terpene lactones are a large group of compounds with noteworthy biological and therapeutic activities [1,2]. They occur in plants and microorganisms. Some of them were obtained also in the chemical synthesis. Recently we have published the synthesis of isoprenoid lactones, which exhibit interesting antifeedant activity [3]. Here we report the results of chemical synthesis of lactones from cyclic natural ketone –  $\beta$ -damascone. We expect that they will show antifungal and antifeedant activity and interesting odoriferous properties [4]. Our expectations arise from the fact that  $\beta$ -damascone itself possesses important valuable odour and has been identified as component of tea aroma. It possesses fruity, roselike odour and is used in perfume compositions. Besides, it is tested as potential insecticidal agent [5,6,7]. In our studies we focused on the chemical synthesis of lactone derivatives of  $\beta$ -damascone and microbial transformations of obtained products using fungal cultures.

Acknowledgements: This Project was financed by European Union from the European Regional Development Found Grant No. POIG.01.03.01-00-158/09-03.

1. Marles, R.J. et al. (1995) *Phytochemistry of Medicinal Plants* 29, pp 333-356.
2. Robles, M. et al. (1995) *Planta Med.* 61, pp 199-203.
3. Gliszczyńska A. et al. (2011) *Przem. Chem.* 90, 5, pp 759-763.
4. Kerrigan, N.J. et al. (2004) *Tetrahedron Lett.* 45, pp 9087-9090.
5. Bauer K. et al. (1997) *Common Fragrance and Flavour Materials*, New York.
6. Schoch E. (1991) *Appl. Environ. Microb.* pp 15-18.
7. Kaufman P.E. et al. (2011) *Pest Manag Sci.* 67, pp 26-35.



**P 144. Effects of the essential oil of *Alpinia zerumbet* leaves in muscle activity in spastic gastrocnemius of patients with stroke**

Kamyla Daniely Lima Cardoso<sup>1</sup>, Natalia Miranda Batista dos Santos<sup>1</sup>; Sheyla Alves Rodrigues<sup>1</sup>; Lauro Xavier-Filho<sup>2</sup>; Edna Aragão Farias Cândido<sup>1</sup>

<sup>1</sup>*Institute of Technology and Research, Tiradentes University, Brazil;*

<sup>2</sup>*Teacher of the Rede Nordeste de Biotecnologia, Brazil*

*shelrodrigues@hotmail.com*

Keywords: Electromyography; stroke; spastic muscle

Stroke occurs due to a disturbance in focal cerebral function, secondary to obstruction or bleeding blood vessels<sup>1</sup>. One of the sequels is the hemiplegia, which reaches the lower limb spasticity associated involving Extenders<sup>2,3</sup>. To investigate the effects of the essential oils of *Alpinia zerumbet* leaves in muscle activity in the medial and lateral gastrocnemius of patients with stroke using electromyography. This is a prospective and analytical study of clinical trial I, realized with six patients with stroke associated with hemiparesis. The spastic gastrocnemius muscle activity of each subject was assessed before and after application of the essential oil of *A. zerumbet*, compared by electromyography. The treatment was realized with 0,10 mL/kg dose of the oil associated to 10 applications of physiotherapy. The analysis of the gastrocnemius muscle activity before and after treatment showed a significant improvement in muscle recruitment ( $p < 0.0087$ ) in the medial portion in all patients, while the lateral portion of the muscle showed a slight recruitment was not statistically significant ( $p < 0.0511$ ). Treatment with essential oil of *A. zerumbet* associated with physiotherapy improved the motor activity in spastic gastrocnemius in stroke patients.

Acknowledgments: We thank the Hebron Pharmaceuticals LTDA for the financial support of research.

1. Patten C, Lexell J, Brown HE. (2004) *Journal Rehabilitation Research Development*. 41(3): 293-312.
2. Lianza S. (2001) *Medicina de Reabilitação*. 3 ed. Rio de Janeiro: Guanabara Koogan. 462p.
3. Cargnin APM, Mazzitelli C. (2003) *Revista Neurociências*, 11(1): 34-9.

**P 145. Chemical composition and biological activity of essential oil of *Coffea arabica* L. In the larvae of *Aedes aegypti* and *Artemia salina***

Sheyla Alves Rodrigues<sup>1</sup>, Clisiane Carla de Souza Santos<sup>2</sup>; Américo Azevedo de Souza<sup>2</sup>; Péricles Barreto Alves<sup>3</sup>; Darlisson de Alexandria Santos<sup>3</sup>; Charles dos Santos Estevam<sup>2</sup>

<sup>1</sup>Institute of Technology and Research, Tiradentes University, Brazil;

<sup>2</sup>Department of Physiology, Federal University of Sergipe, Brazil;

<sup>3</sup>Federal University of Sergipe - Brazil

shelrodrigues@hotmail.com

Keywords: Dengue; toxicity; volatile compounds

*Aedes aegypti* L. is the vector responsible for transmitting the viral infection (dengue)<sup>1</sup>. The most effective proven method for disease prevention has been by vector control by chemical or biological means<sup>2</sup>, but chemical insecticides resistance has already been observed in some insect populations in the Brazil<sup>3</sup>. In this regard, we have investigated the effectiveness of compounds extracted from plants in the control of development of *A. aegypti* larvae stages of development and that are free of toxicity to the environment. This work aimed study the chemical composition, toxicity and larvicidal activity of essential oil of *C. arabica*, in the search of new methods to control dengue.

The essential oil of *C. arabica* was extracted by hydrodistillation and submitted quantitative analysis of chemical constituents that was performed by gas chromatography flame ionization (FID) using a Shimadzu GC-17A. Bioassays followed proposed standards by the WHO<sup>4</sup>, using dosages of 1000, 500, 250, 125, 62.5 mg/mL to calculate the lethal concentration (CL<sub>50</sub>) compared with negative and positive control (distilled water with temephos). The evaluation of toxicity in *Artemia salina* was performed according methodology by McLaughlin<sup>5</sup>. The calculation of the CL50% was estimated using the probit analysis method (Minitab software, version 15), with 95% confidence interval.

The components  $\alpha$ -furfuryl alcohol (24.22%), 5-metilfurfural (10.15%) and furfural (21.9%) were characterized as majoritary in the essential oil of *C. arabica*. The essential oil not showed larvicidal effect on *A. aegypti*, causing mortality at a concentration of 1000 ppm after 48 h of exposure, different of studies with caffeine, which showed results with concentrations of 200 and 500  $\mu$ g/mL<sup>6</sup>. The bioassay toxicity in *A. salina* was found an CL<sub>50</sub> of 358 mg / mL, indicating that low toxicity oil. In conclusion, the essential oil of *C. arabica* not proved to be an effective larvicidal to *A. aegypti*, but it has low toxicity.

1. Mendonça FA, Souza AV, Dutra DA. (2009) Sociedade & Natureza, 21(3):257-269.

2. Lingon BL. (2005) Semin Pediatr Infect Dis, 16:60-65.

3. Barreto CF. (2005) Revista Eletrônica Faculdade Montes Bolos, 1(2):62-73, 2005.

4. WHO. Guidelines for laboratory and field testing of mosquito larvicides.2005.

5. McLaughlin J. L. (1998) Drug Inf J, 24:32-24.

6. Laranja AT, Manzatto AJ, Bicudo HEMC. (2006) Rev Saúde Pública, 40(6):1112-1117.

**P 146. GC/MS analysis of essential oils from the flowers of some *Cirsium* species**

Kozyra M, Głowniak K; Kochmańska J

Chair and Department of Pharmacognosy with Medicinal Plant Unit, Medical University of Lublin, 1 Chodźki St., 20-093 Lublin, Poland

[mkozyra@pharmacognosy.org](mailto:mkozyra@pharmacognosy.org)

Keywords: *Cirsium*, volatile oils

The aim of this study was the phytochemical investigation of essential oils from *Cirsium pannonicum*, *Cirsium decussatum* Janka., *Cirsium ligulare* Boiss., *Cirsium heterophyllum* L. (Chill.), *Cirsium eriophorum* L. (Scop.), *Cirsium hispanicum* (Lam.) Lag, *Cirsium rivulare* (Jacq.) All., *Cirsium erisithales* (Jacq.) All., *Cirsium acaule* (L.) Scop., *Cirsium ferox* (L.) CD flowers by GC/MS method.

In this study, air dried and powdered flowers were used for essential oil distillation, which was carried out in the Deryng apparatus using the method presented in the Polish Pharmacopoeia (6<sup>th</sup> ed.). GC-MS analysis was performed with a Thermo-Finnigan (USA) GCQ GC-MS spectrometer, using the electron impact (E) ionization mode 70 eV. Full-scan mass spectra were recorded in the range  $m/z$  35–500 a.m.u. The identification of individual compounds was based on the comparison of mass spectra obtained from the samples with those of authentic standards, those of the NIST (National Institute of Standards and Technology, Gaithersburg, MD, USA) libraries, those of our own library, and related literature [1]. The compounds were also identified by the comparison of GC retention indices and retention times.

Forty one constituents were identified in the essential oils obtained from the flowers of *Cirsium* species. They were qualitatively and quantitative examined.

Volatile oils contained: thymol, lauric and tetradecanoic acids. Thymol was the predominant constituent identified in flowers of *C. rivulare* (24,5%) *C. pannonicum* (8,8%), *C. decussatum* (5,2%), *C. ligulare* (2,4%). Lauric acid is a predominant constituents in *C. pannonicum* (17,3%), *C. rivulare* (11,1%), *C. hispanicum* (6,8%), *C. eriophorum* (6,2%).

1. Adams R.P.: Identification of Essential Oil Components by Gas Chromatography/Mass Spectroscopy, Allured Publishing Corp., Carol Stream, IL, USA, 2001.

**P 147.  $\beta$ -Myrcene biotransformation by *Pseudomonas* sp. M1**

Rodrigues AM<sup>1</sup>, Roma-Rodrigues C<sup>2</sup>; Santos PM<sup>3</sup>; Sá-Correira I<sup>2</sup>; Figueiredo AC<sup>1</sup>; Pedro LG<sup>1</sup>; Barroso JG<sup>1</sup>

<sup>1</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portugal; <sup>2</sup>Instituto de Biotecnologia e Bioengenharia, CEBQ, Instituto Superior Técnico, Universidade Técnica de Lisboa, Av. Rovisco Pais, 1049-001 Lisboa, Portugal; <sup>3</sup>Centro de Biologia Molecular e Ambiental (CBMA), U. Minho, Campus de Gualtar, 4710-087 Braga, Portugal

amargaridacrodriques@gmail.com

Keywords:  $\beta$ -myrcene; *Pseudomonas*; biotransformation

There are two ways to produce compounds in biotechnology: de novo synthesis and biotransformation. The advantage of biotransformation, in comparison with chemical synthesis, is the reduced environmental impact, either by decreasing the amount of wastes or by their easier elimination [1]. Due to the proved important biological activities of some volatile terpenes, the biotransformation of lead compounds within this group can raise new commercially valued products for industry with enhanced biological activities.  $\beta$ -Myrcene, with validated analgesic and antimicrobial activities, besides its importance as intermediate for the production of a wide variety of flavourings and beverages [2], appears as a paramount target. As part of a broader research project towards the metabolic engineering of  $\beta$ -myrcene pathway of *Pseudomonas* sp. M1, in this work we report the preliminary results on de biotransformation of  $\beta$ -Myrcene by the parental strain of *Pseudomonas* sp. M1.

*Pseudomonas* sp. M1 was grown in liquid mineral medium supplemented with  $\beta$ -Myrcene as sole carbon source. Samples for metabolites isolation and identification were collected along the exponential growth phase. Bacteria were separated from the culture medium and the biotransformation volatile compounds were isolated by distillation-extraction and analyzed by Gas Chromatography (GC) and Gas Chromatography-Mass Spectrometry (GC-MS) as in [3]. The crude volatiles, dissolved in pentane, were extracted with aliquots of a 5% sodium bicarbonate solution (acid fraction) prior to being pre-fractionated by liquid-solid chromatography on a silica gel column. Elution was performed with distilled *n*-pentane (hydrocarbon fraction) followed by dichloromethane (oxygen-containing fraction). *E*-Myrcenol and *E*-Myrcenal were two of the main components of the oxygen-containing fraction whereas *E*-Myrcenoic acid occurred in trace amounts in the acid fraction. The occurrence of these compounds in the volatiles isolated from *Pseudomonas* sp. M1 cultures strongly support the suggested catabolic pathway of  $\beta$ -Myrcene by this strain [4].

Acknowledgements: This work is funded by FCT through project ref. PTDC/EBB-BIO/104980/2008.

[1] A Liese (2002) In: K Drauz, H Waldmann (Eds.) Enzyme Catalysis in Organic Synthesis: A Comprehensive Handbook. Wiley-VCH, New York, pp 1419-1459.

[2] M Miyazawa, T Murata (2000) J. Agric. Food Chem. 48:123-125.

[3] JMS Faria *et al.* (2009) Biotechnol. Lett. 31: 897-903.

[4] S Iurescia *et al.* (1999) Appl. Environ. Microbiol. 65: 2871-2876.

**P 148. Greenhouse production of pineapple [*Ananas comosus* (L.) Merr.] in S. Miguel (Azores, Portugal): production of volatiles from traditional growing beds and their effect on soil microorganisms**

Figueiredo AC<sup>1</sup>, Pedro LG<sup>1</sup>; Barroso JG<sup>1</sup>; Domingos C<sup>2</sup>; Carolino M<sup>2</sup>; Cruz C<sup>2</sup>

<sup>1</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portugal;

<sup>2</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Centro Biologia Ambiental C2, Campo Grande, 1749-016 Lisboa, Portugal

acsf@fc.ul.pt

Keywords: *Ananas comosus*, leiva, *Pittosporum undulatum*, *Cryptomeria japonica*, volatiles, antibacterial

Producing 1483 t in 2010 [1], Portugal ranks 13th among the world's pineapple exporters [2]. Traditional pineapple cultivation [Protected Designation of Origin] in the Azores used a substrate composed of the topsoils from uncultivated fields, locally known as *leiva*, and mulch from branches of incense (*Pittosporum undulatum* Vent.) and *Cryptomeria japonica* (Lf) D. Don. Decline in *leiva* soils has stimulated the search for alternative substrates [3,4]. As part of this search, this work reports on the composition of volatiles from the components of the traditional substrate and their effects on soil microorganisms.

The volatiles were isolated by hydrodistillation, and analysed by GC and GC-MS [5]. The antibacterial properties of the volatiles were studied using common soil bacteria and evaluated by the agar diffusion method [6].

*Leiva* is a complex mixture of bryophytes, lichens and mosses, as well as various herbaceous species, thus presenting a high diversity of volatile compounds. The main volatile from *leiva* was hinesene (15-26%). *P. undulatum* essential oil was dominated by limonene (62%). *C. japonica* samples essential oil was dominated by  $\alpha$ -pinene (28%) and phyllocladene (7-16%). Agar disk diffusion tests showed that the three types of essential oils inhibited Gram+ and Gram- bacteria. *Leiva* essential oil showed antimicrobial effects against a larger number of bacterial species (6) than *P. undulatum* (4) and *C. japonica* (4) essential oils. However, when observed, the inhibition of the essential oils from *P. undulatum* and *C. japonica* was higher than that of the *leiva*.

Understanding the effect of volatiles from traditional pineapple substrates on the rhizospheric microbial community can provide insight into their plant-substrate-microbial interactions; and may become a useful tool to promote pineapple growth and quality.

Acknowledgments: Partially supported by INOVA-4/2010, PROCONVERGÊNCIA – EU/FEDER e Secretaria Regional da Ciência, Tecnologia e Equipamentos

1. INE (2011) Estatísticas Agrícolas 2010. Instituto Nacional de Estatística, Lisboa, Portugal.
2. FAOSTAT (2009) Statistics Division.
3. JP Tavares, MT Silva (1995) Acta Horticult 425: 97-108.
4. JF Ponte Tavares (2004) In: Cultura de ananás em estufa: Ilha S. Miguel - Açores (Portugal). JF Ponte Tavares, JA Bettencourt Baptista (Eds). Profrutos, S. Miguel, Açores.
5. MD Mendes et al. (2011) *Industrial Crops and Products* 33: 710-719.
6. L Faleiro et al. (2005) *J. Agric. Food Chem.* 53: 8162-8168

**P 149. Antibacterial activity of *Eucalyptus* essential oils against pathogenic bacterial strains**

de Vasconcelos MCBM<sup>1</sup>, Dias C<sup>1</sup>; Rosa EA<sup>1</sup>; Saavedra MJ<sup>2</sup>

<sup>1</sup>CITAB/UTAD;

<sup>2</sup>CECAV/UTAD

mcarmov@utad.pt

Keywords: *Eucalyptus*; essential oils; antibacterial activity; infections.

Essential oils (EOs) of several plant species have been used in the pharmaceutical, cosmetics, food, industries and medicinal purposes [1,2]. Also, the antimicrobial properties of EOs have been recognized for many years [3,4]. In the present study *in vitro* antimicrobial activity of 15 plant species EOs of *Eucalyptus* have been evaluated, using disk diffusion bioassay, in triplicate [5,6] and minimum inhibitory concentration (MIC) [7]. Four reference bacteria were tested: *Pseudomonas aeruginosa* (ATCC10145), *Escherichia coli* (CECT434), *Staphylococcus aureus* (CECT976) and *Listeria monocytogenes* (ATCC15313). Gram-positive bacteria revealed higher sensitivity, when compared with the Gram-negative. Among the Gram-positive, the EOs from 5 *Eucalyptus* species (*E. bosistoana*, *E. botryoides*, *E. camaldulensis*, *E. cinerea* and *E. citriodora*) showed a total inhibition. This study reveals higher broad-spectrum antibacterial activity against Gram-positive in EOs than antibiotics (gentamicin and ciprofloxacin). *L. monocytogenes* was the most sensitive bacteria and *P. aeruginosa* was resistant to all EOs. These three bacteria were tested against dilutions (50, 25, 15, 5 and 1 % v/v) of the pure EOs to determine the MIC. The dilutions of 50 and 25% presented the same inhibition areas as the pure EOs, while the 15% dilution revealed inhibition for the three bacteria tested with *S. aureus* showing higher values (10.3-30mm). *E. coli* and *L. monocytogenes* revealed no sensitivity to the 5% dilution. The dilution of 1% did not exert sensitivity in any of the tested bacteria. Our results showed that these EOs could be useful as alternative tool to control pathogenic bacteria with natural compounds.

Acknowledgements: To the Centro de Biotecnologia Vegetal, Instituto de Biotecnologia e Bioengenharia (Uni. Lisboa, Faculdade de Ciências) and the FCT financial support (Carla Dias-SFRH/BGCT/33354/2008 and Maria do Carmo Vasconcelos-SFRH/BPD/70310/2010).

1. E Ben-Arye et al. (2011) Evidence-Based Comp. Alt. Med. doi:10.1155/2011/690346.
2. M H Salari et al. (2006) Clinical Microbiol. and Infection 12:178–196.
3. K H Chung et al. (2007) J. Microbiol. Biotech. 17:1848–1855.
4. S Prabuseenivasan et al. (2006) Comp. Alt. Med. 39: -8.
5. JMS Faria et al. (2011) Acta Hort. 925:61-66.
6. MJ Saavedra et al. (2010) Med. Chem. 6: 174-183.
7. AL Barry, C Thornsberry (1991) In: A Balows et al. (Eds.) Manual of clinical microbiology, 5th ed. American Society for Microbiology, Washington DC pp. 1117-1125.

**P 150. Antimicrobial activity of essential oils from squash pumpkin (*Cucurbita pepo*) seeds**

de Vasconcelos MCBM<sup>1</sup>, Aires A<sup>1</sup>; Dias C<sup>1</sup>; Almeida JA<sup>2</sup>; Santos P<sup>3</sup>; Rosa EA<sup>1</sup>; Saavedra MJ<sup>4</sup>

<sup>1</sup>CITAB/UTAD;

<sup>2</sup>UTAD;

<sup>3</sup>Douromel;

<sup>4</sup>CECAV/UTAD

mcarmov@utad.pt

Keywords: *Cucurbita pepo* seeds; Essential oils; Industrial by-products; Antimicrobial activity

The nutritive value of proteins, fatty acids and vitamins from pumpkin seed oils has been previously well described [1,2]. In this study we aim to evaluate the antibacterial activity of residues from the industrial processing (squash pumpkin seeds essential oils) in order to be used as sources of beneficial compounds evaluating the presence of antibacterial agents. The seed samples were provided from a large Portuguese company (Douromel - Fábrica de Confeitaria, Lda.) that processes fruits. In the present study *in vitro* antimicrobial activity of EOs from *Cucurbita pepo* seeds against two pathogenic bacteria (*Staphylococcus aureus* - CECT976 and *Listeria monocytogenes* - ATCC15313), using disk diffusion bioassay, in triplicate [3,4] and minimum inhibitory concentration (MIC) was determined [5]. The seeds were freeze-dried and the essential oils extraction was performed by hydro-distillation [6]. The pure EO and a diluted sample (120 mg/ml in dimethyl sulfoxide) were tested. The results showed that the pumpkin seeds EOs had an inhibitory effect on the growth of *S. aureus*, while *L. monocytogenes* revealed to be resistant to the EO tested. There were no significant differences between the pure (9 mm) and the diluted (10 mm) EO inhibitory effect on the growth of this bacteria. The *S. aureus* was also tested against dilutions (50, 25, 15, 5 and 1 % v/v) of the pure extracts to determine the MIC. However the dilutions did not exert sensitivity in the tested bacteria. These results reveal that the pumpkin seeds essential oils are potentially a good source of antibacterial agents against pathogenic bacteria.

Acknowledgements: To the company Douromel (Confeitaria Lda.) for the kind supply of the pumpkin residues samples. The authors also acknowledge the financial support provided by QREN (Quadro de Referência Estratégico Nacional) to the project Nutridouro "Healthy crystalised fruits and jams with a traditional flavour" (I&DT in Co-Promoção N° 13111).

1. M Murkovic et al. (1996) Z Lebensm Unters Forsch J. 202:275-278.

2. DG Stevenson et al. (2007) J. Agric. Food Chem. 55:4005-4013.

3. JMS Faria et al. (2011) Acta Hort. 925:61-66.

4. MJ Saavedra et al. (2010) Med. Chem. 6:174-183.

5. AL Barry, C Thornsberry (1991) In: A Balows et al. (Eds.) Manual of clinical microbiology, 5th ed. American Society for Microbiology, Washington DC pp. 1117-1125.

6. (COE) Council of Europe (2007) European Directorate for the Quality of Medicines. European Pharmacopoeia 6th Edition, Strasbourg.

**P 151. Chemical characterization and biological evaluation of essential oils from *Plectranthus* species grown in Portugal**

Simões MF<sup>1</sup>, Matias D<sup>1</sup>; Rijo P<sup>2</sup>; Rodríguez MV<sup>3</sup>; Zacchino SA<sup>4</sup>; Moiteiro C<sup>5</sup>

<sup>1</sup>Faculdade de Farmácia, Universidade de Lisboa, iMed.UL, Av. Prof. Gama Pinto, 1649-003 Lisboa, Portugal; <sup>2</sup>FF-UL, iMed.UL, Lisboa; C BIOS, Univ. Lusófona, Campo Grande 376, 1749-024 Lisboa, Portugal, <sup>3</sup>Fac. de Ciencias Bioquímicas y Farmacéuticas, Univ. Nacional de Rosario, Suipacha 531, (200), Rosario, Argentina; <sup>4</sup>Fac. de Ciências, Universidade de Lisboa, Centro de Química e Bioquímica, Campo Grande, 1149-016, Lisboa, Portugal

fasimoes@ff.ul.pt

Keywords: *Plectranthus grandidentatus*, *Plectranthus porcatius*, *Plectranthus venterii*, essential oils

*Plectranthus* is a Lamiaceae genus with interesting potential in drug discovery due to the bioactivities of its constituents and to the ethnopharmacological uses, namely, in Southern Africa [1, 2]. The essential oils (EOs) constituents of *Plectranthus* spp. are poorly studied. Following our research on *Plectranthus*, now we present a study of the *P. grandidentatus*, *P. porcatius* and *P. venterii* [3] EOs (grown in Lisbon from cuttings provided by the Kirstenbosch Nat. Bot. Gardens, South Africa).

Unlike *P. venterii* oil, the chemical composition of *P. grandidentatus* and *P. porcatius* EOs is varied and complex. Camphor is the most abundant monoterpene (52.8 %) of *P. venterii* together with elemol (5.1%), and sandaracopimarinal (6.7%).  $\beta$ -Caryophyllene (16.3%), caryophyllene oxide (5.2%),  $\alpha$ -humulene (6.5%) and 6,7-dehydroroleanone (6.3%) were some of the compounds of *P. grandidentatus*, while  $\gamma$ -terpinene (8.2%), 4-terpineol (7.6%),  $\delta$ -cadinene (6.6%), caryophyllene oxide (5.7%), cineol (5.9%) and  $\alpha$ -thujene (5.4%) were detected in *P. porcatius* EO. *P. grandidentatus* EO diverges from the one described for the plant collected in South Africa [4]. Diterpenes have rarely been identified on *Plectranthus* oils, being 6,7-dehydroroleanone the only isolated from the oil of *P. madagascariensis* [5]. This diterpene was now identified in *P. grandidentatus* EO, while sandaracopimarinal was identified in *P. venterii* EO for the first time.

The EOs showed low antimicrobial activity against *M. smegmatis*, *S. aureus*, *E. coli*, *P. aeruginosa* and *E. faecalis* (MIC 62.50-125  $\mu\text{gml}^{-1}$ ). *C. albicans* growth inhibition by *P. venterii* EO was higher (44.4%) than *P. grandidentatus* EO (27.1%) at 250  $\mu\text{gml}^{-1}$ . However, the activity of the tested components was lower than EOs, being sandaracopimarinal the only showing some activity (20.18 % at 200  $\mu\text{M}$ ).

Acknowledgments: Grateful to E van Jaarsveld, T Vasconcelos and A Duarte. Supported by FCT-MCES (iMed.UL, SFRH/BD/19250/2004), ANPCyT, PICT 0608, PEst-OE/QUI/UI0612/2011.

1. LJ Rice, et al. (2011) J S Afr Bot. 77: 947-959.
2. CW Lukhoba, MSJ Simmonds, AJ Paton (2006) J Ethnopharmacol. 103: 1-24.
3. E van Jaarsveld (2006) The Southern African *Plectranthus* and the art of turning shade to glade. Fernwood Press, South Africa.
4. K Maistry (2003) The antimicrobial properties and chemical composition of leaf essential oils of indigenous *Plectranthus* Lamiaceae species. U. KwaZulu-Natal, South Africa.
5. L Ascensão, et al. (1998) Int J Plant Sci. 159: 31-38.



**P 152. Composition and antibacterial activity of the essential oil of Peruvian *Dalea strobilacea* Barneby**

Benites J<sup>1,2</sup>, Moiteiro C<sup>3,2</sup>, Bravo F<sup>1,2</sup>, Ríos D<sup>1,2</sup>, López J<sup>1,2</sup>, Rojas M<sup>1</sup>, Torres I<sup>4,2</sup>, Sánchez I<sup>4</sup>

<sup>1</sup>Departamento de Ciencias Químicas y Farmacéuticas, Universidad Arturo Prat. Iquique, Chile;

<sup>2</sup>Instituto de Etnofarmacología (IDE), Universidad Arturo Prat. Iquique. Chile. Avenida Arturo Prat 2120. Iquique, Chile; <sup>3</sup>Faculdade de Ciências, Universidade de Lisboa, Departamento de Química e Bioquímica e Centro de Química e Bioquímica, 1749-016 Lisboa, Portugal; <sup>4</sup>Facultad de Farmacia y Bioquímica, Universidad Privada Antonio Guillermo Urrelo, Cajamarca, Perú

*cmmoiteiro@fc.ul.pt*

Keywords: *Dalea strobilacea* Barneby; Fabaceae; essential oil composition; GC and GC-MS; antibacterial activity

The genus *Dalea*, is highly diversified in the northern Peruvian Andes with some of these species being used as medicines [1]. Our research program is focused on the evaluation of the popular use of medicinal plants of the Chilean and Peruvian altiplano [2-5]. In this work we selected *Dalea strobilacea* Barneby (hierba de chil [6]), because residents use it to reduce gastrointestinal smooth muscle spasm and digestive disorders. Its infusion is highly prized as breakfast tea for its mild flavour that replaces the lemon verbena.

Regarding the chemical composition of essential oils (EOs) from genus *Dalea*, data are scarce in the literature. Moreover, no reports have been published about the chemical composition of the EO from *D. strobilacea*, collected in the region of Cajamarca, Perú. Therefore, we decided to carry out a study to determine its composition and also to explore its antibacterial capacity.

The composition of the EO from *D. strobilacea*, obtained by hydrodistillation of the aerial parts, was analyzed by GC and GC-MS, showing that  $\beta$ -phellandrene is the most abundant monoterpene (43.5%), together with  $\alpha$ -pinene (17.7%).

The *D. strobilacea* oil was tested against two Gram-negative and two Gram-positive bacteria. We observed a selective effect on Gram-positive bacteria, with MIC values of 8.7 and 10.7  $\mu\text{g/mL}$  as compared to the MIC value required to inhibit Gram-negative bacteria (59.5  $\mu\text{g/mL}$ ). A similar profile was observed in the effects caused by the antibiotic Vancomycin, namely low MIC values (2 and 5  $\mu\text{g/mL}$ ) in Gram-positive as compared to MIC value in Gram-negative bacteria (15.4  $\mu\text{g/mL}$ ). This antibacterial activity may be related to the chemical proportion of the main compounds,  $\beta$ -phellandrene,  $\alpha$ -pinene, as well as the minor components also present ( $\beta$ -pinene, myrcene and limonene).

Acknowledgments: Thanks to APEAC and A. C. Figueiredo, J. G. Barroso, L. G. Pedro. This study was supported by DI-UNAP, Conicyt-MEC 80100002 and PEst-OE/QUI/UI0612/2011.

1. S Baldeón et al. (2006) Rev. Peru. Biol. 13(2), 302–337.

2. L Rojo et al. (2009) BLAPCMA 8, 498–508.

3. J Benites et al. (2009) J. Chil. Chem. Soc. 54, 379–384.

4. J Benites et al. (2011) Chem. Nat. Compd. 46, 988–989.

5. J Benites et al. (2012) NPC 7(5), 611–614.

6. I Sánchez (2011) Especies Medicinales de Cajamarca I: Contribución Etnobotánica, Morfológica y Taxonómica. Perú, UPAGU y LCF Editorial. 131–133.

**P 153. Chemical composition and potential antioxidant activity of essential oil and different solvent extracts of discarded tobacco leaves**

Xianzhong Zhang<sup>1</sup>, Jun Yang<sup>2</sup>; Lifen Zhang<sup>1</sup>; Xingqian Ye<sup>1</sup>

<sup>1</sup>*School of Biosystems Engineering and Food Science, Zhejiang University;*

<sup>2</sup>*China Tobacco Zhejiang Industrial CO, LTD*

auhxyz@126.com

Keywords: discarded tobacco leaves, essential oil, solvent extracts, antioxidant activity

Every year in China, almost 20% of tobacco resources are discarded as processing waste. The discarded tobacco leaves not only pollute the environment, but also cause a big waste. In fact, there are abundant bioactive compounds in discarded tobacco leaves [1]. This study is designed to investigate the chemical composition and potential antioxidant activity of the essential oil and different solvent extracts of discarded tobacco leaves.

Essential oil was obtained by hydrodistillation (HD) and its composition was analyzed by GC and GC–MS. Antioxidant activity of essential oil and different solvent extracts (Petroleum ether, diethyl ether, dichloromethane, ethyl acetate and ethanol) were determined by two different systems, the DPPH free radical scavenging and the FRAP (Ferric Reducing Antioxidant Power).

GC and GC–MS analyses were resulted in the detection of 55 compounds in essential oil, representing 81.2% of the oil. Major components of the oil were neophytadiene (33.03%),  $\beta$ -damascenone (4.72%), methyl linoleate (1.86%), methyl linolenate (1.83%), megastigmatrienone (1.58%),  $\beta$ -damascone (1.54%) and solanone (0.22%). Results of the DPPH assay gives an  $IC_{50}$  range value of (2.07–9.88) mg/mL for all the samples studied. BHA (butylated hydroxyanisole) and ascorbic acid were used for comparison purposes. In all systems, ethyl acetate extract exhibited excellent activity potential than those of other extracts (Petroleum ether, diethyl ether, dichloromethane, and ethanol) and the oil. As expected, amount of total phenolics was very high in this extract ( $163.47 \pm 2.119$ g GAE mg/g extract). A positive correlation was observed between the antioxidant activity potential and total phenolic levels of the extracts

These results indicated that the essential oil and solvent extracts of discarded tobacco leaves had potential antioxidant activity, the discarded tobacco leaves could be considered as a potential resource of natural antioxidant.

1. Haiyan Wang et al. (2008) Food Chemistry. 107: 1399-1406.

**P 154. Comparative chemical study of *Uvariadendron angustifolium* (Engl. & Diels) R. E. Fries essential oils, a medicinal plant in benin**

Noudogbessi JP<sup>1</sup>, Adomou A<sup>2</sup>, Alitonou GA<sup>1</sup>, Adjalian E<sup>1</sup>, Avlessi F<sup>1</sup>, Sohounhloue KCD<sup>1</sup>, Menut C<sup>3</sup>

<sup>1</sup>Université d'Abomey-Calavi ; Ecole Polytechnique d'Abomey-Calavi, Laboratoire d'Etude et de Recherche en Chimie Appliquée (LERCA) ; 01 BP 2009 Cotonou, BÉNIN;

<sup>2</sup>Herbier National, Département de Botanique, Faculté des Sciences et Techniques, Université d'Abomey-Calavi, République du Bénin;

<sup>3</sup>Institut des Biomolécules Max Mousseron, Faculté de Pharmacie, 15 Avenue Charles Flahault, Bâtiment E, BP 14491, 34093 Montpellier France

chantal.menut@univ-montp2.fr

Keywords: *Uvariadendron angustifolium*; essential oils; citral; methyl eugenol

*Uvariadendron angustifolium*, syn. *Uvaria angustifolia* (Annonaceae) is a West African forests tree which can reach 15-40 ft high [1]. This species, which is found in the south of Benin, is used in traditional medicine to treat rheumatics and gastric ailments, paludism, or for flavouring local dishes (leaves) [2]. The only chemical investigations on *Uvariadendron* genus concern two species: *U. connivens*, which is characterized by phenylpropene compounds in its seeds (elemicin, cinnamaldehyde and 3,4,5-trimethoxy cinnamyl alcohol) [3] and *U. calophyllum*, which furnished, by hydrodistillation of its wood, stem bark and roots, essential oils rich in sesquiterpenes (mainly (E)- $\beta$ -caryophyllene and  $\alpha$ -santalene) [4].

The present work reports a comparative chemical study of volatile components obtained by hydrodistillation of leaves, stems, bark and roots of *Uvariadendron angustifolium* collected in Ketou (Benin). The essential oils, obtained with yields comprised between 0.12 and 0.66%, were analyzed by GC/FID and GC/MS on two capillary columns of different polarities (HP-5 and Carbowax 20M). A great variability of the chemical composition was observed depending on the plant material part treated. The essential oils from leaves and stems were constituted by high levels of neral (29.7-34.2%) and geranial (42.7-46.5%) while methyl eugenol was the major component of the volatile extracts from bark (68.3%) and roots (85.3%). These chemical characteristics justify the use of leaves as food flavouring.

Acknowledgments: The authors wish to thank the University of Abomey-Calavi Scientific Council members for their agreement for this work and the Government of Benin Republic for its financial support.

[1] J Hutchinson et al. (1954) Flora of West Tropical Africa. 2<sup>nd</sup> edition, Vol 1, Part 1, Crown Agents, London S.W.1.

[2] Flore Analytique du Bénin (2006), A Akoègninou, WJ Van Der Burg, LLG Van Der Maessen (Eds.), Backhuys Publishers, p 1034.

[3] I Mohammad, PG Waterman (1985) J. Nat. Prod. 48: 328-329.

[4] FF Boyom et al. (2005) J. Essent. Oil Res. 17: 128-129.

**P 155. Mycorrhiza biotechnology in the cultivation of essential oil yielding traditional african medicinal plants.**

Koehorst RR, Laubscher CP

*Cape Peninsula University of Technology*

*koehorstr@cput.ac.za*

Keywords: Sustainable, symbiosis, soil biota, mineral supplementation

The cultivation of quality essential oil yielding Traditional African Medicinal (TAM) plants in Africa is threatened by low levels of minerals in local soils, combined with environmental stressors such as water and salt stress. Mycorrhiza is the symbiotic association between plant roots and specific species of soil borne fungi, and it can play a large part in the cycling of mineral nutrients from the soil into the plant, as well as protecting the host plant against stress, both environmental and cultural. The relationship of the fungi and host can take a variety of forms and can have a range of effects upon the growth and development of the host plant. The most commonly utilized mycorrhizal relationship type is arbuscular mycorrhiza (AM). AM is used in the cultivation of many important horticultural crops, and has been shown to have an effect upon the growth, development and yield of both ornamental and medicinal plant species. There is a large amount of information available on the use of AM in an ornamental horticultural situation, and so this paper will outline the current perspectives on the use of AM in the cultivation of essential oil yielding Traditional Medicinal plants, with a specific focus on the use of AM in an African setting (based on soil and environmental qualities that are relevant to Africa). This paper will summarise the history of traditional medicine and its cultivation/ harvesting in the unique African setting, specifically pertaining to nutrient applications and the potential of mycorrhiza to decrease reliance on alternate forms of nutrient supplementation. To further illustrate the potential of AM in the cultivation of TAM in Africa, the paper will also discuss a selection of TAM plants and review the potential for the use of AM in their cultivation in Africa.

**P 156. Gamma irradiation in food preservation and health promotion – influence in the aroma compounds of spices**

Duarte RC<sup>1</sup>, Lopes JF<sup>2</sup>, Santana CJ<sup>2</sup>, Nunes IS<sup>2</sup>, Villavicencio ALCH<sup>3</sup>, Gaspar EMM<sup>2</sup>

<sup>1</sup>IPEN-CNEN/SP-Brazil; CQFB-Requimte, FCT- UNL-Portugal;

<sup>2</sup>CQFB-Requimte, FCT- UNL. Portugal; <sup>3</sup>IPEN-CNEN/SP-Brazil

renatocduarte@yahoo.com.br

Keywords: Spices; Clove; Volatile compounds; Gamma Irradiation; HSPME-GC-MS

History illustrates how spices have played a prevalent part in man's life and death. Due to their strong scent, mankind have used spices to fulfil the absence of hygiene habits, for embalming and for religious purposes. However, the most important role of spices was their ability to heal and perpetuate life, through their use as aphrodisiac, in dieting and as medicines. In fact, at 1500s, when the "Spice Wars" occurred between the Portuguese and the Dutch and later the Dutch and the English, the reason was their medicinal importance, because spices were considered as a miracle cure for the plagues. A list of cures may justify why many of them are exactly the same spices which commonly are found in spice racks for our everyday cooking. For over 2000 years, clove, *Syzygium aromaticum*, was consumed in different parts of Asia and later transported to Europe. Cloves are the dried aromatic flower buds of the evergreen trees, highly flavored and renowned for their health benefits (antiseptic, bactericide and fungicide) and nutritional content. The spice industry is faced with the problem of reducing microbial contamination derived from the techniques used in the post-harvest steps. Gamma irradiation is becoming a food preserving methodology, reducing also the incidence of foodborne disease. Depending on the energy, the irradiation process is capable of extending the shelf life of food, reducing spoilage, inhibiting sprouting and destroying insects. However, the irradiation treatment can induce chemical changes in food [1, 2]. The aim of this work is to investigate the effect of gamma irradiation in the chemical composition of spices. Beginning with the aroma compounds, using the free solvent HS-SPME-GC-MS, the effect of <sup>60</sup>Co irradiation, 5.0, 10.0, 20.0 and 30.0 kGy, in the volatile chemical profile of clove are being studied. Preliminary results show some quantitative differences, but also some qualitative differences in terpenoid composition.

Acknowledgments: The authors acknowledge financial support from Brazilian CNPq, CAPES (Renato Duarte - 201998/2011-9) and to IPEN-CNEN/SP-Brazil and Portuguese FCT (João Lopes - FCT/SFRH/BD/40564/2007; José Santana - SFRH/BNDE/51585/2011 and Inês Nunes - SFRH/BD/33809/2009). Authors also are grateful to Hikari (Brazil) for the samples supply.

1. M Suhaj, J Horvátová (2007). J. Food. Nutr. Res. 46, 112-122.
2. A L C H Villavicencio *et al.* (2000). Radiat. Phys. Chem. 57, 289-293.

**P 157. Bio-valorization of Azorean *Cryptomeria japonica* essential oils**

Moiteiro C<sup>1</sup>, Esteves T<sup>1</sup>; Rojas R<sup>2</sup>; Alvarez SL<sup>3</sup>; Zacchino SA<sup>3</sup>; Bragança H<sup>4</sup>

<sup>1</sup>Faculdade de Ciências da Universidade de Lisboa, Departamento de Química e Bioquímica, Centro de Química e Bioquímica, Campo Grande, 1149-016, Lisboa, Portugal;

<sup>2</sup>Laboratórios de Investigación y Desarrollo, Facultad de Ciencias y Filosofía, Universidad Peruana Cayetano Heredia, Lima, Perú;

<sup>3</sup>Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Suipacha 531, (200), Rosario, Argentina;

<sup>4</sup>Instituto Nacional de Recursos Biológicos I.P., Quinta do Marquês, 2780-159 Oeiras, Portugal

cmmoiteiro@fc.ul.pt

Keywords: *Cryptomeria japonica*; Essential oils; GC and GC-MS; Antimicrobial activity

*Cryptomeria japonica* D. Don wood production is of great economic importance in Azores Archipelago (Portugal), having a high value as a building material, furniture and also as an ornamental tree. Every year Azorean *C. japonica* wood industry generates a large amount of residues (bark, wood and aerial parts) with 30% of the wood wastes derived only from sawmills wood processing [1].

Essential oils (Eos) from aerial parts, bark and heartwood of *C. japonica* from Azores were analyzed by GC and GC-MS. The main compounds found in aerial parts were  $\alpha$ -pinene (9.6-29.5%), (+)-phyllocladene (3.5-26.6%), *ent*-kaur-16-ene (0.2-20.6%), sabinene (0.5-19.9%) and limonene (1.4-11.5%). Heartwood oils were characterized by a high content of cubebol (2.7-39.9%) and *epi*-cubebol (4.1-26.9%), absent in aerial parts. Elemol and eudesmol isomers were found in all aerial parts and heartwood oils, while (+)-phyllocladene was not present in heartwoods. Bark oils were composed of dehydroferruginol and ferruginol diterpenes and a wide variety of sesquiterpenes ( $\delta$ -cadinene,  $\alpha$ -muurolene, *epi*-zonaren, cubenol, T-muurolol,  $\beta$ -eudesmol,  $\gamma$ -eudesmol and hedyariol). Azorean *C. japonica* oils exhibited significant chemical differences comparatively to native plants from Asia [2].

From the several fungi that affect many plant species, *Botrytis cinerea*, was inhibited by all Eos (MIC 100  $\mu$ g/ml) tested with fungistatic activity, while *Fusarium* spp. And *Cryphonectria parasitica* showed resistance towards all oils tested. Furthermore, all the Eos exhibited moderate activity against the wood surface contaminant *Trichoderma harzianum* and the human pathogenic bacteria *Streptococcus aureus* and *S. faecium* (MIC 100-200  $\mu$ g/ml). The Eos of leaves (MIC 25-50  $\mu$ g/ml) were more active than those of the heartwood (MIC 50-200  $\mu$ g/ml) against multidrug-resistant *Mycobacterium tuberculosis*. These results illustrate the potential bio-valorization of the wastes produced by *C. japonica* wood industry, until now only seen as disposable.

Acknowledgments: Collaborative work was performed under the auspices of the CYTED, PIBATUB, INETI and project Pest-OE/QUI/UI0612/2011. The authors thank Engineer J. Belerique, of the DRRF for providing the *C. japonica* and the facilities for this study.

1. JFP Mendes (2008) A Estratégia para a Floresta na Região Autónoma dos Açores, DRRF, *Seminário Certificação Florestal na Região Autónoma dos Açores*, Ponta Delgada.
2. H-J Gu, et al. (2009) *J. Agric. Food Chem.* 57, 11127-11133.

**P 158. Floral scents of *Adansonia*: chemical characterization and role in the pollination process**

Razanamaro O<sup>1</sup>, Menut C<sup>2</sup>; Rasoamanana E<sup>1</sup>; Rakouth B<sup>1</sup>; Danthu P<sup>3</sup>

<sup>1</sup>Département de Biologie et Ecologie Végétales, Faculté des Sciences, Université d'Antananarivo, B.P. 906, Antananarivo (101), Madagascar;

<sup>2</sup>Institut des Biomolécules Max Mousseron, Faculté de Pharmacie, 15 Avenue Charles Flahault, Bâtiment E, BP 14491, 34093, Montpellier, France;

<sup>3</sup>CIRAD, URP Forêts et Biodiversité, BP 853, Antananarivo, Madagascar and Campus de Baillarguet, 33398 Montpellier Cedex 5, France

chantal.menut@univ-montp2.fr

Keywords: Baobab; *Adansonia*; scent; pollination; Madagascar

The genus *Adansonia* (baobab, Malvaceae family) includes eight species among them seven were identified in Madagascar and six are endemic and belong to the two sections Brevitubae and Longitubae [1]. They are emblematic in the island, nevertheless more knowledge is necessary regarding their productive biology; among the different parameters involved in the pollination process, the mechanism of scent release by flowers is of importance [2]. The aim of this study was : (i) the identification of the volatile compounds released by the flowers of the different botanical species and (ii) elucidation of their role in the flowers/pollinating agents interaction.

The study was performed on individual trees of the six species endemic in Madagascar during their flowering period.

Two complementary methods were used to catch these volatile compounds: solvent extraction (using 3M engineered fluid HFE-7200, ethoxy-nonafluorobutane) and solid phase microextraction (using "Mono Trap" disks). Their identification was performed by GC/FID and GC/MS. A great chemical variability was observed according the botanical section. The scents released by flowers of the species belonging to the Longitubae section were dominated by aromatic compounds (mainly 2-phenylacetonitrile) while aliphatic compounds (mainly heptadec-8-ene) characterized the Brevitubae group.

The chemical characteristics of the scents related to the Longitubae section were previously described as possessing entomophilous properties [3]. Other studies on the floral morphology, the nectar and the pollinating agents are in progress.

[1] DA BAUM (1995) The comparative pollination and floral biology of Baobabs (*Adansonia*-Bombacaceae). Ann. Missouri Bot. Gard. 82: 332-348.

[2] C Suchet (2010) Ecologie et Evolution des odeurs florales chez *Antirrhinum Majus*. Thèse de Doctorat, Institut National Polytechnique de Toulouse.

[3] M Takashi, R Yamaoka, T Yahara (1998) Floral scents of hawkmoth- pollinated flowers in Japan. J. Plant. Res. 111: 199- 205.

**P 159. Antifungal and anti-inflammatory potential of *Lavandula stoechas* and *Thymus herba-barona* essential oils**

Zuzarte M<sup>1</sup>, Gonçalves MJ<sup>1</sup>, Cavaleiro C<sup>1</sup>, Cruz MT<sup>2</sup>, Benzarti A<sup>3</sup>, Marongiu B<sup>4</sup>, Maxia A<sup>5</sup>, Piras A<sup>4</sup>, Salgueiro L<sup>1</sup>

<sup>1</sup>Centro de Estudos Farmacêuticos, Faculdade de Farmácia, Pólo das Ciências da Saúde, Universidade de Coimbra, Azinhaga de S. Comba 3000-354, Coimbra, Portugal.

<sup>2</sup>Centro de Neurociências e Biologia Celular, Universidade de Coimbra, 3000-295, Coimbra, Portugal.

<sup>3</sup>Faculty of Science, University of Monastir, Monastir, Tunisia

<sup>4</sup>Dipartimento di Scienze Chimiche e Geologiche, Università degli Studi di Cagliari, Cittadella Universitaria di Monserrato, SS 554, Km 4.500, 09042 Cagliari, Italy

<sup>5</sup>Dipartimento della Scienza della Vita e dell'Ambiente-Macrosezione di Botanica e Orto Botanico, Università degli Studi di Cagliari, Viale Sant' Ignazio, I-09123 Cagliari, Italy

ligia@ff.uc.pt

Keywords: *Lavandula stoechas*; *Tymus herba-barona*; essential oil; antifungal; anti-inflammatory; toxicity

This study reports the chemical composition and antifungal activity of *Lavandula stoechas* and *Thymus herba-barona* essential oils against fungi responsible for human infections and food contamination. Also the anti-inflammatory potential and toxicity of the oils on a macrophage cell line is reported.

*L. stoechas* essential oil was rich in fenchone (38.0%) and camphor (28.1%) whereas *T. herba barona* oil showed high amounts of two phenols, carvacrol (54.4%) and thymol (30.2%). The latter was the most active oil against the tested fungi but evidenced high toxicity on macrophages. *L. stoechas* was active against dermatophyte strains and showed potential anti-inflammatory activity, by inhibiting the production of NO in LPS-stimulated macrophages, at concentrations from 0.16 to 0.64 µl/ml. The oil did not affect cells viability up to 0.32 µl/ml.

These results support the use of *L. stoechas* in the development of phytopharmaceuticals for the management of dermatophytosis and/or inflammatory-related diseases. Regarding *T. herba-barona*, it can be used as a preservative in storage products, due to its ability to inhibit *Aspergillus* growth.

Acknowledgments: This work was funded through national funds from FCT – Fundação para a Ciência e a Tecnologia under the project CEF/POCI2010/FEDER and a PhD fellowship to Mónica R. Zuzarte (SFRH/BD/40218/2007).



**P 160. Essential oil of *Ocimum americanum* L. and its application in perfumery**

Benevides PC, Figueiredo RO; Neto AP; Costa IM; Barbizan DS; Porto C; Gallucci SM

*Natura Inovação e Tecnologia de Produtos Ltda.*

*roseanefigueiredo@natura.net*

Keywords: *Ocimum americanum*; estoraque; essential oil; perfumery

Studies on biodiversity prospection are extremely important for the cosmetic industry and Brazil has a great potential in this field, due to its huge diversity of native and adapted species [1]. Based on previous works of aromatic species prospection in the Amazon region, the aim of this study was to investigate the potential use of essential oil obtained from aerial parts of the endemic methyl cinnamate chemotype of *Ocimum americanum* (popularly known in north of Brazil as Estoraque) for the fragrance industry. Production of Estoraque was done following an organic agricultural model (IBD organic certification) in the community of Campo Limpo - Santo Antonio Tauá, state of Pará. The steam distillation of aerial parts of Estoraque yielded 0.15-0.25% of essential oil that was analyzed by gas chromatography-mass spectrometry (GCMS). The major compounds were: *E*-methyl cinnamate (42.0%), carvone (9.3%) and *trans*-caryophyllene (9.1%). This study reports the industrial scale production, phenological aspects, toxicology and olfactory significance of methyl cinnamate chemotype of *O. americanum*, revealing an innovative ingredient for perfumery use.

Acknowledgements: The authors would like to thank Veronica Kato (Natura Perfumer) and Givaudan.

1. AJ Beattie, W Barthlott, E Elisabetsky, R Farrel, CT Kheng, I Prance, J Rosenthal, Simpson, RRB Leakey, M Wolfson, K Ten Kate, S Laird. 2005. New products and industries from biodiversity (Chapter 10), *In: Ecosystems and Human Well-Being: Vol 1. Current State and Trends*, (Eds R Hassan, R Scholes and N Ash), Findings of the Condition and Trends Working Group of the Millennium Ecosystem Assessment, Island Press, Washington DC, USA, pp. 273-295.

**P 161. Microalgae: Promising source in the production of carotenoids and essential oils**

Kiril B<sup>1</sup>, Espirito Santo C<sup>1</sup>, Costa JR<sup>1</sup>; Mota P<sup>1</sup>

<sup>1</sup>*Microalgae - Investigação e Desenvolvimento, Lda., Bencanta, 3040-316 Coimbra, Portugal*

*kiril@esac.pt*

Keywords: *Haematococcus pluvialis*, *Chlamydomonas* sp., fatty acids, carotenoids.

Microalgae are a potential source of biologically active compounds, namely carotenoids and fatty acid, showing higher efficiency than those obtained from traditional crops. Regarding this they can be considered as functional “ingredients”, justifying the recent increased interest in their commercial, pharmaceutical, environmental and/or agri-food industries applicability. This work aimed to evaluate and determine the total lipid content and fatty acid profile, as well as to quantify the amount of carotenoids, in two species of microalgae *Haematococcus pluvialis* Flotow (Hp), *Chlamydomonas* sp. Ehrenberg (Csp), both belonging to the phylum *Chlorophyta* [1] Their growth was monitored and assessed in two different culture media. Solvent extraction technique was used for lipid content and gas chromatography, spectrophotometry UV-vis and mass spectrometry (MS) to evaluate the profile of methyl esters of fatty acids and carotenoids, respectively [2, 3]. The percentages of total lipid for Hp and Csp were 12,05%, 8,33% and 23,82% and 14,65% in the two media, respectively. The profile of fatty acids proved to be variable, both in size of the hydrocarbon chains and for the position of the unsaturated carbon atom, conditioned by the culture media. The evaluation of the carotenoids content for both species, showed similar variability. Hp (H2) presented higher content of this bioproduct, 28.4% and 39% in both media. Our results were similar to those obtained in other studies, with both species, which have shown the increasing importance of these microorganisms in the production of essential compounds, with specific physiological functions [4].

Acknowledgements: Abílio, Sobral, PhD – Departamento de Química (FCTUC). Jorge Varejão, Professor Adjunto - ESAC.

1. Spolaore *et al.* (2006) – Commercial Applications of Microalgae. *Jornal of Bioscience and Bioengineering*. 101, 2: 87-96;
2. Hossain *et al* (2008) - Biodiesel Fuel Production from Algae as Renewable Energy. *American Journal of Biochemistry and Biotechnology* 4 (3):250-254;
3. Holtin *et al* (2009) - Determination of astaxanthin and astaxanthin esters in the microalgae *Haematococcus pluvialis* by LC-(APCI)MS and characterization of predominant carotenoid isomers by NMR spectroscopy. *Anal Bioanal Chem* 395:1613–1622;
4. Guedes *et al.*(2011) - Microalgae as Sources of Carotenoids. *Marine Drugs*.9: 625-644.

**P 162. Understanding the ecological role of volatiles in the host-plant resistance: the case of the portuguese sweet potato (*Ipomoea batatas*) from Aljezur**

Nunes IS<sup>1</sup>, Lopes JF<sup>1</sup>; Santana CJ<sup>1</sup>; Figueiredo AC<sup>2</sup>; Gaspar EMM<sup>1</sup>

<sup>1</sup>CQFB-Requimte, FCT- UNL. Portugal;

<sup>2</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal

inez.sn@gmail.com

Keywords: *Ipomoea batatas*; Lira; Protected Geographical Indication; volatiles; GC; GC-MS

A major constraint to sweet potato production worldwide is the sweet potato weevil, *Cylas* spp.. Although there are no records of this species in Europe, interceptions of this pest should be considered due the worldwide trade of sweet potato. The development of insect-resistant sweet potato lines is seen as a viable component in the integrated management of this and other pests. Qualitative and quantitative analysis of the host-plant volatiles has been one of the approaches for successful identification of resistance in germplasm resources [1].

Sweet potato, *Ipomoea batatas* (L.), is an important food crop and it is an alternative source of carbohydrates, taking the fourth place after rice, corn and cassava. The tuberous roots and the green tops constitute a important source of vitamins and minerals [2,3].

In Portugal, *Lira* variety is cultivated in Aljezur – a Protected Geographical Indication (PGI) area [4]. Phytosanitary problems have been rarely detected. Only sporadic attacks of red spider mite (*Panonychus ulmi*) and some fungi have occurred, which normally do not involve treatments, since they haven't represented any economic loss for farmers [5].

In order to understand the potential resistance of this Portuguese variety, the volatile composition of all plant is being studied. The volatiles were isolated by hydrodistillation and analyzed by GC and GC-MS as in [6]. Different chemical profiles were obtained for the aerial parts and storage roots.  $\beta$ -Caryophyllene and germacrene are the main compounds of the volatiles from the aerial parts, while hexadecanoic acid is the major component of storage roots.  $\alpha$ -Humulene, a sesquiterpene known to be repellent to insects [7] is also present in the aerial parts and storage roots. Additional studies are needed in order to elucidate the role of sweet potato volatile composition, mainly the sesquiterpene fraction, in the resistance of plant against pests.

Acknowledgements: ISN, JFL and JCS acknowledge FCT (SFRH/BD/33809/2009; SFRH/BD/40564/2007; SFRH/BDE/51535/2011) and Associação de Produtores de Batata Doce de Aljezur.

1. RR Korada et al. (2010) *Curr. Sci.* 99:1597-1601.

2. N Zuraida (2003) *J. Litbang Pertanian* 22:150-155.

3. S Islam (2006) *J. Food Sci.* 71:13-21.

4. Council Regulation (EC) No 510/2006, C 324/31, 19.12.2008.

5. Caderno de especificações Batata doce de Aljezur - Indicação Geográfica Protegida (2008) 1-54.

6. P Barbosa et al. (2010) *J Nematol* 42: 8-16.

7. Y Wang, SJ Kays (2002). *J. Amer. Soc. Hort. Sci.* 127: 656-662

**P 163. Phytochemical and genetic diversity in *Mentha cervina* (L.) Opiz based on essential oils profile and ISSRs markers fingerprinting**

Rodrigues LS<sup>1</sup>, Póvoa O<sup>2</sup>; Van Den Berger C<sup>3</sup>; Figueiredo AC<sup>4</sup>; Monteiro A<sup>1</sup>

<sup>1</sup>Centro de Botânica Aplicada à Agricultura (CBAA), Instituto Superior de Agronomia, Technical University of Lisbon, Tapada da Ajuda 1349 017 Lisboa, Portugal;

<sup>2</sup>Escola Superior Agrária de Elvas, Rua de Alcamim nº 19, 7350-903 Elvas, Portugal.;

<sup>3</sup>Departamento de Ciências Biológicas, Universidade Estadual de Feira de Santana, Av. Transnordestina s.n., 44036-900, Feira de Santana, Bahia, Brazil;

<sup>4</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, IBB, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749 016 Lisboa, Portugal.

liarodrigues@isa.utl.pt

Keywords: *Mentha cervina*; Essential oils; Genetic diversity; ISSRs; Conservation genetics

*Mentha cervina* L. is an aromatic plant traditionally used in Portugal to flavour food dishes and for its medicinal properties. In the last decade, the growth of commercial demands, the excessive harvesting from the wild and the unfavourable conservation status of their habitat, has shrunk the natural resource of *M. cervina* to a narrow distribution [1], leading to the classification of this species as Near Threatened in the IUCN Red List of Threatened Species [2].

In this study, phytochemical and genetic variation was studied to evaluate the level and distribution of diversity in *M. cervina* populations in order to provide guidelines for the conservation and sustainable use of this medicinal species. Gas Chromatography and Gas Chromatography–Mass Spectrometry was used to reveal the essential oil profile and Inter-simple sequence repeats (ISSRs) markers fingerprinting to assess the population's genetic structure and diversity.

*M. cervina* essential oils showed high uniformity ( $S_{corr} \geq 0.95\%$ ), with pulegone has the major essential oils compound in all of the populations (68-83%) collected at full flowering, in different growing conditions (51-82%), and for all the developmental stages studied (47-82%). The *M. cervina* populations exhibited a relatively low genetic diversity ( $PPB = 14.3-64.6\%$ ,  $H_e = 0.051-0.222$ ,  $I = 0.076-0.332$ ), with high structuring between them ( $G_{ST} = 0.51$ ). However, the genetic diversity at species level was relatively high ( $PPB = 97.7\%$ ;  $H_e = 0.320$ ).

The low genetic diversity found within these populations seems to result from genetic drift and inbreeding that lead to diminished fitness in this species, which may be observed in the rather unusual uniformity of the essential oil composition. From a conservation perspective, the low genetic and phytochemical diversity observed is symptomatic and a signal that ecological management of *M. cervina* habitats is necessary to prevent the consequent decline in population size that could increase the risk of extinction due to demographic and environmental stochasticity.

Acknowledgments: Leandra Rodrigues is grateful to the FCT for the grant SFRH/BD/38143/2008.

[1] Rodrigues *et al.* (2008) Flavour Fragr. J. 23, 340-347.

[2] Rhazi L., Grillas P (2010). IUCN Red List of Threatened Species.

**P 164. Anti-candidal activity of 7-hydroxycalamenene isolated from *Croton cajucara***

Azevedo MMB<sup>1</sup>, Chaves FCM<sup>2</sup>; Bizzo HR<sup>3</sup>; Alviano CS<sup>4</sup>; Alviano DS<sup>4</sup>

<sup>1</sup>Instituto de Química, CT, UFRJ, Rio de Janeiro, RJ, Brazil;

<sup>2</sup>EMBRAPA Amazônia Ocidental, Manaus, AM, Brazil;

<sup>3</sup>EMBRAPA Agroindústria de Alimentos, Rio de Janeiro, RJ, Brazil;

<sup>4</sup>Instituto de Microbiologia Paulo de Góes, CCS, UFRJ, Rio de Janeiro, RJ, Brazil

marimbarros@gmail.com

Keywords: 7-hydroxycalamenene; *Croton cajucara*; red sacaca; essential oil; anti-Candidal activity

The leaves and bark from *Croton cajucara* Benth. (family Euphorbiaceae), a shrub from the Amazon, have been used locally used in folk medicine to treat diabetes, malaria, gastrointestinal and liver disorders [1]. A chemotype of this species was found, with an essential oil rich in 7-hydroxycalamenene [2]. 7-hydroxycalamenene is a hydroxylated sesquiterpene of molecular weight 218 found in *Heritiera ornithocephala* [3], *Eremophila drummondii* [4], *Heteroscyphus planus* [5], *Tilia europea*, *Morus alba* [6], *Ulmus thomasii* [7] and other elm species, and methanolic and dichloromethanic extracts of *Bazzania trilobata*. This substance is reported to have antifungal activity against *Botrytis cinerea*, *Cladosporium cucumerinum*, *Phytophthora infestans*, *Pyricularia oryzae* and *Septoria tritici* [8]. During our studies with *C. cajucara* essential oil, we isolated 7-hydroxycalamenene by silicagel column chromatography. The pure compound (+98% by GC) was tested against some *Candida* species. Minimum inhibitory concentration (MIC) was evaluated in triplicate according standard method from the Clinical and Laboratory Standards Institute (CLSI) [9]. The calculated MIC's were 39,06 µg/mL was found to *C. albicans* (ATCC10231), *C. dubliniensis* e *C. albicans* (CaA), of 78,125 µg/mL to *C. albicans* (Cab) e *C. parapsilosis* and *C. albicans* (CaB) > 2500 µg/mL. From these data, it was observed 7-hydroxycalamenene is a compound with good activity against these *Candida* species.

**P 165. Antimicrobial activity of essential oils from *Lippia alba* (Miller) N.E. Brown**

Costa DCM<sup>1</sup>, Azevedo MMB<sup>2</sup>; Amancio CA<sup>1</sup>; Blank AF<sup>3</sup>; Alviano CS<sup>1</sup>; Alviano DS<sup>1</sup>

<sup>1</sup>Instituto de Microbiologia Paulo de Góes, CCS, UFRJ, Rio de Janeiro, RJ, Brazil;

<sup>2</sup>Instituto de Química, CT, UFRJ, Rio de Janeiro, RJ, Brazil;

<sup>3</sup>UFS, SE, Brazil

marimbarros@gmail.com

Keywords: *Lippia alba*; essential oils; antimicrobial activity

*Lippia alba* (Miller) N.E. Brown (Verbenaceae) is an aromatic plant known in Brazil as “Erva-cidreira-do-campo”, and that is native to southern Texas in the United States, Mexico, Caribbean, Central America, and South America [1]. This specie occurs in practically all regions of Brazil and has a large importance in brazilian folk medicine, with its use widely distributed in the country. *L. alba* is locally used as sudorific, expectorant, on the treatment of colds and flu, antidysenteric, stimulant, antireumatic and against hypertension. Its use in popular medicine can be explained, partly, by the bioactive volatile constituents. However, this specie is characterized by variability in morphology and in the chemical composition of the essential oil. Secondary metabolics as limonene, carvone, citral, myrcene, are often found in essential oils of the plant [2]. The present study evaluated the antimicrobial potential of 6 different chemotypes of essential oils (EOs) of *L. alba*, it was analyzing the inhibition of the activity of these oils against pathogenic microorganisms and his virulence factors, like peptidase activity. During the studies with *L. alba* essential oils, were made tests against bacteria, yeasts and filamentous fungi, like *Escherichia coli*, *Candida albicans* and *Trichophyton rubrum*. The minimum inhibitory concentration was based on protocols CLSI [3] and the results showed MIC of 312 µg/mL against *C. albicans* (ATCC10231) and 70 µg/mL to *T. rubrum*. To determining a possible target of antifungal activity, it was performed assays of proteolytic activity inhibition, through to the method described by Buroker-Kilgore and Wang [4]. The results showed that *L. alba* EOs presented proteolytic activity inhibitions to *C. albicans* secreted proteases, mainly serine and aspartic classes. The results show a great antifungal activity of *L. alba* EOs, suggesting that more researches are required to considerate the EOs like alternatives for a new and effective antimicrobial therapy.

**P 166. Inhibition of extracellular peptidases of *Rhizopus oryzae* by *Melissa officinalis* essential oil.**

Almeida CA<sup>1</sup>, Blank AF<sup>2</sup>, Alves PB<sup>2</sup>, Campos-Takaki GM<sup>3</sup>, Alviano CS<sup>1</sup>, Alviano DS<sup>1</sup>

<sup>1</sup>Instituto de Microbiologia Paulo de Góes, CCS, UFRJ, Rio de Janeiro, RJ, Brazil;

<sup>2</sup>UFS, SE, Brazil;

<sup>3</sup>UNICAP, PE, Brazil

catiamancio@yahoo.com.br

Keywords: *Melissa officinalis*; essential oil; *Rhizopus oryzae*; extracellular peptidases.

Lemon balm (*Melissa officinalis* L., Lamiaceae) is a well-known herb used to give fragrance to different food and beverage products. It has also been used as a medicinal plant for treatment of headaches, gastrointestinal disorders, nervousness, and rheumatism [1]. The essential oil is a well-known antibacterial and antifungal agent. Mucormycosis is a life-threatening fungal infection of worldwide distribution and increasing importance affecting mainly immunocompromised hosts. It is caused by fungi of the Mucorales order involved in human or animal diseases. *Rhizopus* spp. usually causes infections of sinuses and rhinocerebral structures, cutaneous tissue or lungs. In the latter infections, the fungi generally disseminates in almost half of all cases. Pathological hallmarks of mucormycosis are vascular invasion with hyphae, thrombosis, infarction, and necrosis of tissue, which lytic enzymes produced by *Rhizopus* spp. are thought to be directly or indirectly involved [2]. In addition to the lack of potent, well-tolerated antifungals, the poor prognosis can clearly be attributed to the exceptionally aggressive behavior of the fungus in the human host. In this study, we evaluated the antifungal and anti-peptidase activities of *M. officinalis* essential oils (EO) against *R. oryzae*. Firstly, the susceptibility of *R. oryzae* to the EO was tested by microdilution, based on protocol M38-A2 [3]. Spores were diluted in RPMI 1640 containing L-glutamine and buffered to pH 7.0 with 0.165 M MOPS, and the MIC obtained for *R. oryzae* was 625 µg/mL. To determining a possible target of this antifungal activity, it was performed assays of proteolytic activity inhibition, through the method described by Buroker-Kilgore and Wang [4]. Cell-free supernatants of *R. oryzae*, grown in RPMI medium, were incubated with bovine serum albumin (BSA) 0.1 mg/mL as proteic substrate and some pHs buffers. The preliminary results showed that the extracellular peptidases were able of hydrolizate the proteic substrate in pHs 3 and 12, and their proteolytic inhibitions were obtained with EO concentration 0.1 µL/mL, suggesting a possible target that justify the antifungal activity this essential oil.

[1] N Mimica-Dukic et al. (2004) J. Agric. Food Chem. 52: 2485-2489.

[2] A Spreer et al. (2006) Medical Mycology. 44: 723-731.

[3] Clinical and Laboratory Standards Institute. (2008). M38-A2. Clinical and Laboratory Standards Institute, Wayne, PA.

[4] M Buroker-Kilgore, KKW Wang (1993) Anal Biochem 208: 387-392.

**P 167. Chemical composition of essential oils of *Ocimum gratissimum* L (Lamiaceae) from Limpopo province, South Africa: A comparative analysis of leaf and stem oil**

Theka T<sup>1</sup>, Mashimbye MJ<sup>2</sup>

<sup>1</sup>Department of Chemistry, University of Cape Town, Cape Town, South Africa,

<sup>2</sup>Department of Science and Technology, Pretoria, South Africa,

takalani.theka@gmail.com

Keywords: *O. gratissimum* L; Lamiaceae; essential oil; South Africa

*Ocimum gratissimum* L is a perennial herb or shrub with medicinal properties [1,2]. It is found in tropical countries including Limpopo Province, South Africa [2]. Our interest arose because the aromatic smell of crushed leaves and stem is used by locals for warding off mosquitoes [3]. Fresh leaves and stem were collected (flowering period, June-July) from Ha-Mutsha. The essential oils were extracted using hydrodistillation and analyzed by GC-MS. The percentage yield of the essential oils were 0.78% and 0.14% (w/w) for leaf and stem oils, respectively. Both leaf and stem oils have a high content of limonene, neral and geraniol (Table 1). A closer look reveals that the non-oxygenated monoterpenes *o*-cymene (3.36%), eugenol (2.37%), germacrene-D (2.88%) and piperitenone (19.04%) are only found in the leaf oil and *d*-3-carene (1.73%), *α*-terpinolene (1.27%) and *α*-copaene (1.24%) are only found in the stem oil. The components that are only found in the stem oil occurs in lesser amounts. The oil is characterized by the absence of thymol, ethyl cinnamate and linalool which have been reported in similar plant species [4].

Table 1. Chemical composition of *O. gratissimum* L leaf and stem oils from Ha-Mutsha

Compound	Leaf oil	Stem oil
$\Delta$ -3-Carene		1.73
<i>o</i> -cymene	3.36	
Limonene	24.80	36.11
<i>cis</i> $\beta$ -Ocimene	6.15	19.81
<i>trans</i> $\beta$ -Ocimene	3.64	1.44
$\alpha$ -Terpinolene		1.27
Nerol	2.60	1.13
Neral	14.39	17.16
Geraniol	1.98	1.26
Geraniol	15.21	16.71
Piperitenone	19.04	
Eugenol	2.37	
$\alpha$ -copaene		1.24
$\beta$ -Caryophyllene	3.59	2.14
Germacrene-D	2.88	

Acknowledgements: Prof Magwa ML (Forthare University) for assistance with GC-MS. Mr Paulus Sebothoma (National Botanical Institute, Pretoria) for plant identification. National Research Foundation for financial assistance. The study was conducted while the authors were still at the University of Venda.

1. Adebolou TT, Oladimeji SA (2005) Afr. J. Biotech. 4(7): 682-684.
2. Rietif E, Herman PPJ (1997) Plants of the Northern Provinces of South Africa: keys and diagnostic characters. 1<sup>st</sup> ed Struik Publishers, Pretoria, pp. 495-642.
3. Mabogo DEN (1990) Thesis, University of Pretoria, South Africa
4. Orwa C et al. (2009) Agroforestry Database: a tree reference and selection guide version 4.0



**P 168. Effect of essential oil of orange peel on germination, growth and production of soybean**

Bruno Reis, Hingrid Ariane da Silva; Heitor Vieira Mateussi; Carlos Moacir Bonato

*State University of Maringa, Brazil*

*cmbonato@hotmail.com*

Keywords: *Glycine max*; photosynthesis; allelopathy; mineral oil

Many essential oils have bioherbicide effect. Their effects on germination and growth occur in very low doses. However, there are commercial formulations in which essential oils are used to replace mineral oil in the agriculture. Thus, the essential oil may be used as alternative to petroleum oils. Therefore, the objective of this study was to evaluate the inhibition effect of a commercial formulation containing 5% essential oil of orange peel on the germination, growth and yield of soybeans. Growth, photosynthesis rate and yield were evaluated in greenhouse condition with nine treatments (commercial mineral oil Assist (BASF), Nimbus (Syngenta) to 0.5% with 2 and 3 applications, and commercial essential oil of orange peel Orobor (Oro Agri) at 0.1 and 0.4% with 2 and 3 applications). Germination test was evaluated with 200 seeds in germitest paper at 25°C for 7 days [1] at the concentration of 0, 2.5, 5, 7.5 and 10% of commercial essential oil of orange peel.

Essential oil of orange peel affected all the germination variables but did not affect the biometric variables, photosynthesis and yield. These results suggest that essential oil of orange peel presented as a potential alternative to replace mineral oil in agriculture.

1. Brasil. Ministério da Agricultura. Regras para análise de sementes. Brasília: MINISTÉRIO DA AGRICULTURA, PECUÁRIA E ABASTECIMENTO, 2009.

**P 169. Phytochemical constituents of leaf oil of *Lantana camara* L (Verbernaceae) from Limpopo province, South Africa**

Theka T<sup>1</sup>, Mashimbye MJ<sup>2</sup>

<sup>1</sup>Department of Chemistry, University of Cape Town, Cape Town, South Africa;

<sup>2</sup>Department of Science and Technology, Pretoria, South Africa,

takalani.theka@gmail.com

Keywords: *Lantana camara* L; Verbernaceae; essential oil; limonene;  $\beta$ -caryophyllene; South Africa

We have recently completed phytochemical investigation of *L. camara* L, a much-branched shrub that is widespread in summer to winter [1]. Its purplish berries are eaten, mostly by young people [2]. Our interest in its leaf oil constituents arose mainly because the leaves are used traditionally for treatment of eye injuries. Fresh leaves were collected from University of Venda campus. The oil, with percentage yield of 0.06% (w/w), was extracted using hydrodistillation and analyzed by GC-MS. The cyclic monoterpenic hydrocarbon limonene is the major component of the oil comprising 46.74% of the total oil while the bicyclic sesquiterpene  $\beta$ -caryophyllene content of the oil is 17.14%. Six other essential oil constituents appear in lesser amounts:  $\alpha$ -curcumene (5.36%), *p*-cymene (4.21%),  $\beta$ -elemene (4.47%),  $\alpha$ -humulene (4.52%), *cis*- $\beta$ -ocimene (4.73%) and *trans*- $\beta$ -ocimene (6.64%). A comparison with the constituents of various essential oils of *L. camara* L with different origin will also be given.

Acknowledgements: Prof Magwa ML (Forthare University) for assistance with GC-MS analysis. Mr Tshisikhawe MP (University of Venda) for plant identification. National Research Foundation for financial assistance. The study was conducted while the authors were still at the University of Venda.

1. Germishuizen G, Fabian A (1997) Wild flowers of Northern South Africa. 1<sup>st</sup> ed Fernwood Press, Cape Town, pp 342-444
2. Mabogo DEN (1990) Thesis, University of Pretoria, South Africa

**P 170. Phytotoxic activity of essential oils from three *Copaifera* L. species**

Gurgel ESC<sup>1</sup>, Souza APS<sup>2</sup>; Santos JUM<sup>3</sup>; Lucas FCA<sup>4</sup>; Mendonça MS<sup>5</sup>; Zoghbi MGB<sup>1</sup>

<sup>1</sup>Museu Paraense Emílio Goeldi, Coordenação de Botânica;

<sup>2</sup>Empresa Brasileira de Pesquisa Agropecuária, Embrapa Amazônia Oriental;

<sup>3</sup>Universidade Federal Rural da Amazônia;

<sup>4</sup>Universidade do Estado do Pará, Departamento de Ciências Naturais;

<sup>5</sup>Universidade Federal do Amazonas

copaldoc@yahoo.com.br

Keywords: Phytotoxic; *Copaifera duckei*; *C. martii*; *C. reticulata*; inhibitory effects.

The rich and diversified Amazonian flora represents an excellent resource for new chemical structures. In this work, the chemical composition of the essential oils from leaves and stems of *Copaifera duckei* Dwyer, *C. martii* Hayne and *C. reticulata* Ducke (Leguminosae – Caesalpinioideae) were characterized. The phytotoxic activity these essential oils have on seed germination, and development of root and hypocotyl of the pasture weeds *Mimosa pudica* L. (malícia) and *Senna obtusifolia* (L.) H. S. Irwin & Barneby (mata-pasto) was analyzed. Inhibitory effects were more intense on root development and less intense on seed germination. *Mimosa pudica* tended to be more sensitive to phytotoxic effects than *S. obtusifolia*. Leaf oils presented a greater potential to inhibit root and hypocotyl development, while stem preferentially inhibited seed germination, although in some cases these differences were not statistically significant. The oils of leaves had a greater number of constituents when compared to those of stem, especially from *C. martii* which could justify the differences observed in the intensity of the phytotoxic effect between stems and leaves. Among the constituents identified, only d-cadinene and linalool had been previously related to phytotoxic activity.

**P 171. Effect of different culture media on the essential oil composition of *in vitro* grown *Thymus caespititius***

Mendes MD, Figueiredo AC; Trindade H

*Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portugal*

*mdmendes@fc.ul.pt*

Keywords: *Thymus caespititius*; essential oils; chemotypes; culture media

*Thymus caespititius* Brot. (Lamiaceae) is an endemic species of the NW Iberian Peninsula and of the Azores and Madeira archipelagos. These plants possess essential oils rich in thymol-, carvacrol-,  $\alpha$ -terpineol-, sabinene- or carvacrol/thymol [1]. Given the interest on this species' essential oil, *in vitro* cultures of selected *T. caespititius* chemotypes were established. In the present study, the effect of different culture media on the essential oil composition of *in vitro* grown *T. caespititius*, was assessed.

Five *T. caespititius* genotypes corresponding to three distinct chemotypes, carvacrol (C), carvacrol/thymol (CT1 and CT2) and sabinene/carvacrol (SC1 and SC2) were used. The essential oils were isolated by hydrodistillation and analyzed by GC and GC-MS as in [2].

*In vitro* cultures of *T. caespititius* were established as in [3]. Six months following culture initiation, the explants were transferred to MS medium [4] with reduced growth regulators for shoot elongation. Prior to the determination of the volatiles composition, the cultures were kept, for at least three subcultures, in each of the two culture media studied, MS and SH [5].

The different nutrient composition of the two media altered quantitatively the composition of the essential oils of the C and CT plantlets. Carvacrol, thymol, carvacryl acetate and thymyl acetate were the main components of the essential oils from MS grown plantlets. In the essential oils from the plantlets grown on SH medium a decrease of the relative amount of the acetate compounds and an increased of  $\gamma$ -terpinene and *p*-cymene was observed. These components are biosynthetically related to carvacrol and thymol, which synthesis proceeds from  $\gamma$ -terpinene to *p*-cymene.

No differences were observed between the SC plantlets essential oils grown in MS and SH medium. Sabinene was the main component identified in these oils, followed by carvacrol.

Acknowledgements: Partially funded by FCT, under PTDC/AGR-GPL/101334/2008. Marta D. Mendes is grateful to FCT for the PhD grant SFRH/BD/60244/2009.

1. AC Figueiredo et al. (2008) *Curr Pharm Des* 14: 3120-3140.
2. MD Mendes et al (2011) *Ind Crops Prod* 33: 710-719
3. MD Mendes et al (2010) *Acta Hort.* 860: 215-218
4. T Murashige, F Skoog (1962) *Physiol Plant.* 15: 473-497.
5. RU Schenk, AC Hildebrandt (1972) *Can J Bot.* 50: 199-204.

**P 172. Essential oil composition from twenty six *Eucalyptus* taxa from Mata Experimental do Escaroupim (Portugal)**

Sena I<sup>1</sup>, Faria JMS<sup>2</sup>; Sanches J<sup>3</sup>; Trindade H<sup>2</sup>; Pedro LG<sup>2</sup>; Barroso JG<sup>2</sup>; Figueiredo AC<sup>2</sup>

<sup>1</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portuga

<sup>2</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portuga

<sup>3</sup>Autoridade Florestal Nacional, Direcção Regional de Florestas de Lisboa e Vale do Tejo, 2001-901 Santarém, Portugal

icsena@fc.ul.pt

Keywords: Myrtaceae; *Eucalyptus* spp.; essential oils; cluster analysis; GC GC-MS; Escaroupim

*Eucalyptus* is one of the world's most important and most widely planted genera. This economically important Myrtaceae genus includes more than 700 species [1]. *Eucalyptus* are an important source of timber, pulpwood and for the production of essential oils, used for medicinal and pharmaceutical purposes [2]. To a minor extent, some are used as ornamental trees. The Mata Experimental do Escaroupim (MEE), located in Salvaterra de Magos, Portugal, includes an *arboretum*, harbouring more than 125 different species of this genus [3]. In continuation of our work on sixteen *Eucalyptus* species [4], and integrated in a study that aims at evaluating essential oils as nematotoxics, we herewith report on the composition of essential oils isolated from additional ten *Eucalyptus* taxa grown in MEE.

The essential oils were isolated by hydrodistillation, analysed by GC and GC-MS and the percentage composition of the essential oils was used to determine the relationship between different samples by cluster analysis as in [4]. Cluster analysis including previously [4], and presently studied essential oils samples showed two poorly correlated clusters ( $S_{\text{corr}} < 0.04$ ). Cluster I that included twenty four out of the twenty six taxa showed several subclusters. Cluster Ia, with twenty two samples from eighteen species (*E. bosistoana*, *E. botryoides*, *E. camaldulensis*, *E. cinerea*, *E. cordieri*, *E. ficifolia*, *E. gigantea*, *E. globulus*, *E. hemiphloia*, *E. macarthurii*, *E. pauciflora*, *E. piperita*, *E. polyanthemos*, *E. radiata*, *E. saligna*, *E. smithii*, *E. urophylla* and *E. viminalis*) and Cluster Ib (two samples of *E. citriodora*) were also poorly correlated ( $S_{\text{corr}} < 0.10$ ). All oils from Cluster Ia showed variable amounts  $\alpha$ -pinene (1-94%), 1,8-cineole (1-83%) and limonene (1-41%), without citronellal, whereas Cluster Ib was citronellal rich (36-47%), with lower  $\alpha$ -pinene (1%), 1,8-cineole (0.1-11%) and limonene (0.2-0.3%) contents. Cluster II which included only the two *E. dives* samples was characterized by a high piperitone content (40-55%), which was absent to 0.5% in the other taxa.

Acknowledgments: Jorge Faria is grateful to the FCT for the PhD grant SFRH/BD/43738/2008. This study was partially funded by FCT, under research contract PTDC/AGR-CFL/117026/2010.

1. Menut C et al. (1995) *J. Agric. Food Chem.* 43: 1267–1271.

2. Ghisalberti E.L. (1996) *Phytochemistry* 41: 7–22.

3. Goes E (1977) Os eucaliptos (ecologia, cultura, produção e rentabilidade), Portucel.

4. Faria JMS et al. (2011) *Acta Horticulturae* 925: 61-66.

**P 173. The phytotoxic effects of the secondary metabolites myrthenal, carvone and carvacrol**

Martínez-Peñalver A, Costas-Gil A, Díaz-Tielas C, Graña E, Reigosa MJ, Sánchez-Moreiras AM

University of Vigo, Vigo, Spain.

anampe3@uvigo.es

Keywords: myrthenal; carvone; carvacrol; phytotoxicity; microtubules; *Arabidopsis*

Myrthenal, carvone and carvacrol are natural compounds commonly found in essential oils of plants of the families Apiaceae and Lamiaceae [1, 2, 3]. These secondary metabolites have been found to possess antibacterial and fungicide activity [3, 4], but their phytotoxic potential has been poorly investigated. Therefore the phytotoxicity of carvone, carvacrol and myrthenal was tested on *Arabidopsis thaliana* (L.) Col-0, a model plant broadly used in phytotoxic studies [5]. *Arabidopsis* seedlings were treated with 0, 50, 100, 200, 400, 800 and 1200  $\mu\text{M}$  of carvone, carvacrol or myrthenal and root length was measured 15 days after allelochemical exposition. A strong inhibition and highly significant effect were detected on the development and growth of *A. thaliana* roots with very low  $\text{IC}_{50}$  (compound concentration required for 50% inhibition). Root thickness and presence of root hairs were analyzed under a magnifier and a left-handed growth, which suggests microtubule alteration [6, 7], was observed in the treated roots. Immunofluorescence assays were done to confirm microtubule malformations after carvone, carvacrol and myrthenal treatments. The results suggest the different mode of action of these metabolites and confirm the multiple targets on the mode of action of natural compounds.

1. R.J.W. Lambert (2001) J. Appl. Microbiol. 91: 453-462.
2. Luiz Fernando Rolim de Almeida (2010) Molecules, 15: 4309-4323.
3. K. Oosterhaven et al (1995) Industrial Crops and Products
4. 23-31. 4. Ilkka M. Helander (1988) J. Agric. Food Chem. 46: 3590-3595.
5. M Pennacchio et al. (2005) J. Chem. Ecol. 31: 1573-1561.
6. I Furutani et al. (2000) Development 127: 4443-4453.
7. D Chaimovitsh et al. (2010) The Plant J. 61: 399-408.

**P 174. *Ammoides verticillata* and *Thymus ciliatus* oils from Algeria: chemical composition, antimicrobial and antioxidant activities**

Tefiani C<sup>1</sup>, Youcefi F<sup>1</sup>; Riazi Ali<sup>2</sup>; Aazza S<sup>3</sup>; Gago C<sup>3</sup>; Faleiro M<sup>4</sup>; Pedro L<sup>5</sup>; Barroso JG<sup>5</sup>; Figueiredo AC<sup>5</sup>; Miguel MG<sup>3</sup>

<sup>1</sup>Département des Sciences Agronomique et des Forêts, Faculté des Sciences de la Nature et de la Vie et des Sciences de la Terre et de l'Univers, Université Tlemcen 13000 Algérie;

<sup>2</sup>Laboratoire des Microorganismes Bénéfiques, des Aliments Fonctionnels et de la Santé, Université Abdelhamid Ibn Badis, 27000 Mostaganem 13000 Algérie; <sup>3</sup>Universidade do Algarve, Faculdade de Ciências e Tecnologia, Departamento de Química e Farmácia, Instituto de Biotecnologia e Bioengenharia, Centro Biotecnologia Vegetal, Campus de Gambelas 8005-139 Faro, Portugal; <sup>4</sup>Universidade do Algarve, Faculdade de Ciências e Tecnologia, Departamento de Biologia e Bioengenharia, Instituto de Biotecnologia e Bioengenharia, CBME, Campus de Gambelas 8005-139 Faro, Portugal; <sup>5</sup>Universidade de Lisboa, Faculdade de Ciências de Lisboa, Departamento de Biologia Vegetal, Instituto de Biotecnologia e Bioengenharia, Centro de Biotecnologia Vegetal, C2, Piso 1, Campo Grande, 1749-016 Lisboa, Portugal;

choukri13@gmail.com

Keywords: *Ammoides*, *Thymus*, Algeria, essential oils, biological activities

*Ammoides verticillata* (Apiaceae) is an Algerian endemic species which flowers are rich in essential oils [1]. A great variability on the chemical composition of the essential oils has been reported depending on the part of the plant as well as on the developmental stage [2,3]. The chemical composition and antimicrobial activity of *Thymus ciliatus* (Desf.) Benth. ssp. *eu-ciliatus* Maire essential oil from Algeria was reported by Bousmaha-Marroki et al [4]. Carvacrol dominated the oil which may explain the good antimicrobial activity detected. In the present work, the chemical composition, antioxidant and antimicrobial activities of the essential oils of those two species collected in Algeria were studied. The essential oils were isolated from the aerial parts by hydrodistillation, and analyzed by Gas Chromatography and Gas Chromatography-Mass Spectrometry as in [5].

Cumin alcohol (44%), *p*-cymene (18%), limonene (14%), thymol (11%) and *g*-terpinene (7%) dominated in the *A. verticillata* oil; whereas carvacrol (71%) predominated in the *T. ciliatus* oil.

The antibacterial activity of the essential oils of *A. verticillata* and *T. ciliatus* was determined by agar diffusion against *Escherichia coli*, *Salmonella enterica* serovar Thyphimurium, methicillin-resistant *Staphylococcus aureus* and *Listeria monocytogenes*. The tested bacteria were more susceptible to the essential oil of *T. ciliatus*.

*A. verticillata* oil was the most effective for scavenging the free radicals 2,2-diphenyl-1-picrylhydrazyl (DPPH); 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS); and peroxy radicals. The same oil was also the best for prevent lipid peroxidation when measured through the TBARS (thiobarbituric acid reactive species).

[1] Oumessaad T (2011) *Current Opinion in Biotechnology* 22S: S15–S152.

[2] Bekhechi C et al (2010) *Natural Product Communications*, 5: 1107-1110.

[3] Mohagheghzadeh et al (2007) *Food Chemistry*, 100: 1217-1219.

[4] Bousmaha-Marroki L et al (2007) *Journal of Essential Oil Research*, 29: 490-493.

[5] MD Mendes et al. (2011) *Industrial Crops and Products* 33: 710-719.

**P 175. Essential oil variation of *Lippia origanoides* with occurrence in the Carajás national forest, PA, Brazil**Alcy Favacho Ribeiro<sup>1</sup>, Eloisa Helena A. Andrade<sup>1</sup>; José Guilherme S. Maia<sup>1</sup><sup>1</sup>Universidade Federal do Pará

gmaia@ufpa.br

Keywords: *Lippia organoides*, essential oil variation, circadian activity

The harvest time can be an important aspect in the production of essential oils. The interactions of environmental conditions that occur throughout the day can influence directly or indirectly the processes of secondary metabolism that result in quantitative and qualitative variations in the essential oils [1]. *Lippia origanoides* Kunth is a small shrub growing up to 3 m, with different aromatic characteristics and some known chemotypes. The species occurs in countries of Central and South America, especially in the Brazilian Amazon. It is used for culinary and medicinal purposes. The aim of this study was to evaluate the circadian activity of the plant, based on the analysis of its leaf essential oil. The leaf samples were collected between 9 h and 18 h, during 3 days (triplicate) in the Carajás National Forest, state of Pará, Brazil, September 2011. The leaf samples were hydrodistilled using Clevenger apparatus and the oils analyzed by GC and GC-MS, both equipped with DB-5ms capillary column (30 m x 0.25 mm; 0.25  $\mu$ m film thickness), temperature programmed to 60-240 °C (3 °C/min) and carrier gas helium and nitrogen, respectively. The leaf oil yields were as follows: 9 h (3.1%), 12 h (2.8%), 15 h (2.8%) and 18 h (2.5%). The main constituents (over 3%) identified at 9 h were *p*-cymene (27.0%), carvacrol (11.8%), (*E*)-methyl cinnamate (8.3%), methyl thymol ether (5.2%),  $\alpha$ -pinene (4.6%), (*E*)-nerolidol (3.9%) and 1,8-cineol (3.5%); at 12 h were (*E*)-methyl cinnamate (26.5%), (*E*)-nerolidol (16.9%), 1,8-cineol (9.9%),  $\alpha$ -pinene (5.0%) and *p*-cymene (4.0%); at 15 h were (*E*)-nerolidol (14.7%), 1,8-cineol (12.3%), (*E*)-methyl cinnamate (10.8%), *p*-cymene (9.1%),  $\alpha$ -pinene (6.9%) and  $\alpha$ -phellandrene (5.0%); and at 18 h were 1,8-cineol (17.6%), *p*-cymene (9.1%), (*E*)-nerolidol (6.5%),  $\alpha$ -pinene (4.7%), methyl thymol ether (4.7%), (*E*)-caryophyllene (3.9%) and  $\alpha$ -copaene (3.4%). In addition, the main components (over 3%) found in the oil of fine stems were (*E*)-nerolidol (25.8%), 1,8-cineol (13.3%), (*E*)-caryophyllene (9.3%),  $\alpha$ -copaene (7.4%) and  $\alpha$ -pinene (3.6%), while in the oil of flowers were 1,8-cineol (25.1%), (*E*)-caryophyllene (12.1%),  $\alpha$ -copaene (5.1%) and  $\alpha$ -pinene (4.5%). The constituents with greater variation during the day were *p*-cymene, (*E*)-methyl cinnamate, 1,8-cineol and (*E*)-nerolidol, specially the carvacrol which has a maximum value at 9 h. The study of seasonal activity in *L. origanoides* is ongoing.



**P 176. Detection of shiikuwasha juice adulteration by calamondin juice using volatile components**

Kenta Yamamoto<sup>1</sup>, Ayami Yoshimoto<sup>1</sup>; Ayumi Yahada<sup>1</sup>; Kumi Sasaki<sup>1</sup>; Hideaki Ohta<sup>1</sup>

<sup>1</sup>Nakamura Gakuen University

s10d101@nakamura-u.ac.jp

Keywords: Citrus volatiles; essential oil;  $\gamma$ -terpinene; adulteration; Shiikuwasha; Calamondin

Shiikuwasha (*Citrus depressa* Hayata) has been a very popular fruitlet as a flavor enhancer in Okinawa, Japan, and contains large amount of nobiletin reported to show anti-tumor activities [1] in the edible part of this citrus [2]. Therefore, market of Shiikuwasha products has grown rapidly by today's health-conscious consumers. Although Shiikuwasha farmers could not keep up with the supply of this fruit, Shiikuwasha juice adulterated with Calamondin (*Citrus madurensis* LOUR.) juice produced in Taiwan and Philippines is widely commercialized. The objective of this work was to investigate the volatile compounds to detect simply the Shiikuwasha juice adulterated with Calamondin juice.

Shiikuwasha juice, calamondin juice and 10 commercial shiikuwasha juice was used for the experiment. One milliliter of shiikuwasha juice was placed in a vial. The vial solution was held at 40°C, and solid phase microextraction (SPME) fiber was introduced into the headspace of vial and kept for 20 minutes. Heating of GC column-oven was stopped and a column head was dipped into liquid nitrogen to collect the volatiles in splitless mode. Then, the SPME fiber was introduced into the injector of GC and kept there for 7 minutes with cryofocusing. Headspace (HS) gas was analyzed by GC and GC/MS.

Chromatograms with high resolution were obtained by HS-SPME-cryofocusing, and 39 aromatic components were identified or presumed. Calamondin contained a slight amount of  $\gamma$ -terpinene (1.7%: composition rate) as compared with shiikuwasha (17.3%), it is supposed  $\gamma$ -terpinene detected shiikuwasha juice adulterated with calamondin juice. We obtained  $\gamma$ -Terpinene ratio ( $\gamma$ -terpinene peak area/total peak area).  $\gamma$ -Terpinene ratio was with a 0.07-0.11 % range for 3 commercial juices, and these juices were doubt adulterations with calamondin juice.

$\gamma$ -Terpinene was a useful volatile marker component to detect shiikuwasha juice adulterated with calamondin juice.

[1] A Murakami *et al.* (2000) *Cancer Res.*, 60, 5059-5066.

[2] S Kawaii *et al.* (1999) *J. Agric. Food Chem.*, 47, 3565-3571.

**P 177. Antifungal activity of some essential oils on pathogenic fungi isolated from medicinal plants**

Savikin K<sup>1</sup>, Stevic T<sup>1</sup>, Godjevac D<sup>2</sup>, Cujic N<sup>1</sup>, Beric T<sup>3</sup>, Stankovic S<sup>3</sup>

<sup>1</sup>Institute for Medicinal Plants Research „Dr Josif Pancic“, Tadeusa Koscuska 1, Belgrade, Serbia;

<sup>2</sup>Institute of Chemistry, Technology and Metallurgy, University of Belgrade, Njegoševa 12, 11000 Belgrade, Serbia;

<sup>3</sup>Institute for Medicinal Plants Research „Dr Josif Pancic“, Tadeusa Koscuska 1, Belgrade, Serbia;

<sup>4</sup>Faculty of Biology, University of Belgrade, Studentski trg 3, Belgrade, Serbia

ksavikin@mocbilja.rs

Keywords: essential oils; medicinal plants; antifungal activity

Microbiological quality of the medicinal plants very often is not good enough to fulfill official criteria (standards) because of the presence of various pathogenic fungi. On the other side, aromatic plants presents significant source of essential oils with various microbiological activities. In this study, the antifungal activity of four widely used essential oils (Rosemary, Oregano, Tea tree and Eucalyptus oil) were determined against eight pathogenic fungi: *Aspergillus flavus*, *A. niger*, *Penicillium sp.*, *Alternaria alternata*, *Chaetomium sp.*, *Gliocladium roseum*, *Trichotecium roseum* and *Phomopsis sp.* All tested fungi were previously isolated and identified in our laboratory [1] from medicinal plants (*Mentha piperita*, *Urticaria dioica*, *Calendula officinalis*) that very often could have insufficient microbiological quality. Essential oils were commercially obtained (Frey + Lau, Germany). The microdilution method was used to establish minimal inhibitory concentrations (MIC) [2]. Standard antimycotic fluconazole was used as a positive control.

The most abundant compounds were limonene (89.9%), carvacrol (75.8%), terpinen-4-ol (40.7%) and 1,8-cineole (45.3%) detected in Eucalyptus, Oregano, Tee tree and Rosemary oil, respectively. All selected oils exhibited antifungal activity against tested pathogenic fungi. Oregano oil showed the highest antifungal activity, followed by Tea tree oil, Rosemary and Eucalyptus oil. Oregano oil inhibited the complete growth of different fungi at the concentration of 0.15 and 0.3 µl/ml. Tea tree oil completely inhibited the growth of *A. niger*, *T. roseum* and *Phomopsis sp.* at the 1.25 µl/ml, while *A. flavus* was most resistant to this oil. The lowest MIC for Rosemary oil was noticed against *Phomopsis sp.* and *A. alternata* (2.5 µl/ml), while *A. flavus* and *Penicillium sp.* were the most resistant. Among these oils, Eucalyptus oil was the least active with MIC values ranged from 5-15 µl/ml. According to our results, essential oils tested may be useful as alternative antifungal agents.

Acknowledgments: The authors acknowledge their gratitude to the Ministry of Education and Science of Serbia for the financial support (Project No. 46013).

1. T Stevic et al. (2012) Arch. Biol. Sci. 64: 49-58.

2. H Hanel and W Raether (1988) Mycoses. 3: 148-154.

**P 178. Volatile components of *Lamium amplexicaule* and *Lamium purpureum* growing wild in Huntsville, Alabama**

Chelsea Jones<sup>1</sup>, William N. Setzer<sup>1</sup>

<sup>1</sup>University of Alabama in Huntsville

cdj0006@uah.edu

Keywords: essential oil composition,  $\alpha$ -pinene,  $\beta$ -pinene, germacrene D, 1-octen-3-ol

The overall goal of the project is to test the hypothesis that *Lamium* species use phytotoxic allelochemicals in interplant competition. Both *Lamium amplexicaule* and *Lamium purpureum* are non-native perennial weeds, originally from Eurasia [1]. We hypothesize that these two *Lamium* species use chemicals to compete with other plant species [2,3].

Several samples of each plant species were collected from Huntsville, Alabama. The fresh aerial parts were hydrodistilled/extracted for four hours using a Lickens-Nickerson apparatus with continuous extraction with dichloromethane to give the volatile fractions. The volatiles were analyzed using gas chromatography – mass spectrometry (GC-MS) [4].

The volatile fraction of *Lamium amplexicaule* was composed largely of germacrene D (33%),  $\alpha$ -pinene (16%),  $\beta$ -pinene (11%),  $\alpha$ -thujene (7%), (*Z*)- $\beta$ -ocimene (4%), 1-octen-3-ol (4%), and germacrene B (4%). *Lamium purpureum* volatile, on the other hand, was dominated by  $\beta$ -pinene (16%),  $\alpha$ -pinene (15%), 1-octen-3-ol (15%), germacrene D (15%),  $\beta$ -ylangene (4%),  $\beta$ -elemene (4%), and (*E*)-phytol (4%).

The high concentrations of monoterpene and sesquiterpene hydrocarbons are probably responsible for the allelopathic effects of the *Lamium* species [5].

1. DJ Mabberly (1997) *The Plant Book*, 2<sup>nd</sup> Ed. Cambridge Univ. Press, UK, p. 389.
2. Y Fujii (2002) *In: Nonpesticide Methods for Controlling Diseases and Insect Pests*. Asian Product. Org., Tokyo, pp. 49-61.
3. S Shiraishi et al. (2002) *Weed Biol. Manag.* 2:133-142.
4. JE Kennedy et al. (2011) *Allelopathy J.* 27:111-122.
5. LCA Barbosa et al. (2007) *Quim. Nova*, 30:1959-1965.

**P 179. Essential oil of peppermint grown in the presence of methanolic leaf extract *Leonurus sibiricus* L.**

Bufalo J<sup>1</sup>, Almeida LFR<sup>1</sup>, Marques MOM<sup>2</sup>, Boaro CSF<sup>1</sup>

<sup>1</sup>Department of Botany, Institute of Biosciences (IB), UNESP - Univ. Estadual Paulista, Botucatu, Sao Paulo, Brazil;

<sup>2</sup>Agronomic Institute of Campinas, IAC, Campinas, Sao Paulo, Brazil

jenniferbufalo@yahoo.com.br

Keywords: essential oil; *Mentha piperita*; terpenes

Peppermint (*Mentha piperita* L.) is an important and commonly used as flavoring agent world-wide. Leaves of *Mentha* species are used as condiments, and essential oils of these plants, are used as flavorings for foods and beverages, fragrances, and fungicides or insecticides in many pharmaceutical and industrial products [1, 2]. This work evaluated the allelopathic activity of different concentrations of a methanolic extract of *Leonurus sibiricus* L. leaves in the chemical compositions of volatile oil of *M. piperita*. The plants were cultivated using complete Hoagland and Arnon [3] n° 2 nutrient solution. The control (0 mg L<sup>-1</sup>) contained only the complete nutrient solution in the absence of the methanolic extract from *L. sibiricus* leaves; the treatments were 25, 50 and 100 mg L<sup>-1</sup> of methanolic extract of *L. sibiricus* leaves in complete nutrient solution. Aerial part of *M. piperita* were collected at 77, 94, 108 and 120 days after transplanting (DAT) and subjected to hydrodistillation in a Clevenger-type apparatus. The composition of oils was analyzed in a gas chromatograph (GC/MS). Principal Component Analysis (PCA) was applied to substances, limonene, pulegone, menthone, menthofuran, menthol and menthyl acetate. The results revealed that at the beginning of the developmental cycle plants *M. piperita* showed a highest correlation with precursors of the metabolic pathway, such as limonene, pulegone and menthone, regardless of the presence leaf extract of *L. sibiricus*. However, with the development the plants the *M. piperita* showed a higher correlation with menthol when these plants were cultivated with 25 mg L<sup>-1</sup> of extract. Thus, the leaf extracts of *L. sibiricus* influenced in chemical compositions of volatile oil of *M. piperita*, because low concentrations of the extract ensured the quality of essential oil.

Acknowledgments: Financial support CNPq.

1. D Ram et al. (2006) Bioresour. Technol. 97: 886-893.

2. P Harrewijn et al. (2001) Kluwer Academic Publishers, London, U.K., 2001, p. 440.

3. DR Hoagland, DI Arnon (1950) California Agriculture Experimental Station, Berkeley, CA.

**P 180. Composition and antimicrobial activity of the essential oil of *Hyssopus seravschanicus* growing wild in Tajikistan**

Thompson RM<sup>1</sup>, Sharopov FS<sup>2</sup>; Kukaniev MA<sup>2</sup>; Satyal P<sup>1</sup>; Setzer WN<sup>1</sup>

<sup>1</sup>Department of Chemistry, University of Alabama in Huntsville;

<sup>2</sup>V. I. Nikitin Institute of Chemistry, Tajik Academy of Sciences

rmt0002@uah.edu

Keywords: *Hyssopus seravschanicus*; essential oil composition; antimicrobial activity; cis-pinocamphone

The volatile compounds were extracted from *Hyssopus seravschanicus* Pazij, a perennial, branched semi-shrub that grows wildly near the high ridge mountainous regions of Varzob, Northern Dushanbe in Tajikistan at an altitude approximately 2500 meters above sea level. The samples were analyzed using gas chromatography-mass spectrometry (GC-MS) in order to study the composition of the essential oil of hyssop. Eighty-seven chemical components of the essential oil were found and characterized representing approximately 95.4% of the oil. The most abundant components were cis-pinocamphone (57.0-88.9%),  $\beta$ -pinene (0.4-6.0%), 1,8-cineole (1.8-3.6%), camphor (0.5-4.0%), and spathulenol (0.1-5.0%). An antimicrobial screening was performed on five essential oil samples against Gram-positive bacteria, *Bacillus cereus* (ATCC No. 14579) and *Staphylococcus aureus* (ATCC No. 29213) as well as Gram-negative bacteria, *Pseudomonas aeruginosa* (ATCC No. 27853) and *Escherichia coli* (ATCC No. 10798). The essential oil showed notable antimicrobial activity against Gram positive *Bacillus cereus* and *Staphylococcus aureus*. cis-Pinocamphone was the most abundant chemical component in this study and in most others. It has been reported that cis-pinocamphone has antimicrobial effects which can be attributed to the low molecular weight and lipophilic constituents of the oil. In this way, the essential oil can alter the cellular composition of bacteria and influence normal cell growth. As a result, the essential oil of hyssop can be cultivated more abundantly and utilized for its antimicrobial activity.

**P 181. Neutralizing effects of two *Nectandra* species essential oils and extracts against *Bothrops neuwiedi* snake venom**

A.M. Torres<sup>1</sup>, N. Martinez<sup>2</sup>; F.J. Camargo<sup>1</sup>; G.A.L. Ricciardi<sup>1</sup>; A.A.I. Ricciardi<sup>1</sup>; M. Minteguiaga<sup>2</sup>; D. Lorenzo<sup>2</sup>; E. Dellacassa<sup>2</sup>

<sup>1</sup>Laboratorio Dr. Gustavo A. Fester, Facultad de Ciencias Exactas y Naturales y Agrimensura, Universidad Nacional del Nordeste, Corrientes, Argentina;

<sup>2</sup>Cátedra de Farmacognosia y Productos Naturales, Facultad de Química-Universidad de la República. Montevideo, Uruguay

edellac@fq.edu.uy

Keywords: *Nectandra* sp; snake venom; anti-snake venom; oil composition; chemotaxonomy

Venomous snakebite remains an important medical problem in both developing and developed countries. Through serum therapy, neutralization of the systemic toxic effects is usually reached, but that of local tissue injury is not. *B. neuwiedi diporus* (yará chica) is the species responsible for the major proportion of the viper accidents that occur in Argentina. Its venom is a complex mixture of hemorrhagic, clot-forming, anticoagulant, proteolytic, neurotoxic and edema-inducing toxins, among others. Although bothropic antiserum therapy has been shown to be effective against the systemic effects of envenomation, hitherto no antidote exists to combat the local damage.

The present study evaluated the potential hemolytic and anti-hemorrhagic effect of the aqueous and alcoholic extracts, and essential oil obtained from *Nectandra angustifolia* and *N. megapotamica* against *B. neuwiedi diporus* snake venom. Additionally, and in order to better characterize the chemical taxonomy of the population studied, we report the chemical composition, including the enantiomeric distribution of different mono and sesquiterpenes, of both *Nectandra* species essential oils. Venom was analyzed by SDS-PAGE [1]. Neutralization of *B. neuwiedi* venom enzymes by plant extracts and essential oils was measured using an indirect hemolytic assay on agarose-erythrocyte–egg yolk gel plate to define the minimum indirect hemolytic dose. Neutralization of coagulant activity was also evaluated. The components of the oil were analyzed as previously reported [2]. Comparison of fragmentation patterns in the mass spectra with those stored on the GC-MS database was also performed. Enantiomeric ratios of selected oil components were obtained by multidimensional GC using the experimental conditions previously described [2]. From these investigations, it may be concluded that, even when oils were active, the ethanol extract of *N. angustifolia* leaves showed a promising venom inhibition effect against *B. neuwiedi*.

[1] Camargo et al. (2011) BLACPMA 10: 429-434.

[2] Torres et al. (2011) Nat. Prod. Comm.6: 1393-1396.

**P 182. Potential use in cosmetic formulations and antimicrobial activity of essential oil of *Ocotea pulchella* (Nees) Mez (Lauraceae) and *Xylopiya aromatica* (Lam.) Annonaceae**

Silva EA<sup>1</sup>, Paula ACCFF<sup>2</sup>; Vicente NF<sup>3</sup>; Bernardes GG<sup>1</sup>; Alvarenga AA<sup>4</sup>

<sup>1</sup>Universidade José do Rosário Vellano- UNIFENAS/ Divinópolis- MG, Brazil.;

<sup>2</sup>Laboratório de Fisiologia Vegetal/ Botânica - Instituto Federal de Educação Ciência e Tecnologia - IFMG/ Bambuí, Brazil;

<sup>3</sup>E. Yoshia Takaoka – São José dos Campos – São Paulo- Brazil;

<sup>4</sup>Departamento de Biologia, Universidade Federal de Lavras, UFLA - Lavras; Brazil

anacard4@yahoo.com.br

Keywords: essential oil, antimicrobial activity, pharmaceutical and cosmetic formulations

The aroma and fragrance industry is a billion-dollar world market which grows annually. Essential oils comprise the majority of compounds used by these industries. Each component of an essential oil has a characteristic profile and each aroma and flavor is one combination of its representatives' profiles. The essential oils have been showing effective in the control of several microorganisms (Santos *et al.*, 2008). The main objective of this work was to evaluate the antimicrobial activity and the technological characteristics of the oil in cosmetics formulations. Although, the chemical composition will be performed can be verified another study of this subject by some authors (Botega *et al.* 1993. Potenza, 1999a, 2006, Pontes *et al.* 2007 and Raggi, 2008). The aerial parts of the plant were collected in the proximities of the city of Divinópolis - Minas Gerais State, area that prevails ecosystems of cerrado. Essential oil of leaves was extracted by hydrodistillation (Clevenger apparatus). Aliquots of those oils were submitted the tests of antimicrobial activity being used Brain Heart Infusion Agar (BHI) medium for *Staphylococcus aureus*, *Bacillus cereus*, *Pseudomonas aeruginosa* and Sabouraud medium for *Candida albicans*. The tested formulations were no-ionic cream-base and antimicrobial ointment. The obtained oil of *Ocotea pulchella* had antimicrobial activity against *C. albicans*, *B. cereus* and *P. aeruginosa*. The essential oil of *Xylopiya aromatica* it was active against *S. aureus* and *C. albicans*. The essential oil of *Ocotea pulchella* it was classified as a transparent liquid to milky appearance. For *Xylopiya aromatica* the essential oil shows a limpid and slightly astringency liquid. There was verified compatibility of the oils with the different ones tested formulations. These partial results stimulate searches for new compounds with potential cosmetics and pharmaceutical use.

Acknowledgement – FAPEMIG, UNIFENAS/ Divinópolis- MG, IFMG/ Bambuí

1. Patrícia O. Santos *et al.* (2008) Chemical composition and antimicrobial activity of the essential oil of *Hyptis pectinata* (L.) Poit. Quím. Nova vol.31 n<sup>o</sup>.7 São Paulo.
2. Botega *et al.* Phytochemistry, v.32, n.5, p.1331-3, 1993. Potenza ET AL., M.; *Arq. Inst. Biol.* 66, 31. 1999a, \_\_\_\_, 2006;
3. Pontes *et al.*, Quim. Nova, V. 30, No. 4, 838-841, 2007;
4. Raggi, Instituto de Botanica da Secretaria do Meio Ambiente Dissertação de mestrado (2008)

**P 183. Antimicrobial activity of the essential oil extracted from *Thymus vulgaris*, *Ocimum basilicum* (Lamiaceae) and *Piper aduncum* (Piperaceae)**

Alves RS<sup>1</sup>, Cardoso MG<sup>1</sup>; Piccoli RH<sup>2</sup>; Paula ACCFF<sup>3</sup>; Lacerda ICA<sup>4</sup>; Natividade MMP<sup>1</sup>

<sup>1</sup>Departamento de Ciências dos Alimentos, Universidade Federal de Lavras, UFLA Lavras; Brazil;

<sup>2</sup>Departamento de Ciências dos Alimentos, Universidade Federal de Lavras, UFLA Lavras; Brazil.;

<sup>3</sup>Laboratório de Fisiologia Vegetal/ Botânica - Instituto Federal de Educação Ciência e Tecnologia - IFMG/ Bambuí, Brazil;

<sup>4</sup>Departamento de Ciências Biológicas, Universidade Federal de Minas Gerais, UFMG - Belo Horizonte, Brazil.

anacard4@yahoo.com.br

Keywords: Essential oil, antimicrobial activity, Food-borne illness

Food-borne illness is a major international problem and an important cause of reduced economic growth. The contamination of the food supply with the pathogens and its persistence, growth, multiplication and/or toxin production has emerged as an important public health. *Piper aduncum* (Piperaceae), *Thymus vulgaris* and *Ocimum basilicum* (Lamiaceae) are seasoning products obtained from plants widely used in foods. Previous study were performed of the effect of essential oil composition and bioactivity of these plants related to toxicological and antimicrobial action (Sousa et al., 2008; Dandlen et al., 2011; Silva et al., 2012). However, there are few study of the synergism between essential oils on the antimicrobial activity against microorganism of the food-borne illness. The aim of this study was to evaluate *in vitro* activity of the essential oil extracted from *Thymus vulgaris*, *Ocimum basilicum* and *Piper aduncum* against *Salmonella enteritidis*, *Staphylococcus aureus* and *Escherichia coli*. The essential oil of this plants was extracted by hydrodistillation (Clevenger apparatus) about 2 hour and was analyzed by GC and GC/MS (SHIMADZU). The oils were mixed in different proportions and the antimicrobial activity were performed with isolated oils and the mixed oils from different proportions using 10µL of the extracts in Agar Muller Hinton medium with 10<sup>8</sup> UFC mL<sup>-1</sup> of each microorganism. The major components of the essential oil were tymol, carvacol, o-cimeno, γ-terpineno and linalol (*Thymus vulgaris*). Linalol, 1,8-cineol and eugenol (*Ocimum basilicum*) and linalol, (E) nerolidol and cariofilene oxyde (*Piper aduncum*). The essentials oil can be able to inhibit growth of microorganism caused of food-borne illness. The essential oil from this plants showed antimicrobial activity against *E. coli*, *Salmonella enteritidis* e *S. aureus*. The essential oil of *T. vulgaris* was the most active. Concerning the mixed essential oil were verified antagonist effect in some mixture and synergistic effect in others. The results of this study showed that these seasoning plants have an important activity isolated and/or in mixture, could be source for the discovery of new antimicrobial products.

Acknowledgement – FAPEMIG, UFLA/ Lavras- MG and IFMG/ Bambuí

1. Patrícia O. Santos et al. (2008) Chemical composition and antimicrobial activity of the essential oil of *Hyptis pectinata* (L.) Poit. Quím. Nova vol.31 n7 São Paulo. Sousa et al., 2008.,
2. Dandlen et al., Rev. Bras. farmacogn. V.21, 2011



**P 184. The use of vibrational spectroscopy to rapidly assess the quality of tea tree oil (*Melaleuca alternifolia*)**

Kamatou GPP, Tankeu YS, Vermaak I, Sandasi M, Viljoen AM

*Department of Pharmaceutical Sciences, Private Bag X680, Tshwane University of Technology, Pretoria, South Africa*

*kamatougpp@tut.ac.za*

Keywords: Tea tree oil, GC/MS, spectroscopy, NIR, MIR, correlation

Tea tree oil is a native Australian essential oil obtained mainly from the leaves of *Melaleuca alternifolia* (Myrtaceae). The oil is a common ingredient in many commercial detergents, cosmetics and herbal remedies. Tea tree oil is also incorporated as active ingredient in various topical formulations used to treat skin infections. The quality of the oil is usually determined by GC/MS based on the major constituents (terpinen-4-ol,  $\gamma$ -terpinene,  $\alpha$ -terpinene,  $\alpha$ -terpineol, terpinolene, limonene and 1,8-cineole). This analytical method is however time consuming and expensive. The need to identify a fast, accurate alternative method for routine analysis is thus important.

The study was aimed at investigating the possibility of using vibrational spectroscopy in combination with chemometric data analyses as alternative tool for the quality assessment of tea tree oil.

Samples of tea tree oil ( $n = 60$ ) obtained from different suppliers and natural populations were analysed using GC/MS as reference method. For spectroscopy analyses, Fourier transform NIR spectra of each oil was recorded on a NIRFlex N500 liquid cell spectrometer. The oil spectra were collected in the transmittance mode between the wavenumbers 10,000 and 4000  $\text{cm}^{-1}$ . Fourier transform MIR spectra of each oil was recorded in the wavenumber range 4000–550  $\text{cm}^{-1}$  on an alpha-P Bruker spectrometer. The spectral data combined with the reference data were analysed using chemometric analysis software (Simca P+, version 12.0).

The partial least squares (PLS) regression method was used to develop calibration models with good correlation coefficients ( $R^2$ ) based on both MIR and NIR spectra for terpinen-4-ol (MIR = 0.91; NIR = 0.78), terpinolene (MIR = 0.90; NIR = 0.80),  $\gamma$ -terpinene (MIR = 0.90; NIR = 0.89), limonene (MIR = 0.89; NIR = 0.88) and  $\alpha$ -terpineol (MIR = 0.85; NIR = 0.90). 1,8-Cineole had the highest  $R^2$  of greater than 0.96 for MIR and NIR. Generally, the error parameters (RMSEE and RMSEP) after external validation ( $n = 18$ ) were low (< 3%). The lower RMSEE and RMSEP values recorded demonstrated accurate predictions of compounds investigated.

The results have shown the potential of MIR and NIR spectroscopy as alternative methods to GC/MS for quality assessment of tea tree oil.

Acknowledgements: The financial support of the National Research Foundation (South Africa) and Tshwane University of Technology is gratefully acknowledged.

**P 185. The phytotoxic activity of the secondary metabolite eugenol**

Costas Gil A<sup>1</sup>, Martínez-Peñalver A<sup>2</sup>; Díaz-Tielas C<sup>2</sup>; Graña E<sup>2</sup>; Reigosa MJ<sup>2</sup>; Sánchez-Moreiras AM<sup>2</sup>

<sup>1</sup>University of Vigo, Vigo, Spain;

<sup>2</sup>University of Vigo, Vigo, Spain.

aitana@uvigo.es

Keywords: Eugenol; phytotoxicity; microtubules; *Arabidopsis*; essential oil; weed management

Eugenol is a natural allylbenzenic compound commonly found in essential oils of plants of the families Lauraceae, Mirtaceae and Miristicaceae [1]. Although the antiseptic, anesthetic and hepatotoxic activity of eugenol has been already demonstrated [2, 3, 4], its phytotoxic activity has been poorly investigated. The phytotoxicity of this essential oil was analyzed on *Arabidopsis thaliana* (L.) Col-0 seedlings at 50, 100, 200, 400, 800 and 1200 µM eugenol concentrations. The results showed a strong growth inhibition with the IC<sub>50</sub> (the concentration required for 50% inhibition) at 246 µM. The root structure, thickness and presence of root hairs was studied under a magnifier. Eugenol-treated roots showed left-handed growth, indicative of microtubule alteration [5, 6], which was confirmed by immunofluorescence. Ultra-structural analysis was performed by electron microscopy on 7 and 14 days eugenol-treated roots. The strong root growth inhibition, the alteration of microtubules and the changes at cellular level confirmed the high phytotoxicity of eugenol, a strong candidate for weed management.

1. Gislene G F Nascimento et al. (2000) Braz. J. Microbiol. 31:247-256.
2. David C Thompson et al. (1998) Tox. Appl. Pharmac. 149: 55-63.
3. K Dallmeier and E A Carlini (1981) Pharmacology 22:113-127.
4. H J D Dorman and S G Deans (2000) J. Appl. Microbiol. 88: 308-316.
5. I Furutani et al. (2000) Development 127: 4443-4453.
6. D Chaimovitch et al. (2010) The Plant J. 61: 399-408.

**P 186. Evaluation anti-*Rhizopus oryzae* activity and synergistic effect of *Hyptis pectinata* essential oil and its bioactive compounds**

Ana Cristina Rivas da Silva<sup>1</sup>, Catia Amancio Almeida<sup>1</sup>; Paula Monteiro Lopes<sup>1</sup>; Maria de Fátima Arrigoni-Blank<sup>2</sup>; Arie Fitzgerald Blank<sup>3</sup>; Péricles Barreto Alves<sup>4</sup>; Celuta Sales Alviano<sup>5</sup>; Daniela Sales Alviano<sup>5</sup>

<sup>1</sup>Instituto de Microbiologia Paulo de Góes; <sup>2</sup>Núcleo de Ciências Biológicas, Campus Professor Alberto Carvalho, Universidade Federal de Sergipe, Brazil; <sup>3</sup>Departamento de Engenharia Agrônômica, Universidade Federal de Sergipe, Brazil; <sup>4</sup>Federal University of Sergipe - Brazil; <sup>5</sup>Instituto de Microbiologia Paulo de Góes, Universidade Federal do Rio de Janeiro, Brazil

catiamancio@yahoo.com.br

Keywords: *Hyptis pectinata*; calamusenone; calamusenone oxidized analogous; dehydroaromadendrene; *Rhizopus oryzae*

In the last few years, the number of cases of zygomycosis has increased, especially among immunocompromised patients, although several authors have also reported infections in patients with unknown underlying conditions. Zygomycetes are a heterogeneous group of fungi with a wide antifungal susceptibility profile. Amphotericin B is the agent of choice to treat zygomycosis. However, its toxicity remains a problem and therefore alternative therapies [1]. Previous works described antimicrobial activity of CHCl<sub>3</sub> extract and also essential oil (EO) from leaves of *H. pectinata*, and only pyrones, and pectinolides A-C (1-3) and H, were isolated of CHCl<sub>3</sub> extract, and their anti-staphylococcal activity was confirmed [2, 3, 4, 5]. No antimicrobial investigation with EO compounds was made to elucidate its bioactivity. So the anti-*R. oryzae* activity of SAM002 genotype of *H. pectinata* EO was evaluated and several bioactive compounds were isolated and tested. The TLC and bioautography showed that three substances were active. After purification of the three substances and GC-MS analysis, they were identified as calamusenone, calamusenone oxidized analogous (COA), and dehydroaromadendrene (DHAD) [5]. The minimal inhibitory concentration (MIC) and minimal microbicidal concentration (MMC) of all these substances were also determined, confirming that *H. pectinata* EO and its purified substances exhibited fungicidal activity with MICs ranging from 375 to 1250 µg/mL [6]. In combination with amphotericin B (AMB) plus essential oil or calamusenone presented additive activity although a synergistic effect was detected when AMB was combined with the COA e DHAD [7]. Once the therapeutic dose of AMB used against zygomycetes is extremely toxic, our results demonstrate that both the oil and its bioactive are promising candidates when it associated with AMB, minimizing side effects in patients with zygomycosis.

Acknowledgement: CNPq, CAPES e FAPERJ.

[1] A Alastruey-izquierdo et al. (2009) Clin. Microbiol. Infec. 15(5): 71–76.

[2] AW Jones (2011) Drug Test. Anal. 3 (6): 337-44.

[3] R Crotea (2000) Rockville: Am. Soc. Plant Physiol. p. 1250-1318.

[4] H Van den Dool, PD Kratz (1963) J. Chrom., 11:463-471.

[5] RP Adams (2007) Allured Publishing Corporation, Carol Stream.

[6] Clinical and Laboratory Standards Institute (2008). M38-A2. CLSI, Wayne, PA.

[7] GB Zore et al. (2011) *Phytomed.* 18:1181–1190.

**P 187. Volatile compounds are involved in the phytotoxicity of *Vicia faba* used as green manure**

Lorena Álvarez-Iglesias<sup>1</sup>, Carlos Cavaleiro<sup>2</sup>; Lígia Salgueiro<sup>2</sup>; Manuel J. Reigosa<sup>1</sup>; Nuria Pedrol<sup>1</sup>

<sup>1</sup>Lab. Ecofisiología Vegetal, Faculdade de Biología, Universidade de Vigo.;

<sup>2</sup>Centro de Estudos Farmacêuticos / Faculdade de Farmácia, Universidade de Coimbra.

lorena.alvarez@uvigo.es

Keywords: Faba bean; essential oil; in vitro volatile assay; bioactivity; weed control

Cover crops grown as green manure are keystone in sustainable management of agricultural systems. Some of the crops used traditionally as green manure are known to possess allelopathic activity so they can provide natural weed control, in addition to their environmental services such as soil conservation and fertility building. Faba bean (*Vicia faba* L.) is a legume crop usually included in rotations with maize in temperate areas. Our previous *in vitro*, greenhouse and field studies revealed strong phytotoxic effects of *V. faba* on some problematic weeds (Álvarez-Iglesias *et al.* submitted). Nonetheless, references about the bioactivity of the volatiles emitted by faba bean on other plants are absent in literature. The aim of our study was to determine if the volatile compounds from complete blossom aerial parts of *V. faba*, as used for green manure, could take part in the observed global phytotoxic effects, as well as to determine the composition of its essential oil.

Biological activity of fresh blossom aerial parts of *V. faba* was evaluated against several species by placing fresh material at a dose of 2 g dw L<sup>-1</sup> air in sealed containers together with seeds and seedlings of some target species<sup>1</sup>. Seedling growth of the model species *Lactuca sativa* and the weeds *Amaranthus retroflexus* and *Digitaria sanguinalis* was reduced over 50 %; moreover, germination of *D. sanguinalis* was strongly inhibited.

In order to elucidate which compounds were potentially responsible of these effects, the essential oil from blossom aerial parts of *V. faba* was obtained by continuous water distillation / solvent extraction using a Likens-Nickerson type apparatus<sup>2</sup>. Analysis was carried out by combination of gas chromatography (GC) and gas chromatography-mass spectroscopy (GC/MS). Aliphatic compounds and oxygenated monoterpenes represented the main fractions. Several compounds with known biological activities, such as terpinen-4-ol, geraniol, and linalool were identified. These compounds, acting individually or by synergistic interactions<sup>3</sup>, could be partially responsible of the phytotoxic activity of *V. faba* against weeds observed in our previous investigations.

Acknowledgements: Thanks are due to Fundação para a Ciência e Tecnologia (FCT), POCI 2010/FEDER for financial support, as well as to the FPU grant from the Spanish Ministry of Education AP2008-03101.

1. JN Barney, AG Hay, LA Weston (2005) J Chem Ecol 31: 247-265.

2. ST Likens, GB Nickerson (1964) Am Soc Brew Chem Proc 5: 13-19

3. D Vokou *et al.* (2003) J Chem Ecol 29: 2281-2301.

**P 188. Fast-RP-HPLC/PDA analysis of oxygen heterocyclic compounds in citrus essential oils**

Russo M<sup>1</sup>, Torre G<sup>1</sup>; Bonaccorsi I<sup>1</sup>; Dugo P<sup>2</sup>; Mondello L<sup>2</sup>

<sup>1</sup>Dipartimento Farmaco-chimico, University of Messina, viale Annunziata, 98168 – Messina, Italy;

<sup>2</sup>Dipartimento Farmaco-chimico, University of Messina, viale Annunziata, 98168 – Messina, Italy;  
Università Campus Bio-Medico (CIR), via Alvaro del Portillo, 21, 00128 Roma, Italy

mrusso@unime.it

Keywords: coumarins, furocoumarins, fast-RP-HPLC, qualitative and quantitative analysis.

Oxygen heterocyclic components present in the non volatile fraction of cold pressed *Citrus* essential oils are mainly represented by coumarins, psoralens and polymethoxylated flavones.

An important role in the characterization of *Citrus* oils has been attributed to these components. In fact, qualitative and quantitative composition of the fraction is characteristic of each oil. Moreover, many pharmacological and toxicological activities have been demonstrated for most of them. These components exhibit strong absorption in the ultraviolet region ( $\lambda_{max}$  315 nm).

Due to their non volatile nature, liquid chromatography has been considered the ideal technique for their analysis. Most of the methods found in literature propose the RP-HPLC analysis of oxygen heterocyclic compounds in an analysis time higher than 40 minutes [1,2]. The aim of this work is to reduce the total analysis time under 15 minutes.

One of the two methods investigated in this work allows a total baseline separation and a correct quantification of all the oxygen heterocyclic compounds within 10 minutes with an Ascentis Express C18 column (50 x 4.6 mm x 2.7  $\mu$ m). The second one permits a good baseline separation of coumarins, furocoumarins and polymethoxyflavones within a 3 minutes analysis time, with the same column and under the same chromatographic conditions.

Both the two methods present numerous advantages: first, the total analysis time is greatly reduced thus allowing the characterization of a larger number of sample in the same day. Moreover, a conventional HPLC instrumentation can be used along with a lower amount of mobile phase with respect to conventional analysis.

Acknowledgments: The Project was funded by the “Italian Ministry for the University and Research (MIUR)” within the National Operative Project “Hi-Life Health Products from the industry of foods”. Project ID: PON01\_01499.

1. P Dugo et al. (2009) J. Agric. Food Chem. 57: 6543-6551.

2. M Russo et al. (2012) J. Ess. Oil Res. 24: 119-129.

**P 189. Exploiting the alternative selectivity of a novel medium-polar ionic liquid stationary phase in the gas chromatographic analysis of essential oils**

Ragonese C<sup>1</sup>, Sciarrone D<sup>2</sup>; Grasso E<sup>1</sup>; Dugo P<sup>3</sup>; Mondello L<sup>3</sup>

<sup>1</sup>Dipartimento Farmaco-chimico, University of Messina, viale Annunziata, 98168 Messina, Italy;

<sup>2</sup>Dipartimento Farmaco-chimico, University of Messina, viale Annunziata, 98168 Messina, Italy; Chromaleont s.r.l. A spin-off of the University of Messina, c/o University of Messina, viale Annunziata, 98168 Messina, Italy;

<sup>3</sup>Dipartimento Farmaco-chimico, University of Messina, viale Annunziata, 98168 Messina, Italy; Università Campus Bio-medico, Via Alvaro del Portillo, 21 - 00128 Roma, Italy

[pdugo@unime.it](mailto:pdugo@unime.it)

Keywords: ionic liquid stationary phase, terpene compounds, peppermint essential oil

Recently a new class of compounds has received increasing attention for their exploitation as GC stationary phases, namely room temperature ionic liquids (ILs), that have recently demonstrated great suitability in the analysis of essential oils [1] being capable of unravelling coelutions that could arise on the most widely used columns in such an application. Essential oils represent a valuable product, widely employed in several fields such as the flavor, fragrance and food industries. The volatile fraction of essential oils is usually rather complex, since it can be composed of hundreds of compounds, mainly monoterpenes, sesquiterpenes and their oxygenated counterparts, which are difficultly separated in a single GC run. For such a reason, it is common, both for qualitative and quantitative purposes, to analyze essential oils on both apolar and polar columns [2].

In this work a novel IL GC column, recently commercially available, and possessing an overall polarity close to that of polyethylene glycol phase [3], has been evaluated towards the separation of terpene compounds, in terms of peak symmetry and resolution of target volatile analytes and, furthermore, it has been applied to the analysis of a peppermint essential oil; the results have been compared to those attained on the polyethylene glycol phase demonstrating the great suitability of this novel IL GC stationary phase in this kind of application, since the overall separation performance and, thus, selectivity of the IL column was better than that obtained on the most widely used polar column.

Acknowledgements: The Project was funded by the "Italian Ministry for the University and Research (MIUR)" within the National Operative Project "Hi-Life Health Products from the industry of foods". Project ID: PON01\_01499.

1. C. Ragonese et al. *Anal. Chem.* (2011) 83: 7947-7954.

2. G. Dugo et al. (2010) *Citrus oil: composition, advanced analytical techniques, contaminants and biological activity*, 1st ed.;

3. C. Ragonese et al. *J. Chromatogr. A* <http://dx.doi.org/10.1016/j.chroma.2012.04.069>

**P 190. Biological investigations by rapid TLC methods on essential oils from *Hua gabonii* grown in Gabon**

Agnaniet H<sup>1</sup>, Makani Th<sup>1</sup>; Bikanga R<sup>1</sup>; Widmer V<sup>2</sup>; Ankle A<sup>2</sup>; Eike R<sup>2</sup>; Mbot E<sup>1</sup>; Bongui JB<sup>1</sup>; Anguille JJ<sup>1</sup>; Lebibi J<sup>1</sup>; Morère A<sup>3</sup>; Menut C<sup>3</sup>

<sup>1</sup>Université des Sciences et Techniques de Masuku; Faculté des Sciences, Laboratoire de Substances Naturelles et de Synthèse Organométalliques (LASUNASOM) ; BP 941 Franceville, GABON;;

<sup>2</sup>CAMAG Laboratory; Sonnenmattstrasse 11, CH- Muttenz Switzerland;

<sup>3</sup>Equipe Glyco et nanovecteurs pour le ciblage thérapeutique, IBMM, Faculté de Pharmacie, 15 Avenue Charles Flahault, BP 14491, 34093 Montpellier France

ahuguette2001@yahoo.fr

Keywords: *Hua gabonii*, essential oils, sulfide compounds, HPTLC, DPPH method, acetylcholinesterase essay

The plant *Hua gabonii* Pierre ex De Willd. (Huaceae), also called “garlic tree” for the characteristic garlic aroma of different parts of the plant, is widespread in dense tropical and humid forest, especially in Gabon, Cameroon and Congo [1]. In Cameroon, the seeds are used with other spices for the flavoring of various local dishes as alternative to common garlic (*Allium sativum*) [2]; in Gabon, the bark is used in alimentary for the same properties; seeds and bark are also traditionally utilized as infusions against colds or in fumigation against rheumatism and headaches [3].

Leaves, barks and roots collected on a specimen of *Hua gabonii* grown in Franceville (Gabon) were submitted to hydrodistillation to give essential oils with 0.21%, 0.37% and 0.39 % yields respectively. All samples presented characteristic odors reminiscent of sulfide compounds.

These essential oils were screened by rapid TLC according to two different methods: a bioautographic enzyme assay for the screening of acetylcholinesterase inhibition [4], involved in Alzheimer’s disease, as well as a DPPH assay, for their antiradical properties evaluation.

The three samples showed some faint acetylcholinesterase (Rf = 0.52, 0.73) and DPPH (Rf = 0.71) active zones.

[1] EJ Adjanohoun, AMR Ahyi, LA Assi et al. (1988) Contribution aux études ethnobotaniques et floristiques en République Populaire du Congo. ACCT, Paris, pp. 211.

[2] L Jirovetz, G Buchbauer, MB Ngassoum, M Geissler (2002) Eur. Food Res. Technol. 214 : 212-215.

[3] A Raponda Walker, R Sillans (1961) Les plantes utiles du Gabon, (Eds.), Paul Lechevalier, Paris, p 203

[4] A Marston, J Kissling, K Hostettmann (2002) Phytochem. Anal. 13: 51-54.

**P 191. Use of computer simulation in steam distillation of essential oils – novel design strategies for enhanced isolation of volatile plant extractives**

Phinn Masango

*Vorgartenstrasse 129-143/1/4, A1020, Wien, Austria*

*mutape2002@yahoo.co.uk*

Keywords: bubble point, essential oils, simulation, stripping.

Computer simulation is widely used in all the different areas the chemical industry such as refining and petrochemicals, pharmaceuticals, agrochemicals, and specialty chemicals. However its application in small to medium scale industries such as steam distillation of essential oils is limited. This limitation of the use of computer simulators or absence of it in the distillation the essential oils is mainly due to the fact that that this industry often operates as a cottage industry with expertise transferred as an art as well as limited resources such computing resources and expertise in advanced process design technology. The advent of the computer (simulation) applications in the design and analysis of chemical industry can also be applied to essential oils production. This application will result in optimized, cleaner and more economic operations. The work in this paper seeks to outline a methodical approach and tools that can be used based on Aspen Hysys computer simulation platform. The results are an improved process that results in a seventh of the energy use and a corresponding reduction in equipment size. This equipment includes condensers and all other equipment downstream to the packed bed or still to include piping. The reduction in hydraulic loading of the process units means a reduction in capital cost as well as operational costs. Though the reduction is not linear for all process parameters, the utilities consumption are reduced by about seven times that of the current process units. One of the key findings from this work is steam distillation units as currently operated operate at the lower end of the process window. That uses more energy and more water in the distillate. The increased aqueous fraction in the distillate most likely increases the solubility of the polar components. This changes the quality of the final oil product. If these polar components make up the top notes constituents, then they are lost in the aqueous fraction. The work improves the process by eliminating such losses and improving the utilities consumption and capital outlay.



**P 192. On the study of some essential oils of Lauraceae family from Vietnam**Dai DN<sup>1</sup>, Thai TH<sup>1</sup>; Thang TD<sup>2</sup>; Son LC<sup>1</sup>; Huyen DD<sup>1</sup>; Ogunwande IA<sup>3</sup><sup>1</sup>*Institute of Ecology and Biological Resources, Vietnam Academy of Science and Technology, 18-Hoang Quoc Viet, Cau Giay, Ha Noi, Vietnam;*<sup>2</sup>*Department of Food Technology, Vinh University, 182-Le Duan, Vinh City, Nghe An Province, Vietnam;*<sup>3</sup>*Natural Products Research Unit, Department of Chemistry, Faculty of Science, Lagos State University, Badagry Expressway Ojo, P. M. B. 0001, Lasu Post Office, Ojo, Lagos, Nigeria*

isiaka.ogunwande@lasu.edu.ng

Keywords: Lauraceae, Essential oils, Terpenoids

We have developed a systematic approach at the analysis of volatile compounds of Vietnamese [1]. The essential oils obtained by hydrodistillation from seven plants of Lauraceae family were analysed by GC and GC/MS. The leaf oil of *Phoebe angustifolia* Meisn., had spathulenol (17.0%), 1,2-Benzenedicarboxylic acid (13.0%) and sabinene (6.4%) in abundance; while the fruit consist mainly of 1,2-Benzenedicarboxylic acid (15.0%) and spathulenol/ cis- $\alpha$ -bisabolene (ca. 5.1%). The main constituents of *Machilus velutina* Champ. ex Benth., were (*E*)- $\beta$ -ocimene (9.5%), (*Z*)- $\beta$ -ocimene (8.2%), bicycloelemene (7.1%), germacrene D (6.8%) and *allo*-ocimene (6.4%). However, (*E*)- $\beta$ -ocimene (85.6%) was the most singly abundant component of *Neolitsea polycarpa* Liou.

The main compounds of *Cinnamomum sericans* Hance, were the sesquiterpenes spathulenol (14.5%), caryophyllene oxide (9.3%),  $\alpha$ -pinene (9.3%), sabinene (8.0%) and  $\beta$ -caryophyllene (7.1%). The monoterpenes- *p*-cymene (15.6%), limonene (13.9%) and  $\alpha$ -phellandrene (9.2%) - were the dominant class of compounds in *Cinnamomum durifolium* Kosterm. *Cinnamomum magnificum* Kosterm., was devoid of monoterpenes, while the major sesquiterpens were bicyclogermacrene (33.9%),  $\beta$ -caryophyllene (25.5%), bicycloelemene (7.2%) and caryophyllene oxide (7.5%).  $\beta$ -Caryophyllene (35.9%), caryophyllene oxide (12.6%) and spathulenol (5.2%) were the dominant compounds of Reinw. ex Blume. The low content of (*E*)-cinnamaldehyde in the *Cinnamomum* species is typical for majority of other species already reported from Vietnam [2-3].

1. DN Dai et al. (2012) Nat. Prodt. Comm. 7: 231-234

2. 2. XD Nguyen et al. (1995) J. Essent. Oil Res. 7: 53-55

3. 3. XD Nguyen et al. (1997) J. Essent. Oil Res. 9: 57-65

**P 193. Chemotaxonomy of essential oils of some *Cassia* species from Nigeria**

Ogunwande IA<sup>1</sup>, Guido F<sup>2</sup>; Avoseh NO<sup>1</sup>; Banwo ID<sup>1</sup>; Hassan AS<sup>1</sup>; Tihamiyu AZ<sup>1</sup>; Ogunsanwo AA<sup>1</sup>

<sup>1</sup>Natural Products Research Unit, Department of Chemistry, Faculty of Science, Lagos State University, Badagry Expressway Ojo, P. M. B. 0001, Lasu Post Office, Ojo, Lagos, Nigeria;

<sup>2</sup>Dipartimento di Scienze Farmaceutiche, sede Chimica Bioorganica e Biofarmacia, Università di Pisa, Via Bonanno 33, 56126 Pisa, Italy

isiaka.ogunwande@lasu.edu.ng

Keywords: Fabaceae, Essential oils; Terpenoids; Chemical markers

In continuation of our study on the chemical composition of the volatile oils obtained from the genus *Cassia* growing in Nigeria [1], we report herein the compounds identified from four *Cassia* species (Fabaceae family). The oils were by obtained hydrodistillation of the air-dried leaves samples and then analyzed by GC and GC/MS. The main compounds of *Cassia lutea* L. were hexahydrofarnesylacetone (20.0-22.5%), (*E*)-geranyl acetone (11.5-11.8%), farnesylacetone (4.8-5.8%), (*E*)-2-decenal (4.3-4.5%). There are significant amounts of (*Z*)-nerolidol (2.6-2.9%), (*E*)- $\alpha$ -ionone (2.1 and 2.5%) and 6-methyl-5-hepten-2-one (2.4-2.8%). The oil of *Cassia ferruginea* (Schrad.) DC consisted mainly of farnesylacetone (4.3%), 1-octen-3-ol (5.5%), pentadecanal (6.1%), phytol (5.8%), (*E*)- $\alpha$ -ionone (9.5%) and (*E*)-geranyl acetone (13.7%). We have identified (*E*)-geranyl acetone (5.8%), 1-octen-3-ol (5.8%), linalool (7.8%), *iso*-italicene (15.4%) and (*E*)- $\beta$ -damascenone (11.0%) as the abundant constituents in *Cassia siemens* L. On the other hand, the oxygenated compounds dominated in the oil of *Cassia occidentalis* L. These are (*E*)-geranyl acetone (8.0%), hexahydrofarnesylacetone (24.0%) and (*E*)-phytol acetate (40.7%). All the studied oil samples contained (*E*)- $\beta$ -ionone (1.1-3.7%), (*E*)-geranyl acetone (5.8-13.7%), (*E*)- $\beta$ -damascenone (0.2-11.0%), hexahydrofarnesylacetone (1.2-22.5%) and farnesylacetone (2.8-5.8%). In addition, the oil samples except *C. siemens* had phytol (0.4-5.8%). These compounds may serves as chemical markers of Nigerian grown *Cassia* species.

1. IA Ogunwande et al (2010) Rec. Nat. Prod. 4: 211-217

**P 194. Yield and composition of the essential oil of *Origanum vulgare* in function of the different types of fertilizer, systems and seasons of planting**

Luz JMQ<sup>1</sup>, Soares JS<sup>1</sup>; Silva SM<sup>1</sup>; Oliveira RC<sup>1</sup>; Facanali R<sup>2</sup>; Marques MOM<sup>2</sup>

<sup>1</sup>Federal University of Uberlândia;

<sup>2</sup>Agronomic Institute of Campinas

*jmagno@umuarara.ufu.br*

Keywords: *Origanum vulgare* L., aromatic plant, fertilization, greenhouse, seasons

*Origanum vulgare* L. better known as marjoram or oregano is a herbaceous plant belonging to the family Lamiaceae. It is native to the Euro-Siberian and Irano-Siberian, which determines long-range chemical and morphological diversity. The content and composition of essential oil of herbs like oregano, depends on different factors such as climate, geographical origin, harvest season, fertilizer and mineral nutrition can significantly affect the production and oil quality. The objective of the work was to analyze the production of biomass, content, yield and composition of the essential oil of oregano under different systems and seasons of planting. The experimental design used in both experiments was a randomized block design in a factorial 2 x 2 x 2 factorial with five replications, two tillage systems (greenhouse and field), two types of fertilizers (mineral and organic) and two growing seasons (autumn and spring). The field plots and greenhouse were three rows with three meters in length and the useful plot of each replicate consisting of seven plants in the central line. The row spacings were 40 cm and 35 cm between plants. The essential oil was extracted by hydrodistillation in a modified Clevenger apparatus. The chemical composition was analyzed by GC/MS. The essential oil presented the same compounds for all treatments; however, the relative proportion of some chemical constituents was altered according to the treatment. Carvacrol, terpinene, ortho-cymene were the major constituents. Both the seasons and the plantation in the field had favored greater essential oil yield and production of carvacrol. The biomass production was better at the hot and humid season. At the season spring/summer, the relative percentages of carvacrol and ortocimeno had been lesser in the field. But, for terpinene, the percentage was bigger in the field using chemical or organic fertilizer. At the season autumn winter, the relative percentage of the orto-cimene component was bigger in the field, but for carvacrol and gamma-terpinene the percentages had been bigger in the greenhouse.

Acknowledgements: FAPEMIG and CNPq for the support to this work.

**P 195. Tomato aroma volatiles measured by PTR-MS following artificial chewing**

Brian Farneti<sup>1</sup>, Simona Cristescu<sup>2</sup>; Guglielmo Costa<sup>3</sup>; Frans J.M. Harren<sup>2</sup>; Ernst J. Woltering<sup>4</sup>

<sup>1</sup>Wageningen UR and Bologna University;

<sup>2</sup>Radboud University;

<sup>3</sup>Bologna University;

<sup>4</sup>Wageningen UR

brian.farneti@gmail.com

Keywords: Tomato; PTR-MS; SPME-GC-MS; flavor

The availability of rapid and accurate methods to assess food flavor is of utmost importance to support quality control measures.

The perception of food flavor and aroma is a complicated process since flavor compounds are released from the food matrix during eating and then transported to the receptors in the mouth and nose. *In vivo* measurements of volatile patterns are therefore most desirable since they reflect the volatile profile reaching the olfactory receptors and therefore they relate better with sensory perception. However, due to the high variability generally observed in oral physiology parameters, flavor release kinetics and sensitivity to flavor stimuli, *in vivo* measurements are not suitable as standard and repeatable method.

The aim of this work was to develop a fast and reliable system to study the volatile aroma profile as close as possible to human perception, using tomatoes as model fruit.

The volatiles emitted by fruit of nine tomato varieties (at ripe stage) were analyzed by using two solvent-free headspace methods: solid-phase micro extraction/gas chromatography-mass spectrometry (SPME/GC-MS) and proton transfer reaction / mass spectrometry (PTR-MS) coupled with an artificial chewing device. In addition, volatiles were analysed from human mouth following natural chewing.

Parallel with the chemical analysis, a quantitative descriptive analysis (QDA) by trained panellists was carried out to qualitatively characterize the flavor of the nine tomato varieties.

Volatile patterns measured after artificial chewing showed good correlation with patterns produced during chewing in the human mouth. Multivariate statistical analysis (principle component, cluster and partial least square regression) of the PTR-MS results allow an unambiguous fingerprinting separation between different tomato varieties. A good correlation with scored sensory parameters, particularly for "tomato fragrance" and "tomato flavour" was shown.

In this study, it is shown that PTR-MS coupled with an artificial chewing device is a suitable method to monitor at high sensitivity the emission of volatiles determining the tomato aroma profile.

**P 196. Conformational analysis of the predominant antistaphylococcal sesquiterpene lactones from *Inula helenium* L. root essential oil**

Polina Blagojević, Niko Radulović

*University of Niš, Faculty of Science and Mathematics, Serbia*

*blagojevicpolina@gmail.com*

Keywords: alantolactone; isoalantolactone; diplophyllin; *Inula helenium* L. (Asteraceae); conformational analysis

It was recently shown that eudesmane-type sesquiterpene lactones: alantolactone (**1**), isoalantolactone (**2**) and diplophyllin (**3**), the dominant constituents of *Inula helenium* L. (Asteraceae) root essential oil, possess a very potent antistaphylococcal activity, with obvious membrane-damaging effects [1]. Although **1-3** are isomers without any essential structural difference, the activity of diplophyllin seems to be significantly higher than that of the other two isomers [1]. Alongside with the presence of some specific functional groups, three-dimensional structure (conformation) can play a significant role when speaking of biological/pharmacological activity of a certain compound [2]. Having this in mind, we decided to compute (MM+ and MMFF94 molecular mechanics and AM1 and PM3 semi-empirical methods; semi/empirical computations were performed with Polak-Ribiere (conjugate gradient) minimization method and with an energy convergence criterion of 0.01 kJ/mol) energetically favorable conformations of the dominant biologically active *I. helenium* oil constituents, and to seek for possible differences between them. Energetically favorable conformations of **1-3** were additionally described in the terms of selected 2D and 3D molecular descriptors (topological indices, conformational, molecular surface and partitioning descriptors). According to the obtained results, alantolactone and isoalantolactone mainly adopt “U”-shaped conformations (>99%, calculated at 298 K; “closed” geometries), with the carbonyl moiety being rather sterically hindered from one side. Contrary to that, in diplophyllin, the cyclohexene and lactone rings are mutually oriented in such a way as to give the overall “S”-shape to the molecule (“open” geometry). These conformational differences (that may influence optimal placement of **1-3** in the binding region of the target biomolecule(s)) could be, at least partially, responsible for the higher observed antistaphylococcal activity of diplophyllin in comparison to the other two isomeric lactones.

Acknowledgments. This work was funded by the Ministry of Education and Science of Serbia (Project 172061). PB was supported by UNESCO-L'OREAL National Fellowships Programme for Young Women in Life Sciences.

[1] Z Stojanović-Radić et al. (2011) Eur. J. Clin. Microbiol. Infect. Dis. 31:1015–1025.

[2] N Radulović et al. (2011) J. Chem. Crystallogr. 41: 545–551.

**P 197. What is nanjamonjagoke (*Takakia* sp.)? Taxonomical position of *T. lepidozoides* and *T. ceratophylla***

Nii K<sup>1</sup>, Higuchi M<sup>2</sup>, Asakawa Y<sup>1</sup>

<sup>1</sup>Faculty of Pharmaceutical Sciences, Tokushima Bunri University, Yamashiro-cho 180, 770-8514 Tokushima, Japan,

<sup>2</sup>National Museum of Nature and Science, Department of Botany, Ueno Park 7-20, Taito-ku, 110-8718 Tokyo, Japan

niiik@ph.bunri-u.ac.jp

Keywords: liverwort; *Takakia*; sesquiterpene lactones; coumarin; chemosystematics

*Takakia lepidozoides* was first discovered in Japan. Hattori and Inoue [1] classified it into the Marchantiophyta (liverwort) although Bryologists confused to where this plant should be classified; to algae, bryophyte or fern. In 1979 we concluded that *T. lepidozoides* could be classified into liverwort because of the presence of sesquiterpene lactone, diplophyllolide [2]. However, since the forms of sporophytes of *T. ceratophylla* later discovered was similar to the moss, *Andereaobryum macrosporum*, *Takakia* species was reclassified into the moss [3]. To reconfirm the above new discussion, a large amount of *Takakia lepidozoides* (1.3g) was analyzed by TLC, GC/MS and NMR to identify the presence of diplophyllolides, coumarin (major), the strong scent of this species, dihydrocoumarin, 1,4-benzoquinone, hydroquinone, dihydrobenzofurane, eudesm-4,11-diene and alpha-asarone (**9**). These compounds, especially, sesquiterpene eudesmanes, are one of the most important chemical markers of liverworts. In conclusion, *T. lepidozoides* is chemically more close to the Marchantiophyta than Bryophyta. *T. ceratophylla* shows the chemical character both of liverwort and moss because it produces anastreptene (**10**), a significant chemical indicator of many liverworts and hopane-type triterpenoids of mosses.

[1] Hattori S, Inoue H. (1958) J. Hattori Bot. Lab. 58, 19, 133-137

[2] Asakawa Y., Hattori S, Mizutani M, Tokunaga N, Takemoto T. (1979) J. Hattori Bot. Lab. 46, 77-90


[3] Smith DK, Daviso PG. (1993) J. Hattori Bot. Lab. 73, 263-271.

**P 198. Biotransformation of *p*-cymene by *Neurospora crassa* and *Aspergillus niger***Noma Y<sup>1</sup>, Asakawa Y<sup>2</sup><sup>1</sup>Faculty of Human Life Sciences;<sup>2</sup>Faculty of Pharmaceutical Sciences, Tokushima Bunri University.

ynoma@tokushima.bunri-u.ac.jp

Keywords: biotransformation; *p*-cymene; *Neurospora crassa*; *Aspergillus niger*

In the continuing studies on the microbial transformation of terpenoids and aromatic compounds[1,2], we report the biotransformation of *p*-cymene (**1**), a naturally occurring aromatic organic compound and a constituent of the essential oil of cumin and thyme, by 2 kinds of *Neurospora crassa* strain F40 and F42 and *Aspergillus niger* TBUYN-2. *N. crassa* was cultivated in the czapek-pepton medium at 30°C for 3 days. After full growth of microorganisms **1** was added to the cultured medium of microorganisms everyday at ca. 20mg in the 200ml cultured medium to 29 days. Totally 12.48g (2.08%) of substrate was added to the 6 flask. The ether extract of medium is 373.5mg (85.5%), ethyl acetate extract of medium 20.1mg (4.6%), ether extract of mycelium 21.4mg (4.9%) and ethyl acetate of mycelium 21.8 (5.0%). Each cultured broth was filtrated, and the resulting broth and the mycelium were extracted with ether, and then with ethyl acetate. These extracts were chromatographed on silica gel and each fraction was applied to GC-MS. Compounds shown in Fig.1 are identified by GC-MS with comparison of authentic compounds. The identification of each metabolite and the metabolic pathway of **1** will be discussed.

Fig.1. The biotransformation of **1** by *N. crassa* and *A. niger* TBUYN-2.

[1] Noma Y and Asakawa Y (2009) Handbook of Essential Oils. K. H. C. Baser and G. Buchbauer (Eds), CRC Press. Boca Raton, pp. 583-853.

[2] Noma Y and Asakawa Y (2010) Comprehensive Natural Products II, Chemistry and Biology, Mander, L.,Lui, H-W. (Eds), Elsevier: Oxford, Vol. 3, pp. 669-801.

**P 199. Chemical composition and antibacterial activity of the essential oils isolated from leaves and fruits of *Peucedanum austriacum* (Jacq.) W. D. J. Koch**

Jovanović O<sup>1</sup>, Zlatković B<sup>2</sup>, Simonović S<sup>1</sup>, Đorđević A<sup>1</sup>, Palić I<sup>1</sup>, Stojanović G<sup>1</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science and Mathematics, University of Niš, Višegradska 33, 18000 Niš, Serbia

<sup>2</sup>Department of Biology and Ecology, Faculty of Science and Mathematics, University of Niš, Višegradska 33, 18000 Niš, Serbia

gocast@pmf.ni.ac.rs

Keywords: *Peucedanum austriacum* (Jacq.) W. D. J. Koch; Umbelliferae, essential oil composition;  $\beta$ -phellandrene; caryophyllene oxide; antibacterial activity

Chemical composition and antibacterial activity of the essential oils isolated from fruits and leaves of *Peucedanum austriacum* (Jacq.) W. D. J. Koch were determined in order to define their medical potential. An analyses of leaves and fruits essential oils by GC and GC/MS, resulted in identification of 141 different components, representing 93.6% and 96.1% of the total oils. The most abundant class of compounds in fruits essential oil were monoterpenoids (64.0%), while in the leaves essential oil were sesquiterpenoids (82.0%). The major contributors of fruits essential oil were  $\beta$ -phellandrene (45.2%) and  $\alpha$ -pinene (10.1%). Major constituents of leaves essential oil were caryophyllene oxide (23.1%), germacrene D (12.2%) and (*E*)-caryophyllene (10.2%). Minimal inhibitory (MIC) and minimal bactericidal concentrations (MBC) were determined by performing broth microdilution assay against two Gram-positive bacteria (*Bacillus subtilis* ATCC 6633 and *Staphylococcus aureus* ATCC 6538) and three Gram-negative bacteria (*Escherichia coli* ATCC 8739, *Pseudomonas aeruginosa* ATCC 9027 and *Salmonella typhimurium* ATCC 14028). In the antibacterial activity assay, both samples showed moderate activity against *Bacillus subtilis* and *Staphylococcus aureus* with inhibitory effect at 0.625 and 1.25 mg mL<sup>-1</sup>, weak activity against *Pseudomonas aeruginosa* with inhibitory effect at 5.00 mg mL<sup>-1</sup> and no effect on the growth of bacteria *Escherichia coli* and *Salmonella typhimurium*. The results showed that essential oils composition of *P. austriacum* leaves and fruits differed considerably. Tested oils were proven to be selective antibacterial agents.

Acknowledgements: The authors are grateful to the Serbian Ministry of Education and Science Ministry for financial support (Project 172047).



**P 200. The composition of *Hypericum umbellatum* A. Kern. essential oil from Serbia**Đorđević A<sup>1</sup>, Lazarević J<sup>2</sup>, Stojanović G<sup>1</sup>, Palić I<sup>1</sup> and Petrović G<sup>1</sup><sup>1</sup>Department of Chemistry, Faculty of Science and Mathematics, University of Niš, Višegradska 33, 18000 Niš, Serbia<sup>2</sup>Department of Pharmacy, Faculty of Medical Sciences, University of Niš, Bul. dr Zorana Đinđića 81, 18000 Niš, Serbia

gocast@pmf.ni.ac.rs

Keywords: *Hypericum umbellatum*, Hypericaceae, essential oil composition

*Hypericum* L. (Guttiferae/Clusiaceae/Hypericaceae) is a genus represented by more than 400 species, widespread in warm-temperate areas throughout the world, as well as on the Balkan Peninsula [1]. Plants of the genus *Hypericum* have traditionally been used as highly esteemed medicinal plants particularly *H. perforatum* (St. John's wort) [2]. The aim of this study was to perform a detailed analysis of the *H. umbellatum* essential oil originating from Serbia, and establish chemotaxonomic correlation within the members of the section *Drosocarpium* Spach (*H. umbellatum* is a member of this section). The essential oil of fresh aerial parts of *H. umbellatum* obtained by hydrodistillation was analyzed by GC and GC/MS. One hundred and twenty-six identified compounds accounted for 93.9% of the total oil. The main oil components were: germacrene D (6.1%), (*E*)-nerolidol (4.4%), n-nonane (4.0%), (*E*)-caryophyllene (3.0%) and caryophyllene oxide (3.0%). Dominant class of compounds - terpenoids (75.9%), was unevenly distributed between mono- and sesquiterpenoids (14.9% and 59.2%, respectively). Monoterpenoids were mostly made of hydrocarbons (12.3 %), opposed to sesquiterpenoids made of relatively comparable amounts of hydrocarbons (27.3%) and oxygenated derivatives (31.9%). Non-terpenoid compounds (17.0%) consisted mainly of n-alkanes (9.6%). To the best of our knowledge the essential oil of *H. umbellatum* has not been studied previously. The results of this study showed that chemical composition of *H. umbellatum* oil is in agreement with previous reports on the species belonging to the section *Drosocarpium* Spach.

Acknowledgments: This work was funded by the Ministry of Education and Science of Serbia (Project No. 172047).

1. NKB Robson, A Strid (1986) *In*: Strid, A. (Ed.) Mountain Flora of Greece, Cambridge University Press, Cambridge, vol. 1, pp. 594-608.
2. K Yazaki, T Okada (1994) *In*: Bajaj, Y.P.S. (Ed.) Biotechnology in agriculture and forestry, Springer-Verlag, Berlin, vol. 26, pp. 167-178.

**P 201. Enantioselective GC-MS analysis of aroma components of essential oils and hydrosols extracted from rosemary (*Rosmarinus officinalis*)**

Tomi K<sup>1</sup>, Kitao M<sup>1</sup>; Hayashi T<sup>1</sup>; Nakagawa M<sup>1</sup>; Utsunomiya N<sup>1</sup>; Murakami H<sup>2</sup>; Matsumura Y<sup>2</sup>

<sup>1</sup>Faculty of Agriculture, Kinki University;

<sup>2</sup>Graduate School of Agriculture, Kyoto University

tomi@nara.kindai.ac.jp

Keywords: optical isomer; enantioselective GC-MS; essential oil; hydrosol; rosemary; principal component analysis

When essential oils are extracted by hydrodistillation from plants, hydrosols in the form of aqueous solution are obtained as byproducts. While essential oils are often used for food production as well as aromatherapy, utilization of hydrosols is under way because of the absence of the data about their chemical composition. In this study, essential oils and hydrosols were separated from rosemary harvested in several seasons, and chemical composition of aroma components in the two fractions was analyzed by GC-MS. Optical isomer composition of some aroma components was also analyzed.

Leaves of rosemary were collected every two months from February 2010 to February 2011. Leaves were soon subjected to hydrodistillation after harvesting. Essential oils and hydrosols were analyzed using GC-MS. For the analysis of the aroma component compositions, DB-1MS column was used. Optical isomer composition was determined using MEGA-DEX DET-beta column, which can separate optical isomers of aroma components. Results were analyzed by using principal component analysis.

The main components of the rosemary essential oil were eucalyptol and alpha-pinene. On the other hand, camphor, borneol and eucalyptol were predominant in the hydrosols. Classification of aroma components based on chemical groups revealed that essential oils contained the high amount of monoterpene hydrocarbons but hydrosols not. There is no difference in the ratio of enantiomer for optically active aroma components between essential oils and hydrosols. Principal component analysis indicated some seasonal variations of aroma component compositions in both essential oils and hydrosols.

The main aroma components of rosemary essential oils and hydrosols were different with each other, suggesting the different biological functions of the two fractions. Similar results were obtained for the optical isomer composition of some aroma components between essential oils and hydrosols.

**P 202. Synergistic antimicrobial activities of the combination essential oils of *Cymbopogon citratus* and *Alpinia galanga***

Sarin Tadtong<sup>1</sup>, Rith Watthanachaiyingcharoen<sup>1</sup>; Narisa Kamkaen<sup>2</sup>

<sup>1</sup>Faculty of Pharmacy, Srinakharinwirot University, Nakhon-Nayok, 26120 THAILAND;

<sup>2</sup>Faculty of Science and Technology, Suan Dusit Rajabhat University, Dusit, Bangkok 10300 THAILAND

[nkamkaen@gmail.com](mailto:nkamkaen@gmail.com)

Keywords: Synergistic antimicrobial activity; Essential oils; *Cymbopogon citratus*; *Alpinia galanga*

To evaluate antimicrobial activities of the single and combined essential oils of *Cymbopogon citratus* and *Alpinia galanga* against four pathogenic microorganisms: *Staphylococcus aureus* ATCC6538, *Pseudomonas aeruginosa* ATCC9027, *Streptococcus bovis* DMST18769 and *Candida albicans* ATCC10231.

Fresh leaves of *C. citratus* and rhizome of *A. galanga* were submitted to hydrodistillation in a Clevenger-type apparatus for 2 hours [1]. Qualitative analyses were conducted with Finnigan Trace GC ultra equipped with Finnigan DSQ Quadrupole detector [2, 3]. Minimum inhibitory concentration (MIC) of essential oils was determined by broth microdilution method [4, 5, 6].

The light yellow essential oils of *C. citratus* (lemongrass) and *A. galanga* (galanga) were obtained in yields of 0.24% and 0.11%, respectively, based on dried extracted material. Nineteen compounds were identified in lemongrass oil sample that included geraniol (16.4%), neral (10.2%) and myrcene (7.67%) as major components. Thirty-six compounds were identified in galanga oil. The major constituents were 1,8-cineole (68.18%), limonene (2.5%), terpinen-4-ol (2.23%),  $\alpha$ -terpineol (1.16%), and  $\alpha$ -pinene (1.1%). The MIC of lemongrass oil and galanga oil were observed against *S. aureus* (0.5% and 4% v/v), *P. aeruginosa* (40%v/v and >40%v/v), *S. bovis* (0.25%v/v and 0.5%v/v) and *C. albicans* (0.25%v/v and 0.5%v/v). The combination profiles of lemongrass oil with galanga oil (ratios 3:7, 5:5, and 7:3) were tested against four pathogenic microorganisms. Synergy was best noted for only one ratio (ratio 7:3).

Results of the present investigation provide evidence that the utilization of two essential oils combination could be assessed for synergistic antimicrobial activity in order to reduce their minimum effective dose.

Acknowledgements: National Research Council of Thailand (NRCT) is acknowledged for financial support.

1. BT Schaneberg, IA Khan. (2002) J. Agr. Food Chem. 50: 1345-1349.
2. Z Cui. (2003) Lixueban. 38: 104-107.
3. VS Rana, M Verdeguer, MA Blazquez. (2010) J. Essen. Oil Res. 22: 521-524.
4. CK Hindumathy. (2011) World Acad. Sci. Engineering Technol. 74: 193-197.
5. J Oonmetta-areea et al. (2006) LWT-Food Sci. Technol. 39: 1214-1220.
6. S Tadtong et al. (2011) J. Health Res. 25(1): 35-37.

**P 203. Larvicidal activity of Apiaceae essential oils against *Anopheles atroparvus* Van Thiel (Diptera, Culicidae) a former european malaria vector**

Sousa RMOF<sup>1</sup>, Novo MT<sup>2</sup>; Fernandes-Ferreira M<sup>3</sup>

<sup>1</sup>Centre for the Research and Technology of Agro-Environmental and Biological Sciences, Quinta de Prados, Apartado 1013, 5001-801 Portugal; Department of Biology, University of Minho, Campus Gualtar, 4710-057 Braga, Portugal; <sup>2</sup>Unidade de Parasitologia e Microbiologia Médicas, Instituto de Higiene e Medicina Tropical, Universidade Nova de Lisboa, Rua da Junqueira 100, 1349-008 Lisboa, Portugal; <sup>3</sup>Centre for the Research and Technology of Agro-Environmental and Biological Sciences, Quinta de Prados, Apart 1013, 5001-801 Portugal; Department of Biology, Faculty of Science, University of Porto, 4169-007 Porto, Portugal; Mapprod Lda, Braga, Portugal.

rosemarie@bio.uminho.pt

Keywords: bio-pesticides; fennel; parsley; cumin; phenylpropanoids; mosquito

In the context of research for new bio-pesticides, extracts from plants have received special attention and volatile constituents of essential oils (EOs) are among the candidates with higher potential [1]. *Anopheles atroparvus* is a European mosquito with proved transmission capacity for malaria parasite, which, in a scenario of climate change, can raise concerns of malaria re-emergence in regions actually not affected by this disease [2]. EOs of three well-known Apiaceae species, wild fennel (*Foeniculum vulgare* Miller subsp. *vulgare* var. *vulgare*), parsley (*Petroselinum crispum* (Miller) A.W. Hill) and cumin (*Cuminum cyminum* L.), were included in this study as potential sources of bio-insecticides and characterized by gas chromatography (GC) and mass spectrometry (GC-MS). Mosquito larvicidal assays were conducted against third-instar larvae of *An. atroparvus*. Lethal concentrations causing 50 and 90% of mortality after 24 hours of exposure were determined by log-Probit regression. Typical constituents of the fennel fruits EO, estragole and *trans*-anethole, were also evaluated. The wild fennel fruits EO was mainly constituted by estragole (65%), fenchone (16%), phellandrenes (10%) and pinenes (5%). A high content of phenylpropanoids, myristicin (32%), apiole (16%) and 1-allyl-2,3,4,5-tetramethoxybenzene (9%), and a considerable amount of pinenes (30%) were found in parsley fruits EO. Cumin fruits EO was rich in cumin aldehyde (39%), *p*-mentha-1,4-dien-7-al (10%) and several monoterpenes hydrocarbons. Both parsley and cumin EOs were more effective against *An. atroparvus* third-instar larvae (LC50= 13 ppm/LC90= 28 ppm and LC50= 17 ppm/LC90= 43 ppm, respectively) than fennel fruits EO (LC50= 61 ppm/LC90= 107 ppm). The larvicidal activity of fennel EO was, in some extend, justified by its high estragole and low *trans*-anethole contents (65% and 2%, respectively), since the former was less active (estragole LC50= 56 ppm and *trans*-anethole LC50= 17 ppm).

Acknowledgments: This work was financially supported by the FCT (Fundação para a Ciência e Tecnologia) with a PhD grant SFRH/BD/66041/2009 and through the CITAB research center.

1. E Shaaya and A Rafaeli (2007) In, I Ishaaya, R Nauen and AR Horowitz (Eds.): Insecticides Design Using Advanced Technologies. Springer-Verlag Berlin Heidelberg, pp. 249-261.
2. TH Jetten and W Takken (1994). Wageningen Agricultural University Papers. 94.5.

**P 204. Antimycoplasmatic evaluation of essential oils, isolated compounds and semi-synthetic indoles**

Rebello RA<sup>1</sup>, Begnini IM<sup>1</sup>, Meneghetti EK<sup>1</sup>, Dognini J<sup>1</sup>, Tizziani T<sup>1</sup>, Cordova CMM<sup>1</sup>, Zeni ALB<sup>1</sup>, Sgrot F<sup>2</sup>

<sup>1</sup>Universidade Regional de Blumenau- FURB, 89012-900, Blumenau, SC, Brazil;

<sup>2</sup>Vegetal Brasil, 88101-280, Palhoça, SC, Brazil.

ricardorebello@furb.br


Keywords: *Mycoplasma mycoides*; essential oils; myristicin; sesquiterpenes; indoles; safrole.

Mycoplasmas are obligately parasitic bacteria with no cell wall causing diseases in a wide range of hosts, including humans [1]. In the search for new antimicrobial agents, phytochemicals represent a potential source for such class of compounds [2], therefore several natural products and two semi-synthetic indoles derived from safrole were assayed against *Mycoplasma mycoides* subsp. *capri*.

Essential oils (EOs) from *Drimys angustifolia* (leaf and branch), *Aloysia gratissima* (leaf) and *Melaleuca alternifolia* (leaf) were obtained by hydrodistillation under nitrogen atmosphere using a Clevenger type apparatus. The oils were characterized by means of gas chromatography (GC-FID) and gas chromatography coupled to mass spectrometry (GC-MS). The sesquiterpenes cyclocolorenone and bicyclogermacrene, and the arylpropanoid myristicin were isolated from leaf essential oil of *D. angustifolia* by column chromatography on silicagel. Pure hexane and the binary mixture (hexane:dichloromethane) were used as eluents. Drimenol was obtained from branch essential oil of *D. angustifolia* using crystallization in cold hexane. Semi-synthetic indoles (1) were obtained using piperonal, a safrole derivative, as starting material. EOs, isolated natural products and semi-synthetic indoles were assayed against *M. mycoides* using the broth microdilution method.

The tested samples showed activity against *M. mycoides* with Minimum Inhibitory Concentration (MIC) values of 50 mg mL<sup>-1</sup>. The most active sample was the indole ester 1, which showed MIC value of 17.5 mg mL<sup>-1</sup>.

EOs of several plant species as well as their isolated compounds showed weak activity against *M. mycoides*. The most promising compound, the semi-synthetic indole ester, may be suitable for chemical transformations in order to obtain more active molecular structures.



Acknowledgments: FURB, UNIVALI, FAPESC, Vegetal Brasil, INCT Catalysis.

1. AV Markov, IA Zakharov (2009) Russ. J. Genet. 45: 781-787.

2. M Saleem et al (2010) Nat. Prods. Rep. 27: 238-254.

**P 205. The essential oil of *Eryngium duriae* subsp. *juresianum* inhibits IL-1 $\beta$ -induced NF-KB AND MAPK activation and iNOS expression in human chondrocytes**

Rufino AT<sup>1</sup>, Cavaleiro C<sup>2</sup>; Judas F<sup>3</sup>; Salgueiro L<sup>2</sup>; Lopes MC<sup>1</sup>; Mendes AF<sup>1</sup>

<sup>1</sup>Center for Neuroscience and Cell Biology; Faculty of Pharmacy, University of Coimbra;

<sup>2</sup>Centre of Pharmaceutical Studies; Faculty of Pharmacy; University of Coimbra;

<sup>3</sup>Univ. Hosp. of Coimbra and Faculty of Medicine, Univ. of Coimbra

ana.t.rufino@gmail.com

Keywords: *Eryngium duriae*; Anti-inflammatory properties; Osteoarthritis; IL-1 $\beta$ ; MAPK

Osteoarthritis (OA) is characterized by cartilage and joint destruction. The nuclear factor-kappaB (NF-kB) and mitogen-activated protein kinase (MAPK) pathways mediate the increased expression of inflammatory factors and matrix metalloproteinases. Nitric Oxide (NO) produced by the inducible NO synthase (iNOS) is increased in OA chondrocytes and is involved in inhibition of matrix synthesis and induction of matrix degradation, leading to cartilage loss and joint destruction. IL-1 $\beta$  is a major pro-inflammatory and pro-catabolic cytokine that is increased in OA and activates the NF-kB and MAPK pathways and iNOS expression [1,2].

This work reports on the ability of the essential oil of *Eryngium duriae* subsp. *juresianum* (M. Lainz) (Ej) to decrease IL-1 $\beta$ -induced responses in human chondrocytes.

Knee cartilage samples were obtained from multi-organ donors (n=8) at the University Hospital of Coimbra. The essential oil of the aerial parts of Ej was diluted in DMSO. Final concentrations of 5 to 200 mg/mL were added to chondrocyte cultures 30 min before of IL-1 $\beta$  (10 ng/mL) for 5 min, 30 min or 18 h. The MTT reduction assay was used to rule out cytotoxic effects. NO levels were measured by the Griess reaction. Protein levels of iNOS, I $\kappa$ B- $\alpha$  and the phosphorylated forms of I $\kappa$ B- $\alpha$  and of the MAPK, p38, JNK and ERK1/2, were assessed by western blot. Results were considered statistically significant for P<0.05.

Ej significantly decreased IL-1-induced NO and iNOS levels in a concentration-dependent manner. The greatest inhibition of IL-1-induced NO ( $38.4 \pm 8.4\%$ , P<0.001) and iNOS protein ( $53.1 \pm 6.3\%$ , P<0.001) levels was achieved with a concentration of 25 mg/mL. 200 mg/mL were required for maximal inhibition of I $\kappa$ B- $\alpha$  ( $70.9 \pm 5.6\%$ , P<0.001), JNK ( $36.4 \pm 0.6\%$ , P<0.001), p38 ( $57.1 \pm 4.4\%$ , P<0.001) and ERK1/2 ( $24.4 \pm 3.3\%$ , P<0.001) phosphorylation.

These results suggest that by acting on various signalling pathways, Ej elicits the synergistic inhibition of iNOS expression and show the potential of Ej as a promising source of compounds with anti-inflammatory and anti-catabolic properties that may be useful as anti-OA drugs.

Acknowledgments: This work was supported by grants PTDC/EME-PME/103578/2008 and PTDC/EME-TME/113039/2009 from FCT.

1. Marcu, K.B., et al.(2010), Curr Drug Targets, 11(5): 599-613.

2. Berenbaum, F.(2008) Arthritis ResTher, 10(Suppl 2): p. S1

**P 206. Composition and antioxidant activity of essential oil from Portuguese myrtle (*Myrtus comunis* L.) through the vegetative cycle**

Paula Cristina Pereira<sup>1</sup>, Maria-João Cebola<sup>1</sup>; Maria Gabriela Bernardo-Gil<sup>2</sup>; Anabela Romano<sup>3</sup>

<sup>1</sup>Institute for Biotechnology and Bioengineering, Centre for Biological and Chemical Engineering, Instituto Superior Técnico, Av. Rovisco Pais, 1049-001 Lisboa, Portugal; Faculdade de Engenharias e Ciências Naturais, ULHT, Av. Campo Grande, 376, 1749-024 L;

<sup>2</sup>Institute for Biotechnology and Bioengineering, Centre for Biological and Chemical Engineering, Instituto Superior Técnico, Av. Rovisco Pais, 1049-001 Lisboa, Portugal;

<sup>3</sup>Institute for Biotechnology and Bioengineering, University of Algarve, Laboratory of Plant Biotechnology, Campus de Gambelas, 8005-139 Faro, Portugal

[m.joao.cebola@gmail.com](mailto:m.joao.cebola@gmail.com)

Keywords: *Myrtus communis* L.; essential oil; Antioxidant activity; Folin-Ciocalteu; TEAC; ORAC.

*Myrtus communis* L., commonly known as myrtle, possesses a set of qualities that make it very interesting for the pharmaceutical, nutraceutical and cosmetic industries. Previous studies have shown that some differences arise in the composition of the essential oil and in the plant extract due to the different stages of the vegetative cycle of the plant and also due to different geographical locations.

In this work a time perspective is also investigated. Thus, Portuguese Myrtle (*Myrtus communis* L.) is being studied over a period of three years covering the main stages of the vegetative cycle of the plant, and the chemical composition of the essential oil was determined. The oil, extracted from leaves and berries separately, was obtained by Clevenger distillation and analysed by GC and GC/MS. The results show that the major components are limonene+1,8-cineole, myrtenyl acetate,  $\alpha$ -pinene and linalool, and, over the three-year period, a decrease in the composition of the most volatile compounds was observed.

The aqueous phase obtained from the Clevenger distillation was extracted with diisopropylether obtaining what is hereby designated as the Liquid Phase extract. The extracts were subject to a determination of antioxidant activity using three different methods: the Folin-Ciocalteu, the TEAC and the ORAC methods. The results obtained for the first year of collection showed that the leaves at the flowering stage and the ripe fruit presented better antioxidant activity, thus the study of the remaining two years was focused on them. The antioxidant activity varies between 1404- 2052  $\mu\text{mol trolox/g}$  for ORAC, 4554-5457  $\mu\text{mol trolox/g}$  for TEAC and 843 – 1831  $\mu\text{mol GAE/g}$  for Folin-Ciocalteu, with a wider range of variability on the leaves than in the fruits. In addition, there seems to be a similar trend in the Folin-Ciocalteu and TEAC methods differing from the ORAC method.

The fruit antioxidant activity was similar in the three years; however, the leaves presented higher activity in the first year.

**P 207. Lab-on-a-chip: a technique for detection of antibacterial effect of essential oils on outer membrane proteins**

Horváth Gy<sup>1</sup>, Bencsik T<sup>1</sup>, Godányi R<sup>1</sup>, Felső P<sup>2</sup>, Felső P<sup>3</sup>, Lemberkovic É<sup>4</sup>, Böszörményi A<sup>4</sup>, Takátsy A<sup>5</sup>, Molnár P<sup>1</sup>, Kocsis B<sup>2</sup>

<sup>1</sup>Department of Pharmacognosy, Medical School, University of Pécs, 7624 Pécs, Rókus u. 2., Hungary

<sup>2</sup>Institute of Medical Microbiology and Immunology, Medical School, University of Pécs, 7624 Pécs, Szigeti út 12., Hungary

<sup>3</sup>Institute of Bioanalysis, Medical School, University of Pécs, 7624 Pécs, Szigeti út 12., Hungary

<sup>4</sup>Institute of Pharmacognosy, Faculty of Pharmacy, Semmelweis University, 1085 Budapest, Üllői út 26., Hungary

<sup>5</sup>Department of Biochemistry and Medical Chemistry, Medical School, University of Pécs, 7624 Pécs, Szigeti út 12., Hungary

gyorgyi.horvath@aok.pte.hu

Keywords: Essential oil, *Erwinia* sp., *Pseudomonas aeruginosa*, capillary electrophoresis, chip technology, outer membrane protein

The structural changes of outer membrane protein (OMP) composition may have an effect on the adhesive ability and pathogenic properties of bacteria. Previously, it has been demonstrated that some factors, e.g. antibiotics or iron restriction, can change the bacterial OMP composition [1, 2]. The aim of our study was to examine how the essential oils of thyme, cinnamon bark and clove modify the OMP composition of the plant pathogen *Erwinia* and the human pathogen *Pseudomonas aeruginosa* strains. In case of thyme oil, we wanted to analyse which components might play a significant role in the changes induced.

The essential oils were obtained from a Hungarian pharmacy. Chemical composition of the oils was controlled with GC and GC-MS. The minimum inhibitory concentration (MIC) values of the oils were determined by a modified tube dilution method. 0.5 and 2 x MIC concentrations of the oils were administered to the culture and incubated for 60 min. OMP preparation was performed according to ref [3]. Measurements were performed in the Protein 230 Plus LabChip Kits of the 2100 Bioanalyzer System of Agilent.

Thymol was the main component of the thyme oil (49.9%), eugenol (83.7%) of clove oil and trans-cinnamic aldehyde (73.2%) of cinnamon bark oil. Thymol as the main component of thyme oil caused major changes in the protein composition of *Erwinia*. Cinnamon and clove oil also influenced the OMP composition of *Pseudomonas* strains. After the treatment, the protein peak with molecular weight 42.7 kDa and 79.4 kDa disappeared in case of cinnamon oil and clove, respectively.

Quantitative changes in the protein profile may contribute to the explanation of antibacterial effect of thyme, cinnamon bark and clove essential oils on pathogenic *Erwinia* and *Pseudomonas* strains. The structure of proteins considerably reduced will be analysed with MALDI-TOF/MS.

1. I Kustos et al. (2005) Electrophoresis 26: 3789-3795.

2. L Babujee et al. (2007) J. Proteome Res. 6: 62-69.

3. I Kustos et al. (1998) Electrophoresis 19: 2324-2330.



**P 208. Transfer of volatiles from oregano or caraway essential oils into cow's milk**

Lejonklev J<sup>1</sup>, Larsen MK<sup>1</sup>; Mortensen G<sup>1</sup>; Petersen MA<sup>2</sup>; Weisbjerg MR<sup>3</sup>

<sup>1</sup>Aarhus University, Department of Food Science;

<sup>2</sup>University of Copenhagen, Department of Food Science;

<sup>3</sup>Aarhus University, Department of Animal Science

JohanS.Lejonklev@agrsci.dk

Keywords: aroma; caraway; milk; oregano

Essential oils have a number of applications as a feed additive in ruminant production of milk and meat (1). The objective of this experiment was to study the transfer of volatile compounds, mainly terpenes, from oregano and caraway essential oils into cow's milk. During feeding and rumination the animals lungs and intestines are both exposed to volatile compounds present in the feed (2). In order to study the differences between respiratory and gastrointestinal exposure Holstein cows equipped with a duodenum cannula were used in two setups with two animals receiving each treatment. In the first the animals were placed in a controlled environment to inhale vapours of the essential oils for 9 hours. In the second essential oils diluted in deodorised sesame oil was injected through the cannula over a period of 9 hours, with two different levels being tested. Milk was collected prior to and immediately after treatment, as well as the following morning. Commercially available essential oils from *Origanum vulgare* plants and *Carum carvi* seeds were used. The aroma profile of essential oils and milk was analysed using purge-and-trap coupled with GC/MS.

The results show that milk contains a number of terpenes naturally at very low levels. When the animal is exposed to essential oils several terpenes, present in the essential oils, increase or appear in the milk, suggesting that the aroma compounds are absorbed through the lungs as well as the intestine. It also indicates that the absorption, and subsequent transfer from blood into milk, is very fast. In addition to the terpenes two esters were identified that increased in the milk after exposure, despite the essential oils contained no or insignificant amounts of them. This indicates that these esters are more readily absorbed or synthesised within the animal following exposure to essential oils. Little or none of the increased amounts of aroma compounds, terpenes and esters, could be found in the milk one day after exposure.

1. Benchaar C et al. (2008) Anim. Feed Sci. and Technol. 145: 209-228

2. Shipe WF et al. (1962) J. Dairy Sci. 45: 472-476

**P 209. Flavor active compounds from minor fruits grown in Uruguay**

Martínez N<sup>1</sup>, Bellucci I<sup>1</sup>; Vignale B<sup>2</sup>; Cabrera D<sup>3</sup>; Carrau F<sup>4</sup>; Boido E<sup>4</sup>; Fariña L<sup>4</sup>; Dellacassa E<sup>5</sup>

<sup>1</sup>Cátedra de Farmacognosia y Productos Naturales, Facultad de Química-UdelaR. Montevideo, Uruguay.;

<sup>2</sup>EEFAS, Facultad de Agronomía-UdelaR. Salto, Uruguay.;

<sup>3</sup>INIA-Las Brujas;

<sup>4</sup>Sección Enología, Facultad de Química-UdelaR. Montevideo, Uruguay.;

<sup>5</sup>Cátedra de Farmacognosia y Productos Naturales, Facultad de Química-UdelaR. Montevideo, Uruguay; Sección Enología, Facultad de Química-UdelaR. Montevideo, Uruguay

edellac@fq.edu.uy

Keywords: flavor, Uruguayan native fruits, isoprenoids

The perception of fruits flavor and aroma is the result of a multitude of interactions between a large number of chemical compounds and sensory receptors. In the case of isoprenoids, their role in the aroma of exotic fruits is a determining factor in their acceptance by consumers. Aroma compounds are present in fruits in free volatile form but also as non-volatile precursors such as glycosides

Considering the complexity associated with the composition of the different aromatic fractions (free and glycosidated) present in exotic fruits, the development of chromatographic techniques make available not only to separate these components, but also to provide information about the main biological activity to be considered: their aroma description [1]. Gas chromatography coupled to specific detectors as mass spectrometry and olfactometry, are able to respond to this challenge [2].

In this work we present the results obtained in the developing of the required methodology to make possible to evaluate the aromatic potential represented by the glycosylated isoprenoids present in native Uruguayan fruits. The fruits used as a model of study were: *Acca sellowiana* (Berg.) Burret (Myrtaceae, "guayabo del país"), *Psidium cattleianum* Sab. (Myrtaceae, "arazá") and *Butia capitata* (Mart.) Becc. (Arecaceae, "butiá"). The results will be applied to develop new strategies in order to generate original elements of quality and authenticity for the fruits consumption and transformation.

Acknowledgments. This research was financially supported by the Agencia Nacional de Investigación e Innovación (ALI\_1\_2011\_1\_2545) and CSIC-UdelaR, Uruguay (Group Project 656).

[1] Versini et al. (2008) *In*: R Flaminio (Ed.) *Hyphenated Techniques in Grape and Wine Chemistry*. John Wiley & Sons, Chichester, pp. 173-217.

[2] Carrau et al. (2008) *Nat. Prod. Commun.* 3: 1-16.

## P 210. Mathematical Modeling of Supercritical CO<sub>2</sub> extraction of volatile oils from pennyroyal (*Mentha pulegium*)

Palavra AF<sup>1</sup>, Fareleira JMNA<sup>1</sup>, Nobre BP<sup>1</sup>, Cristino AF<sup>1</sup>, Matos P<sup>1</sup>, Coelho JAP<sup>2</sup>

<sup>1</sup>Centro de Química Estrutural, Instituto Superior Técnico, Av. Rovisco Pais, 1, 1049-001 Lisboa, Portugal; <sup>2</sup>Chemical Engineering and Biotechnology Research Center, Instituto Superior de Engenharia de Lisboa, Rua Conselheiro Emídio Navarro, 1, 1959-007 Lisboa, Portugal

jcoelho@deq.isel.ipl.pt

Keywords: Supercritical extraction, volatile oils, aromatic plants, mass transfer models.

The modelling of the experimental data of supercritical extraction of volatile oils from aromatic plants is very important, since it can be used as a tool for the design, improvement and scale up of these processes from laboratory to pilot and industrial scales. A discussion of the modelling of our data on supercritical fluid extraction of volatile oil from pennyroyal was carried out. The most successful models in this scientific area describe the experimental process by using differential mass balances for the fluid and solid phases, as shown with the modelling studies carried out using the model proposed by Reverchon. This model was used successfully to describe our data on pennyroyal [1] and other aromatic plants [2]. Another successful model with two directly adjustable parameters, which are the external mass transfer coefficient,  $k_f$ , and the internal mass transfer coefficient,  $k_s$ , was developed by Sovová [3]. A development of this model, considering the extended Lack's plug flow model for vegetable oil extraction, was reported by Sovová [4], using one adjustable parameter, the external mass transfer coefficient,  $k_s$ . The yields of volatile oil from pennyroyal as function of extraction time described by both models [3,4] for different particle sizes are presented in Figures 1 and 2, respectively.




Fig. 1. Pennyroyal essential oil yield for different mean particle size and a CO<sub>2</sub> flow rate of  $4.7 \times 10^{-4} \text{ kg.s}^{-1}$ . Continuous curves from model [3],  $k_s = 0.89 \times 10^8 \text{ m.s}^{-1}$  and  $k_f = 1.89 \times 10^6 \text{ m.s}^{-1}$




Fig. 2. Pennyroyal essential oil yield for different mean particle size and a CO<sub>2</sub> flow rate of  $4.7 \times 10^{-4} \text{ kg.s}^{-1}$ . Continuous curves from model [4],  $k_s = 1.50 \times 10^8 \text{ m.s}^{-1}$

The experimental results obtained at three different particles sizes using two or only one adjustable parameter, provided a fairly good agreement between the model curves and the experimental data, as shown in Figures 1 and 2, which evidence the capability of these models.

1. E Reis-Vasco et al. (2000) Chem. Eng. Sci. 55: 2917-2922.
2. C Grosso et al. (2010), Chem. Eng. Sci. 65: 3579-3590.
3. H Sovová (1994) Chem. Eng. Sci. 49: 409-414.
4. H Sovová (2005). J. Supercritical Fluids 33, 35-52.

**P 211. Extraction of Volatile Oils from Aromatic Plant with Supercritical Carbon Dioxide**

Palavra AF<sup>1</sup>, Matos P<sup>1</sup>, Nobre BP<sup>1</sup>, Fareleira JMNA<sup>1</sup>, Barroso JG<sup>3</sup>, Coelho JP<sup>2</sup>

<sup>1</sup>Centro de Química Estrutural, Instituto Superior Técnico, Av. Rovisco Pais, 1, 1049-001 Lisboa, Portugal

<sup>2</sup>Chemical Engineering and Biotechnology Research Center, Instituto Superior de Engenharia de Lisboa, Rua Conselheiro Emídio Navarro, 1, 1959-007 Lisboa, Portugal

<sup>3</sup>UL, Faculdade de Ciências de Lisboa, DBV, IBB, Centro de Biotecnologia Vegetal, C2, Campo Grande, 1749-016 Lisboa, Portugal

[jcoelho@deq.isel.ipl.pt](mailto:jcoelho@deq.isel.ipl.pt)

Keywords: Supercritical extraction, hydrodistillation, volatile oils, aromatic plants

In recent years supercritical fluid extraction (SFE) has been used to obtain oils from aromatic plants, as an alternative to the traditional techniques, namely, hydrodistillation (HD) and steam distillation (SD).

The aim of this communication is to discuss an overall view of the supercritical extraction of volatile oils of the aromatic plants, rosemary (*Rosmarinus officinalis*) [1], pennyroyal (*Mentha pulegium*) [2], fennel (*Foeniculum vulgare*) [3], coriander (*Coriandrum sativum*) [4], savory (*Satureja fruticosa*) [5], winter savory (*Satureja montana*) [6], cotton lavender (*Santolina chamaecyparissus*) [7] and thyme (*Thymus vulgaris*) [8] carried out in our laboratories.

This separation technique was performed using a flow apparatus, provided with a 1L-extraction vessel and two separators of 0.27 L each. The oil obtained by supercritical extraction is actually designated as volatile oil to differentiate it from essential oil, which, by definition, is mainly produced by HD and SD.

Different conditions of pressure (90 and 100 bar), temperature (40 and 50 °C), mean particle sizes (0.4, 0.6 and 0.8 mm) and CO<sub>2</sub> flow rate (0.8, 1.1 and 1.3 kg/h) were studied in order to understand the influence of these parameters on the composition and yield of the oils. The results were compared with those obtained for the essential oil isolated by HD. The volatile and essential oils were analysed by GC and GC-MS.

One major difference between those types of oils respect to the absence of waxes in the essential oils. Another important difference is the higher concentration of thymoquinone in volatile oils from thyme (*Thymus vulgaris*) and savoury (*Satureja montana*). In the latter aromatic plant the content in volatile oil can reach a value 15 times higher (SFE- 3%) than the essential oil (HD-0.2%). It is noteworthy that thymoquinone has shown to have important antioxidant and anti-cancer activities.

1. JAP Coelho et al (1996) ACS Symposium Series 670: 101–109.

2. E Reis-Vasco et al. (2000) Chem. Eng. Sci. 55: 2917-2922.

3. JAP Coelho et al. (2003) Flavour and Fragrance J. 148: 316–319.

4. C Grosso et al. (2008) Food Chemistry 111: 197–203.

5. JA Coelho et al. (2007) Flavour and Fragrance J. 22: 438–442.

6. C Grosso et al. (2009) J. Separation Science 32: 328–334.

7. C Grosso et al. (2009) J. Separation Science 32: 3215–3222.

8. C Grosso et al. (2010) J. Separation Science 33: 2211–2218.

**P 212. Chemical composition and trypanocidal activity evaluation of essential oils from *Hedychium coronarium* against *Trypanosoma brucei* strains**

Rodrigues DF<sup>1</sup>, Arenas Velásquez AM<sup>1</sup>, Salgueiro L<sup>4</sup>, Cavaleiro C<sup>4</sup>, Martins GZ<sup>2</sup>, Magalhães NO<sup>2</sup>, Santos LE<sup>2</sup>, Martins MBG<sup>3</sup>, Cicarelli RMB<sup>1</sup>, Moreira RRD<sup>2</sup>

<sup>1</sup>Estadual Paulista University, Faculty of Pharmaceutical Science, Department of Biological Sciences, 14.801-902, SP, Brazil.

<sup>2</sup>Estadual Paulista University, Faculty of Pharmaceutical Sciences, Department of Active Principles Natural and Toxicology, 14.801-902, SP, Brazil.

<sup>3</sup>Estadual Paulista University, Experimental Campus of Litoral Paulista, 11.330-900, SP, Brazil

<sup>4</sup>C.E.F./ Faculty of Pharmacy, University of Coimbra, Portugal

cavaleir@ff.uc.pt

Keywords: *Hedychium coronarium*, Caryophyllene oxide, 1,8-cineole, *Trypanosoma brucei*, Sleeping sickness.

*Trypanosoma brucei* is the etiologic agent of the sleeping sickness transmitted by flies of *Glossina* genus, known as tsetse flies. The search of new trypanocidal compounds is important to develop new drugs more effective and with less adverse effects than pentamidine. In this work, the trypanocidal activity of essential oils from the leaves and rhizomes of *Hedychium coronarium* was investigated using *T. brucei* procyclic strains. The essential oils were prepared by hydrodistillation (4 h) in a Clevenger-type apparatus and the compositions established by gas chromatography and gas chromatography-mass spectroscopy analysis. Caryophyllene oxide and 1,8-cineole are, respectively, the major constituents of the oils from the leaves and rhizomes. Procyclic forms of *T. brucei* (427 and 29-13 strains) were treated with essential oils to determine the cytotoxicity index (CI<sub>50</sub>), using the colorimetric method of MTT. Both essential oils revealed a weak activity (CI<sub>50</sub>>100 µg.ml<sup>-1</sup>) when compared to that of pentamidine (CI<sub>50</sub> = 2.19 µg.ml<sup>-1</sup>), used as control. However, when testing individually the major constituents of the oils, caryophyllene oxide showed a promising activity on both *T. brucei* strains (*T. brucei* 427, CI<sub>50</sub>=65.77 µg.ml<sup>-1</sup> and *T. brucei* 29-13, CI<sub>50</sub>=24.53 µg.ml<sup>-1</sup>).

Acknowledgements: Authors are grateful to Fundação para a Ciência e Tecnologia (FCT), POCI 2010/FEDER and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for financial support.

1. MBG Martins et al. (2010) Rev. Bras. Plant Med., 12: 179-187.

2. F Cotinguiba et al. (2009) Med. Chem. Res., 18: 703-711.

**P 213. Chemical composition and immunomodulatory activity of the essential oil of *Melampodium divaricatum* (RICH. IN PERS.) DC (Asteraceae) and its major compound**

Moreira RRD<sup>1</sup>, Magalhães NO<sup>1</sup>, Santos LE<sup>1</sup>, Botelho VT<sup>1</sup>, Martins GZ<sup>1</sup>, Nepomuceno ALA<sup>1</sup>, Santos AG<sup>1</sup>, Carlos IZ<sup>2</sup>, Matos DC<sup>2</sup>, Andrade G<sup>3</sup>, Martins CHG<sup>3</sup>, Pelissari GP<sup>4</sup>, Salgueiro L<sup>5</sup>, Cavaleiro C<sup>5</sup>

<sup>1</sup>Department of Active Principles Natural and Toxicology, 14.801-902, SP, Brazil.

<sup>2</sup>Departament of Clinical Analysis, Scholl of Pharmaceutical Science/UNESP – Universidade Estadual Paulista - CEP 14901-502 - Araraquara – SP - Brazil. [moreirar@fcar.unesp.br](mailto:moreirar@fcar.unesp.br)

<sup>3</sup>Research Laboratory of Applied Microbiology, LaPeMA – Unifran - University of Franca – Franca-SP - Brazil

<sup>4</sup>University Lutheran Center of Palmas, Palmas- TO - Brazil

<sup>5</sup>C.E.F./ Faculty of Pharmacy, University of Coimbra, Portugal

[ligia@ff.uc.pt](mailto:ligia@ff.uc.pt)

Keywords: *Melampodium divaricatum*, essential oil, nitric oxide, hydrogen peroxide, E-Caryophyllene

*Melampodium divaricatum* (Rich. In Pers.) DC (Asteraceae) is a native plant of South America, popularly known as falsa-calendula and commonly used as diaphoretic and diuretics in the treatment of leucorrhoea. The main goal of the present study was the evaluation of the immunomodulatory activity of essential oil of *M. divaricatum* by determining nitric oxide (NO) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) released in Swiss mice peritoneal macrophages cells. E-Caryophyllene (56.0%), germacrene D (12.7%) and bicyclogermacrene (9.2%) were found to be the major constituents. NO production was inhibited at 80% and 42% on the concentrations of 0.075 µl.ml<sup>-1</sup> and 0.04 µl.ml<sup>-1</sup>, respectively. Inhibition process is dose dependent. The production of H<sub>2</sub>O<sub>2</sub> resulting from PMA stimulation was also strongly inhibited in the presence of essential oil of *M. divaricatum*, but not in a dose dependent way. E-Caryophyllene, the major constituent of the oil, did not affect significantly NO and H<sub>2</sub>O<sub>2</sub> release. Therefore, results confirm that the immunomodulatory activity of essential oil of *M. divaricatum* can be due to minor constituents or to the complex of the substances. The dose-dependent inhibition of macrophage NO suggests immunomodulatory and / or anti-inflammatory potential to be confirmed in further mechanistic and in vivo studies.

Acknowledgements: Authors are grateful to Fundação para a Ciência e Tecnologia (FCT) and POCI 2010/FEDER for financial support.

1. MF Agra et al. (2007) Rev. Bras. Farmacog. 17: 114 -121.

2. E Pick, D Mizel (1981) J. Immunol. Methods. 46: 211-226.

**P 214. Extraction and identification of volatile compounds from mature fruit *Averrhoa bilimbi* L. by solid phase microextraction (SPME) and GC-MS**

Torquilha HS<sup>1</sup>, Bobeda CRR<sup>1</sup>; Machado AMR<sup>1</sup>; Godoy RLO<sup>2</sup>; Pacheco S<sup>2</sup>; Andressa MS<sup>2</sup>

<sup>1</sup>IFRJ-Campus Nilópolis-Brazil;

<sup>2</sup>Embrapa-Brazil

ronoel@ctaa.embrapa.br

Keywords: *Averrhoa bilimbi*; solid phase microextraction; volatile compounds.

The fruit *Averrhoa bilimbi* L. belonging to the family of Oxalidaceae and is popularly known as biri-biri, gooseberry from China and bilimbi, among others, being useful in the food industry [1]. In Asia it is widely used in traditional medicine for various ills cure as it has anti-bacterial pharmacological properties [2] and have been also used in diabetes treatment [3]. Volatile compounds were investigated with hydrodistillation followed by solvent extraction [4]. The bilimbi, we had studied was collected in the northwestern of State of Rio de Janeiro (Brazil), near the city of St. Anthony of Padua and transported under refrigeration to the Embrapa's Laboratory of Liquid Chromatography where the analysis took place. The aim of our study was to perform the extraction and identification of volatile constituents of fresh ripe fruit bilimbi using solid phase microextraction (SPME) with polydimethylsiloxane fiber and GC-MS techniques. We found that under analytical conditions used the chromatographic profile revealed only the region of monoterpene hydrocarbons, ethers and esters and some of these have not cited in previous studies. The mass spectra were compared with those of literature and NIST mass spectra library, had been identified 25 substances which six are the majority, among them the monoterpene limonene and substances: ethyl acetate; ethoxyethane, butyl acetate, methyl butanoate, ethyl butanoate. The result of this work shows that it is possible to study the bilimbi volatile compounds by SPME minimizing time of analysis, sample size and possible losses caused by the technique of solvent extraction.

Acknowledgments: To IFRJ by financial and institutional support

1 PA Souza et al. (2009) Rev. Bras. Frutic. 31(4):1190-1195.

2 MN Norhana et al. (2009). Int. J. Food Microbiol. 136 (1):88-94.

3 P Pushparaj et al. (2001) Life Science, 70(5): 535-547.

4 AJ Pino, R Marbot, A Bello. (2004) J. Essent.Oil Res. 16:241-242.

**P 215. Chemical composition and analgesic effect of the essential oil of *Origanum vulgare* L.**

Sheyla Alves Rodrigues<sup>1</sup>, Deisylaine Maria dos Santos<sup>2</sup>, Américo Azevedo de Souza<sup>2</sup>, Péricles Barreto Alves<sup>3</sup>, Darlisson de Alexandria Santos<sup>3</sup>, Charles dos Santos Estevam<sup>2</sup>

<sup>1</sup>Institute of Technology and Research, Tiradentes University, Brazil; Federal Institute of Sergipe, Brazil.;

<sup>2</sup>Department of Physiology, Federal University of Sergipe, Brazil;

<sup>3</sup>Federal University of Sergipe - Brazil

americosouza\_bio@yahoo.com.br

Keywords: Volatile compounds; oregano; nociceptive

Essential oils are compounds produced by secondary metabolism of plants and used for their pharmacological activities and aromatic value<sup>1</sup>. *Origanum vulgare* L., known as oregano or wild marjoram, is an aromatic plant widely used in cookery. Infusions of the leaves are recommended against liver diseases; digestive, spasmolytic and respiratory disorders<sup>2</sup>. The essential oil this plant is known as an interesting source of antimicrobial compounds to be applied in food conservation and some of its activities related to the compounds carvacrol and thymol<sup>3</sup>. The aim of this work was to investigate the chemical composition and the antioxidant and analgesic activities of essential oil of *O. vulgare* L.

The essential oil of dried leaves from *O. vulgare* was obtained by hydrodistillation on a Clevenger-type apparatus for approximately 3h. The Chemical and physical properties the essential oil was analyzed for Gas chromatography-mass spectrometry (GC-MS) and gas chromatography (GC-FID)<sup>4</sup>. The antioxidant activity of extracts was evaluated using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) assay system<sup>5</sup>. For the abdominal writhings tests, the writhings were induced by the intraperitoneal administration of acetic acid 0,6 %, in male albino Swiss mice (20-25 g) in 100, 200 mg/kg dosage, that was intraperitoneal administered, thirty minutes before performing the experiment in six-animal groups<sup>6</sup>.

The major compounds of essential oil of *O. vulgare* were Terpinen-4-ol (35.73%), g-terpinene (15.58%), thymol (9.01%) and a-terpinene (8.39%). The oil did not show antioxidant activity, but exhibited an analgesic effect in the doses tested with inhibition of 100% in the number of writhings compared with the saline control group (34.83 ±1.94) with p <0.001.

The essential oil of *Origanum vulgare* showed be not only a similar literature composition<sup>3</sup>, but also a potential analgesic.

1. Kintzios SE. (2004) In: Peter, K.V. (ed.) Handbook of herbs and spices (vol. 2). Cambridge: Woodhead Publishing Ltda., 215-229p.
2. Meneses R et al. (2009) Annals of Clinical Microbiology and Antimicrobials, 8:1-8.
3. Mitchell TC et al. (2010) Ciênc. Tecnol. Aliment., Campinas, 30(3): 755-760.
4. Serafini MR et al. (2012) Journal of Biotechnology and Pharmaceutical, 3(1): 1-9.
5. Sousa CMM et al. (2011) Quim. Nova, 30(2): 351-355.
6. Almeida RN. (2006) Psicofarmacologia: fundamentos práticos. 1ª Ed. Guanabara Koogan, Rio de Janeiro.



**P 216. Comparative GS-MS volatile profile of different origin chrism samples**

Claudia-Crina Toma<sup>1</sup>, Sonia Socaciu<sup>2</sup>; F.I. Oros<sup>1</sup>; D. Hanganu<sup>3</sup>; A. Tofana<sup>2</sup>; C. A. Cotoraci<sup>4</sup>; C. Morgovan<sup>5</sup>; Neli K. Olah<sup>6</sup>

<sup>1</sup>Department of Pharmacognosy, Phytochemistry, Phytotherapy, Faculty of Pharmacy, "Vasile Goldis" Western University of Arad, 1 Feleacului Street, 300041, Romania;

<sup>2</sup>Department of Food Science and Technology, University of Agricultural Sciences and Veterinary Medicine, 64 Calea Floresti street, Cluj-Napoca, Romania;

<sup>3</sup>Department of Pharmacognosy, Faculty of Pharmacy, "Iuliu Hațieganu" University of Medicine and Pharmacy, Cluj – Napoca, 15 I. Creangă Street, Romania;

<sup>4</sup>Faculty of Medicine, "Vasile Goldis" Western University of Arad, 1 Feleacului Street, 300041;

<sup>5</sup>Department of Pharmaceutical Management, Marketing and Legislation, Faculty of Pharmacy, "Vasile Goldis" Western University of Arad, 1 Feleacului Street, 300041;

<sup>6</sup>Department of Pharmaceutical Industry, Faculty of Pharmacy, "Vasile Goldis" Western University of Arad, 1 Feleacului Street, 300041 and SC PlantExtrakt SRL, 407059, Rădaia, Jud. Cluj, Romania

claudiatoma2004@yahoo.com

Keywords: Great Chrism, Nard Chrism, Greek Orthodox Chrism, GC-MS

There are several types of chrism used in the different orthodox religious practice. The most important is the Great and Holy Chrism because of the complexity of the chemical composition and religious significations. The preparatory rites began on the morning of Great and Holy Monday, 9<sup>th</sup> of April, and continued through Great and Holy Wednesday. Clergy stirred the Chrism — a mixture of dozens of oils, herbs, fragrances — continuously, day and night, as Scripture was read. In this study we intended to compare four different Chrism types from four different origin and give some explanations for that.[1]

The volatile oil profile was evaluated by GC-MS using headspace injection at 85<sup>o</sup>C, for 15 min, ZB-5MS 50 m x 0,32 mm x 0,25 microm capillary column respectively helium as carrier gas. The separated compounds were identified using an MS spectra library. The quantitative determination was performed by normalization.[2]

The volatile compounds found in chrism samples varied from sample to sample. Thus, in the case of Great Chrism sample, 32 compounds were separated the main ones being alpha-pinene, 1,8-cineol and beta-linalool. Instead, from the Nard Chrism (Israel) sample, 35 compounds were separated from which 32 identified, the major ones being: diphenyl ether, benzyl acetate, phenylethyl alcohol, alpha-terpineol, limonene. From the Greek Orthodox Chrism sample, 36 compounds were identified from the total of 38 separated and beta-linalool, cyclohexane isothiocyanate, as well as limonene being found in high concentrations. The Orthodox Chrism sample from Arad-Gai was characterized by high contents of phenylethyl alcohol, beta-linalool, benzyl acetate and alpha-terpineol.

The difference between chemical composition of the studied chrism samples can be explain by using different herbs or the local variation in composition of the used herbs.

1. Arhieraticon, adica rinduiala slujbelor savirsite de arhiereu, Ed. Institutului Biblic si de Misiune al Bisericii Ortodoxe Romane, Bucuresti, 1993, p.190-196

2. European Pharmacopoeia 7 (2011) Medpharm Scientific Publisher, Stuttgart.

**P 217. Analysis on floral scent of 9 *Syringa* species and varieties by ATD-GC/MS**

Zhao jing, Hu Zenghui, Leng Pingsheng, Li Xiaolong, Liu Jianbin

College of Forestry and Landscape Architecture. Beijing University of Agriculture

[lengpsh@tom.com](mailto:lengpsh@tom.com)

Keywords: *Syringa*; Floral scent; ATD-GC/MS

*Syringa* is an important aromatic woody flower, its dry flowers are traditionally used in tea and spices in China[1]. This study is to determine the volatile components and emission pattern of *Syringa* floral scent. The floral volatiles of *Syringa chinensis*, *S. protolaciniata*, *S. oblata*, *S. oblata* var. *Giraldii*, *S. oblata* var. *plena*, *S. vulgaris* 'Mrs Harry Bickle', *S. vulgaris* 'Bright Centennial', *S. vulgaris* 'White Spires', and *S. vulgaris* 'Pres Lincoln' were collected in vivo by dynamic headspace, and then were identified by ATD-GC/MS[2]. The results showed that except *S. chinensis* (29) and *S. vulgaris* 'Pres Lincoln' (38), more than 50 compounds were detected in floral volatiles of other 7 *Syringa* plants, and the *S. oblata* var. *Giraldii* emitted the most compounds with 61. Among the volatile components of all *Syringa* studied, terpenoids showed the highest relative amount with more than 50% of total release amount, even arrived at 88% in *S. vulgaris* 'Bright Centennial'. And the relative amount of  $\alpha$ -ocimene was over 70% in *S. protolaciniata*, *S. oblata* var. *Giraldii*, and *S. vulgaris* 'Bright Centennial'. Benzenoids and their derivatives were also important volatile components, whose relative amounts varied from 8%-22%, benzaldehyde with 6%-18% of total release amount was the most abundant benzenoids for those 6 *Syringa* plants except *S. oblata* and its varieties. Aldehydes, alcohols, esters, and fatty hydrocarbons were all detected in all *Syringa*, but their release amounts were below 8%. And ketones (1.8%) were only detected in *S. oblata* var. *Giraldii*. In addition, 4 cultivars of *S. vulgaris* were chosen to evaluate the aromatic degrees of the floral scents through smelling by persons. The flowers of *S. vulgaris* 'White Spires' emitted the richest fragrance, followed by *S. vulgaris* 'Mrs Harry Bickle' and *S. vulgaris* 'Bright Centennial', and the floral fragrance of *S. vulgaris* 'Pres Lincoln' was too low to be smelled. Moreover,  $\alpha$ -pinene and alcohols, especially 2-ethyl-1-hexanol, were considered to contribute mostly to aromatic difference of these 4 *Syringa* cultivars. The results provided basis for revealing the main components and their role in aromatic degree, as well as the intraspecific and interspecific difference of floral scent in *Syringa*.

Acknowledgments: The work was supported by National Natural Science Foundation of China

1. Defeng Wang(1979), Biomass Chemical Engineering, 6:369-370

2. HJ Chen, R Hong, YJ Jin et al(2003), J. Instru. Analysis, 22(supp.):226- 228

**P 218. Changes in the essential oil composition and their biological activities during the maturation of *Schinus terebinthifolius* fruits**

Asma Ennigrou

Faculty of Science of Tunis

asma.ennigrou@gmail.com

Keywords: *Schinus terebinthifolius*; fruit maturation; essential oil composition; GC-MS; antioxidant; antibacterial

*Schinus terebinthifolius* commonly known as Brazilian pepper tree is a potential producers of essential oils which have numerous biological activities such as antibacterial, antifungal, antiviral and antitumoral, among others. The chemical composition of the essential oil of the fruits and leaves of *S. terebinthifolius* has been the subject of numerous investigations, and monoterpene hydrocarbons namely  $\alpha$ -phellandrene and  $\beta$ -phellandrene were the prominent compounds. Usually the essential oil is isolated from the leaves and/or the mature fruits. The influence of the stage of maturation of the fruits on the chemical composition of the essential oil of *S. terebinthifolius* has not been reported yet. The aim of the present contribution were to (a) analyze the chemical composition of the essential oils collected at three stages of fruit maturation, and (b) to evaluate the antioxidant and antibacterial activities of the three different oils.

Essentials oils were isolated by hydrodistillation from the air dried fruits collected at three stages of maturation (immature: green; intermediate: green-red and mature: fully red). The oils were dried over anhydrous sodium sulphate and analyzed by gas chromatography-masse spectrometry. At the same time, the essential oils were evaluated for their antiradical property against the DPPH and ABTS radicals and their antibacterial activity.

From the fruits collected at different stages of maturity, a pale yellowish oils with pungent and pepper like aroma were obtained in a yields of 4.6, 4.6 and 4.5% (w/w on dry weight basis), for immature, intermediate and mature fruits, respectively. In all oil samples, Germacrene-D (57.8-59.2%), elixene (9.5-14.5%), terpinolene (7.4-9.08%) and  $\alpha$ -cadinene (3.27-3.9%) were the main components. The oil samples were unable to reduce the DPPH and ABTS radicals, while they found to possess good antibacterial activity against *Enterococcus faecalis*, *Enterococcus faecium* and *staphylococcus aureus*. Among the oils samples, those from the intermediate stage of maturation were found to be the most efficient.

The results obtained herein indicate that *S. terebinthifolius* could be considered as a potential source of volatile oils which could serve not only as flavour agents but also as antiseptic supplements preventing deterioration of foodstuff and beverage products and pharmaceuticals.

**P 219. Method to study the repellent, irritant and toxic effects of essential oils on *Bemisia tabaci* for a combination with insect proof net**

Deletre E<sup>1</sup>, Chandre F<sup>2</sup>; Menut C<sup>3</sup>; Martin T<sup>1</sup>

<sup>1</sup>Cirad, Montpellier;

<sup>2</sup>IRD, Montpellier;

<sup>3</sup>Institut des biomolécules de Max Mousseron, Montpellier

emilie.deletre@cirad.fr

Keywords: behavior; whitefly; tomato; insect proof net; natural product; pest management

*Bemisia tabaci*, vector of the tomato leaf curl virus, is a serious problem in tomato production in many parts of the world. In tropical countries, the use of netting to protect horticultural crops is an effective and sustainable tool against Lepidoptera but not against small pests on the contrary of the Mediterranean region [1]. A previous study showed the efficiency of a repellent net impregnated with alphacypermethrin against the whitefly [2]. But needs to reduce the use of pesticide and resistance in population bring up the issue of finding a natural alternative to alphacypermethrin. The objective of this study was to evaluate the repellent, irritant and toxic effects of 10 essential oils on *B. tabaci* adults in laboratory. The repellent effect due to volatiles compounds was evaluated thanks to a still-air olfactometer. It was expressed as a distance to the essential oil support [3]. The irritant effect which is the contact effect with essential oil was evaluated thanks to a choice test in tube and was illustrated by a different cross rate through an essential oil treated net and a solvent treated net [2]. The toxicity effect was evaluated thanks to a no-choice assay in tube. The toxicity was quantified by the mortality after 24h of *B. tabaci* which crossed an essential oil treated net. Nets were dipped in essential oils at concentrations of 0.01%, 0.1% and 1%. Responses varied according to oil type and dose. The three effects were dose dependent. If one effect showed for an oil it did not mean there was another effect. For example an essential oil can be irritant but not repellent so we could expect that the repellent mechanism are different than the irritant or toxic effect. Several essential oils or a combination of them could be an alternative to alphacypermethrin.

1. P G Weintraub (2009), Physical control : an important tool in pest management programs, Biorational control of arthropods pests, I Ishaaya & A R Horowitz (eds), Springer science
2. Thibaud Martin, Aldi Kamal, Emilie Delétré Romain Bonafos and Serge Simon, (in progress) Repellent effect of an alphacypermethrin treated ent against the whitefly *Bemisia tabaci* Gennadius (Hemiptera :Aleyrodidae)
3. W Zhang, H J McAuslane & D J Schuster (2004) Repellency of ginger oil to *Bemisia argentifolii* (Homoptera: Aleyrodidae) on tomato, *J. Eco Entom.*, 97(4) :1310-1318

**P 220. Synergy between the main compounds of savory and thyme essential oils on control of some plant pathogens**

Lopez-Reyes JG, Spadaro D, Prella A, Gullino ML, Garibaldi A

Centre of Competence for the Innovation in the agro-environmental sector – AGROINNOVA  
University of Turin  
Via Leonardo da Vinci, 44. 10095 Grugliasco - (Turin) – Italy

[giovanny.lopez@unito.it](mailto:giovanny.lopez@unito.it)

Keywords: Bacteria, fungi, pathogen, savory, synergy, thyme.

Among the interesting features of the application of essential oils in post-harvest of plant products, it is significant that their effectiveness in controlling pathogens may be due to a synergy between their components [1]. So the aim of this study was to evaluate the synergy between the main compounds of savory (*Satureja montana*) and thyme (*Thymus vulgaris*) essential oils against two fungal pathogens (*Fusarium fujikuroi* and *Penicillium expansum*) and a bacterial pathogen (*Pseudomonas syringae* pv. *actinidiae*) isolated from diseased plant tissues (from rice cv. Galileo, apples cv. Golden Delicious, and actinidia cv. Hayward, respectively). After a GC-MS analysis, the main components of the savory essential oil resulted to be carvacrol and *p*-cymene (45.1% and 16.1%), while for the thyme essential oil the main components were thymol and  $\alpha$ -pinene (53.7% and 19.8%). Different mixtures of the two main components of both essential oils were tested *in vitro* for their inhibition of pathogen growing using the *agar diffusion* technique [2]. From the obtained results it was possible to highlight the synergy between carvacrol and *p*-cymene, since *F. fujikuroi* and *P. syringae* pv. *actinidiae* inhibition due to the application of the mixtures at 75% -25% and 50% -50% of carvacrol and *p*-cymene was statistically higher than that shown by both compounds applied alone at 100%. Nevertheless, the application of thymol at 100% showed an inhibition statistically higher than the mixtures of thymol and  $\alpha$ -pinene, that behaviour confirmed the thymol-based antimicrobial activity of thyme essential oil. *In vitro* analysis was efficient for screening the potential of essential oils application on plant pathogen control, but *in vivo* trials under commercial conditions are necessary to achieve a better response for the agricultural sector.

1. P Tripathi, NK Dubey (2004) Postharvest Biol. Tec. 32: 235-245.
2. S Inouye et al. (2006) Jpn. J. Med. Mycol. 47:91-98.

**P 221. Antifungal activity of savory essential oil by fumigation in apple storage.**

Lopez-Reyes JG, Spadaro D, Gullino ML, Garibaldi A.

Centre of Competence for the Innovation in the agro-environmental sector – AGROINNOVA  
University of Turin  
Via Leonardo da Vinci, 44. 10095 Grugliasco - (Turin) – Italy

[giovanny.lopez@unito.it](mailto:giovanny.lopez@unito.it)

Keywords: Apple, *Botrytis cinerea*, fumigation, *Penicillium expansum*, savory.

Application of essential oils in post-harvest of apples has been reported as an alternative method to control fungal pathogens such as *Botrytis cinerea* and *Penicillium expansum* [1]. Besides application by spraying or dipping, the antimicrobial activity of essential oils in the vapour phase makes them possible fumigants within the structures of fruit storage [2]. So the aim of this study was to evaluate the antifungal activity of savory (*Satureja montana*) essential oil as a fumigant in storage of apples cv. Golden Delicious against *B. cinerea* and *P. expansum*. The apples were disinfected, punctured with a sterile plastic tip at the equatorial region, and inoculated by spraying a suspension of  $1 \times 10^5$  conidia  $\text{ml}^{-1}$  of each pathogen. Inoculated apples were stored separately in hermetic cabinets (60x75x75cm) for 30 days: a. with 50 ml of a 10% emulsion of savory essential oil (10% essential oil, 88% sterilized water and 2% Tween20); b. with 50 ml of a 10% emulsion of savory essential oil (10% essential oil, 88% sterilized water and 2% Tween20) gelified with agar-agar (15g per liter); c. alone as a control. After storage, apples were brought out and kept 7 days in *shelf-life* at room temperature. From the obtained results it was possible to observe that the diameter of the fungal rot on the apples stored with the essential oil emulsion was statistically lower than the control. The emulsion of savory essential oil was lightly less effective on pathogen control than the gelified one, but better results were found against *B. cinerea* than against *P. expansum*, even after *shelf-life*. A better comprehension of the fungicidal and fungistatic activity of essential oils could define their real efficacy against post-harvest pathogens of fruit and their possible commercial use.

1. Lopez-Reyes JG et al. (2010) Flavour Fragr. J. 25: 171–177.

2. P Tripathi, NK Dubey (2004) Postharvest Biol. Tec. 32: 235-245.

## Indexes







## Species

<i>A. ageratifolia</i> .....	96
<i>A. baumannii</i> .....	91
<i>A. niger</i> .....	171, 221, 242
<i>Acca sellowiana</i> .....	253
<i>Achillea clypeolata</i> .....	81
<i>Achillea serbica</i> .....	96
<i>Acinetobacter</i> .....	69
<i>Adansonia</i> .....	202
<i>Aedes aegypti</i> .....	115, 116, 189
<i>Ageratum conyzoides</i> .....	85, 181
<i>Agrobacterium tumefaciens</i> .....	52
<i>Aldama filifolia</i> .....	156
<i>Aldama linearifolia</i> .....	153
<i>Aloysia gratissima</i> .....	248
<i>Aloysia polystachya</i> .....	134
<i>Alphitobius diaperinus</i> .....	130
<i>Alpinia galanga</i> .....	246
<i>Alpinia zerumbet</i> .....	188
<i>Alternaria alternata</i> .....	168, 221
<i>Amaranthus palmeri</i> .....	25
<i>Amaranthus retroflexus</i> .....	231
<i>Ammoides verticillata</i> .....	218
<i>Ananas comosus</i> .....	192
<i>Andereaobryum macrosporum</i> .....	241
<i>Anethum graveolens</i> .....	131
<i>Annona crassiflora</i> .....	101
<i>Annona paludosa</i> .....	101
<i>Anopheles atroparvus</i> .....	247
<i>Anthemis mixta</i> .....	62
<i>Arabidopsis thaliana</i> .....	217, 229
<i>Aristolochia cymbifera</i> .....	186
<i>Aristolochia trilobata</i> .....	120
<i>Armoracia rusticana</i> .....	95
<i>Artemisia abrotanum</i> .....	93
<i>Artemisia absinthium</i> .....	93, 108
<i>Artemisia afra</i> .....	137
<i>Artemisia dracunculul</i> .....	93
<i>Artemisia dubia</i> .....	181
<i>Artemisia indica</i> .....	181
<i>Artemisia vulgaris</i> .....	93, 181
<i>Aspergillus flavus</i> .....	221
<i>Aspergillus fumigatus</i> .....	84, 137
<i>Aspergillus niger</i> .....	154, 171, 242
<i>Astrantia major</i> .....	73
<i>Averrhoa bilimbi</i> .....	258
<i>Baccharis dracunculifolia</i> .....	60, 152
<i>Bacillus cereus</i> .....	154, 224, 226
<i>Bacillus subtilis</i> .....	47, 55, 243
<i>Bemisia tabaci</i> .....	263
<i>Biomphalaria glabrata</i> .....	115
<i>Blainvillea dichotoma</i> .....	85
<i>Blumea lacera</i> .....	181
<i>Bothrops neuwiedi</i> .....	225
<i>Botrytis cinerea</i> .....	48, 117, 201, 208, 265
<i>Brassica nigra</i> .....	25
<i>Bursaphelenchus xylophilus</i> .....	72, 183
<i>Butia capitata</i> .....	253
<i>C. albicans</i> .....	38, 91, 195, 208, 209, 226, 246
<i>C. chaetocalyx</i> .....	99
<i>C. draconoides</i> .....	99
<i>C. eriocladius</i> .....	99
<i>C. glandulosus</i> .....	99
<i>C. krusei</i> .....	91
<i>C. limon</i> .....	78
<i>C. martii</i> .....	214
<i>C. reticulata</i> .....	214
<i>C. torulosa</i> .....	135
<i>C. tropicalis</i> .....	91
<i>Callistemon viminalis</i> .....	110
<i>Callitris intratropica</i> .....	139
<i>Calycolpus goetheanus</i> .....	71
<i>Candida</i> 38, 47, 55, 69, 84, 138, 154, 158, 159, 208, 209, 246	
<i>Candida albicans</i> .....	69, 158
<i>Carum carvi</i> .....	252
<i>Cassia ferruginea</i> .....	237
<i>Cassia lutea</i> .....	237
<i>Cassia occidentalis</i> .....	237
<i>Cassia siemens</i> .....	237
<i>Centaurea vlachorum</i> .....	147
<i>Chaetomium sp.</i> .....	221
<i>Chamaemelum mixtum</i> .....	62
<i>Chamomilla recutita</i> .....	158
<i>Chlamydomonas sp.</i> .....	205
<i>Choisya ternata</i> .....	82
<i>Chrysanthemum cinerifolium</i> .....	181
<i>Cinnamomum durifolium</i> .....	236
<i>Cinnamomum magnificum</i> .....	236
<i>Cinnamomum sericans</i> .....	236
<i>Cinnamomum zeylanicum</i> .....	168
<i>Cinnamomun zeylanicum</i> .....	134, 145
<i>Cirsium acaule</i> .....	190
<i>Cirsium decussatum</i> .....	190
<i>Cirsium eriophorum</i> .....	190
<i>Cirsium erisithales</i> .....	190
<i>Cirsium ferox</i> .....	190
<i>Cirsium heterophyllum</i> .....	190
<i>Cirsium hispanicum</i> .....	190
<i>Cirsium ligulare</i> .....	190
<i>Cirsium pannonicum</i> .....	190
<i>Cirsium rivulare</i> .....	190
<i>Citrus depressa</i> .....	220
<i>Citrus madurensis</i> .....	220
<i>Cladanthus mixtus</i> .....	62
<i>Cladosporium cucumerinum</i> .....	117, 208
<i>Cladosporium sphaerospermum</i> .....	102, 103
<i>Cleistopholis patens</i> .....	124
<i>Conium maculatum</i> .....	74
<i>Conyza canadensis</i> .....	160
<i>Copaifera duckei</i> .....	214
<i>Copaifera langsdorffii</i> .....	133
<i>Croton cajucara</i> .....	208
<i>Croton campestris</i> .....	99
<i>Cryptomeria japonica</i> .....	192, 201
<i>Cucurbita pepo</i> .....	194
<i>Cuminum cyminum</i> .....	247
<i>Cupressus tonkinensis</i> .....	135
<i>Cylas spp.</i> .....	206
<i>Cymbopogon citratus</i> .....	51, 72, 134, 168, 246
<i>Cymbopogon winterianus</i> .....	58, 114, 162
<i>Dalea strobilacea</i> .....	196
<i>Digitaria sanguinalis</i> .....	231
<i>Drimys angustifolia</i> .....	178, 248
<i>Duguetia echinophor</i> .....	101
<i>E. absinthoides</i> .....	83
<i>E. amethystinum</i> .....	141
<i>E. bosistoana</i> .....	193, 216
<i>E. botryoides</i> .....	193, 216
<i>E. camaldulensis</i> .....	193, 216
<i>E. ciconium</i> .....	83
<i>E. cicutarium</i> .....	83
<i>E. cinerea</i> .....	193, 216
<i>E. cordieri</i> .....	216
<i>E. faecalis</i> .....	38, 91, 195
<i>E. ficifolia</i> .....	216
<i>E. gigantea</i> .....	216
<i>E. globulus</i> .....	216
<i>E. hemiphloia</i> .....	216
<i>E. macarthurii</i> .....	216
<i>E. patrisii</i> .....	71
<i>E. pauciflora</i> .....	216
<i>E. piauhiensis</i> .....	71
<i>E. piperita</i> .....	216
<i>E. polyanthemos</i> .....	216
<i>E. radiata</i> .....	216
<i>E. saligna</i> .....	216
<i>E. smithii</i> .....	216
<i>E. urophylla</i> .....	216
<i>E. viminalis</i> .....	216
<i>Enterococcus</i> .....	69, 262
<i>Eremanthus erythropapus</i> .....	107

<i>Erodium</i> .....	83	<i>Lippia microphylla</i> .....	163
<i>Erwinia</i> sp. ....	251	<i>Lippia organoides</i> .....	219
<i>Eryngium alpinum</i> .....	141	<i>Lippia sidoides</i> .....	179
<i>Eryngium dilatatum</i> .....	155	<i>Listeria monocytogenes</i> 76, 113, 142, 193, 194, 218	
<i>Eryngium duriae</i> .....	249	<i>Lychnophora pinaster</i> .....	142
<i>Eryngium pandanifolium</i> .....	155	<i>Lycopus europaeus</i> .....	67
<i>Eucalyptus</i> .....	72, 137, 145, 193, 216, 221	<i>M. atramentifera</i> .....	71
<i>Eucalyptus citriodora</i> .....	72	<i>M. avium</i> .....	91
<i>Eucalyptus grandis</i> .....	137	<i>M. bracteata</i> .....	71
<i>Eucalyptus</i> spp. ....	145, 216	<i>M. tuberculosis</i> .....	91
<i>Eugenia punicifolia</i> .....	71	<i>Machilus velutina</i> .....	236
<i>Eugenia uniflora</i> .....	170	<i>Margotia gummifera</i> .....	148
<i>Eupatorium adenophorum</i> .....	181	<i>Matricaria chamomilla</i> .....	181
<i>Eupatorium squalidum</i> .....	85	<i>Melaleuca alternifolia</i> .....	228, 248
<i>Euphorbia heterophylla</i> .....	58, 114	<i>Melampodium divaricatum</i> .....	257
<i>Ferula ovina</i> .....	77	<i>Melissa officinalis</i> .....	210
<i>Foeniculum vulgare</i> .....	49, 116, 247	<i>Mentha arvensis</i> .....	72
<i>Fusarium culmorum</i> .....	117	<i>Mentha cervina</i> .....	207
<i>Fusarium fujikuroi</i> .....	264	<i>Mentha longifolia</i> .....	109
<i>Fusarium oxysporum</i> .....	89, 137	<i>Mentha pulegium</i> .....	116, 254, 255
<i>Fusarium solani</i> .....	100	<i>Mentha</i> spp. ....	134
<i>G. columbinum</i> .....	83	<i>Mentha suaveolens</i> .....	109
<i>G. lucidum</i> .....	83	<i>Micrococcus luteus</i> .....	55
<i>G. macrorrhizum</i> .....	83	<i>Mimosa pudica</i> .....	214
<i>G. phaeum</i> .....	83	<i>Mosiera bullata</i> .....	111
<i>G. purpureum</i> .....	83	<i>Mycoplasma mycoides</i> .....	248
<i>G. robertianum</i> .....	83	<i>Myrcia fallax</i> .....	71
<i>G. sanguineum</i> .....	83	<i>Myrtus communis</i> .....	75, 250
<i>Galanthus nivalis</i> .....	90	<i>N. megapotamica</i> .....	225
<i>Geranium</i> .....	83	<i>Nectandra angustifolia</i> .....	225
<i>Gliocladium roseum</i> .....	221	<i>Neolitsea polycarpa</i> .....	236
<i>Glycine max</i> .....	212	<i>Neurospora crassa</i> .....	242
<i>Guatteria schomburgkiana</i> .....	101	<i>Nigella sativa</i> .....	91
<i>H. alutacea</i> .....	70	<i>O. sipyleum</i> .....	61
<i>H. dilatata</i> .....	70	<i>Ocimum americanum</i> .....	204
<i>H. parkeri</i> .....	70	<i>Ocimum basilicum</i> .....	46, 87, 121, 227
<i>H. recurvata</i> .....	70	<i>Ocimum campechianum</i> .....	112
<i>H. suaveolens</i> .....	70	<i>Ocimum gratissimum</i> .....	211
<i>Haematococcus pluvialis</i> .....	205	<i>Ocotea pulchella</i> .....	226
<i>Hedychium coronarium</i> .....	256	<i>Origanum onites</i> .....	61
<i>Helianthus tuberosus</i> .....	66	<i>Origanum virens</i> .....	72, 89
<i>Hornungia petraea</i> .....	84	<i>Origanum vulgare</i> 61, 72, 76, 78, 134, 145, 160, 161, 168, 238, 252, 259	
<i>Hua gabonii</i> .....	234	<i>Ormenis mixta</i> .....	62
<i>Hutchinsia alpina</i> .....	84	<i>Otanthus maritimus</i> .....	45
<i>Hypericum umbellatum</i> .....	244	<i>P. aegerinosa</i> .....	91
<i>Hypericum undulatum</i> .....	174	<i>P. callosum</i> .....	100
<i>Hypochoeris maculata</i> .....	180	<i>P. divaricatum</i> .....	100
<i>Hyptis crenata</i> .....	70	<i>P. krukoffii</i> .....	100
<i>Hyptis pectinata</i> .....	119, 173, 226, 227, 230	<i>Paenibacillus larvae</i> .....	134
<i>Hyssopus seravschanicus</i> .....	224	<i>Papaver rhoeas</i> .....	169
<i>Ichthyothere cunabi</i> .....	85	<i>Papaver somniferum</i> .....	169
<i>Ichthyothere rufa</i> .....	85	<i>Parthenium hysterophorus</i> .....	181
<i>Inula helenium</i> .....	65, 240	<i>Pelargonium graveolens</i> .....	137
<i>Ipomoea batatas</i> .....	206	<i>Pelargonium</i> sp. ....	98
<i>Ipomoea grandifolia</i> .....	59	<i>Penicillium digitatum</i> .....	78
<i>Juniperus communis</i> .....	79, 117	<i>Penicillium expansum</i> .....	137, 264, 265
<i>L. alba</i> .....	57, 209	<i>Penicillium italicum</i> .....	171
<i>L. origanoides</i> .....	102, 219	<i>Penicillium oxalicum</i> .....	171
<i>L. viridis</i> .....	33, 40, 157	<i>Penicillium</i> sp. ....	221
<i>Lactuca sativa</i> .....	231	<i>Penicillium expansum</i> .....	265
<i>Lamium amplexicaule</i> .....	222	<i>Petroselinum crispum</i> .....	247
<i>Lamium purpureum</i> .....	222	<i>Peucedanum austriacum</i> .....	243
<i>Lantana camara</i> .....	213	<i>Philodendron imbe</i> .....	143
<i>Laurus nobilis</i> .....	145	<i>Phoebe angustifolia</i> .....	236
<i>Lavandula luisieri</i> .....	32, 48	<i>Pimpinella pruatjan</i> .....	106
<i>Lavandula pedunculata</i> .....	33, 40	<i>Pinus peuce</i> .....	69
<i>Lavandula</i> spp. ....	33, 41, 157	<i>Pinus pinaster</i> .....	72
<i>Lavandula stoechas</i> .....	157, 203	<i>Pinus</i> spp. ....	183
<i>Lavandula viridis</i> .....	33, 40	<i>Pinus sylvestris</i> .....	136
<i>Leishmania amazonensis</i> .....	115, 162, 172	<i>Piper aduncum</i> .....	47, 227
<i>Leishmania chagasi</i> .....	179	<i>Piper angustifolium</i> .....	47
<i>Leonurus sibiricus</i> .....	223	<i>Piper auritum</i> .....	47
<i>Lepista nuda</i> .....	184	<i>Piper nigrum</i> .....	100
<i>Lippia alba</i> .....	209	<i>Pittosporum undulatum</i> .....	192
<i>Lippia gracilis</i> .....	102, 140, 179	<i>Plectranthus grandidentatus</i> .....	80, 195
<i>Lippia graveolens</i> .....	97		
<i>Lippia javanica</i> .....	137		

<i>Plectranthus porcatus</i> .....	195	<i>Tymus herba-barona</i> .....	203
<i>Plectranthus venteri</i> .....	195	<i>U. connivens</i> .....	198
<i>Pogostemon cablin</i> .....	88	<i>Unxia camphorata</i> .....	85
<i>Porophyllum angustissimum</i> .....	176	<i>Uvaria angustifolia</i> .....	198
<i>Portulaca oleracea</i> .....	160	<i>Uvari dendron angustifolium</i> .....	198
<i>Pseudomonas</i> 55, 76, 142, 150, 154, 158, 191, 193, 224, 226, 243, 246, 251		<i>Vernonia scabra</i> .....	85
<i>Pseudomonas aeruginosa</i> 55, 76, 142, 150, 154, 158, 193, 224, 243, 246, 251		<i>Vetiveria</i> .....	26
<i>Pseudomonas syringae</i> .....	264	<i>Vicia faba</i> .....	231
<i>Psidium cattleianum</i> .....	253	<i>Viguiera filifolia</i> .....	156
<i>Psidium myrsinites</i> .....	71	<i>Viguiera linearifolia</i> .....	153
<i>Pyrus communis</i> .....	94	<i>Xanthium strumarium</i> .....	181
<i>Rhizopus oryzae</i> .....	210, 230	<i>Xylopi aethiopica</i> .....	126
<i>Riencourtia angustifolia</i> .....	85	<i>Xylopi aromatica</i> .....	101, 226
<i>Rosa rugosa</i> .....	128	<i>Xylopi laevigata</i> .....	53, 164
<i>Rosmarinus officinalis</i> .....	76, 122, 145, 245, 255	<i>Xylopi rubescens</i> .....	123
<i>Ruta graveolens</i> .....	72	<i>Yarrowia lipolytica</i> .....	76
<i>S. austrocaledonicum</i> .....	182		
<i>S. cuneifolia</i> .....	104		
<i>S. epidermidis</i> .....	55		
<i>S. innota</i> .....	104		
<i>S. intricate</i> .....	104		
<i>S. montana</i> .....	55, 72, 104		
<i>S. oblata</i> .....	261		
<i>S. protolaciniata</i> .....	261		
<i>S. spicatum</i> .....	182		
<i>S. vulgaris</i> .....	261		
<i>Salicornia ramosissima</i> .....	146		
<i>Salmonella abony</i> .....	158		
<i>Salmonella choleraesius</i> .....	142		
<i>Salmonella enterica</i> .....	218		
<i>Salmonella enteritidis</i> .....	69, 113, 227		
<i>Salvia lavandulaefolia</i> .....	76		
<i>Salvia nemorosa</i> .....	144		
<i>Salvia officinalis</i> .....	56, 76		
<i>Salvia sclarea</i> .....	76		
<i>Santalum album</i> .....	29, 182		
<i>Santolina chamaecyparissus</i> .....	255		
<i>Sarcocornia perennis</i> .....	146		
<i>Satureja</i> .....	55, 72, 104, 255, 264, 265		
<i>Satureja montana</i> .....	55, 72, 255, 264, 265		
<i>Scandix pecten-veneris</i> .....	63		
<i>Schinus molle</i> .....	159		
<i>Schinus terebinthifolius</i> .....	262		
<i>Senecio vernalis</i> .....	64		
<i>Senna obtusifolia</i> .....	214		
<i>Seseli rigidum</i> .....	154		
<i>Solandra maxima</i> .....	54		
<i>Sphagneticola trilobata</i> .....	103		
<i>Spodoptera litoralis</i> .....	137		
<i>Staphylococcus epidermidis</i> .....	69		
<i>Stenocarus ruficornis</i> .....	169		
<i>Streptococcus agalactiae</i> .....	69		
<i>Streptococcus bovis</i> .....	246		
<i>Streptococcus pyogenes</i> .....	69		
<i>Syringa chinensis</i> .....	261		
<i>Syzygium aromaticum</i> .....	72, 168, 200		
<i>Syzygium cumini</i> .....	115		
<i>Syzygium malaccense</i> .....	175		
<i>Takakia lepidozoides</i> .....	241		
<i>Tanacetum balsamita</i> .....	167		
<i>Tanacetum vulgare</i> .....	130		
<i>Tetradenia riparia</i> .....	172		
<i>Tetranychus ludeni</i> .....	143		
<i>Thuja occidentalis</i> .....	130		
<i>Thymbra capitata</i> .....	72, 138		
<i>Thymus caespititius</i> .....	72, 129, 215		
<i>Thymus ciliatus</i> .....	218		
<i>Thymus mastichina</i> .....	127		
<i>Thymus vulgaris</i> .....	19, 72, 227, 255, 264		
<i>Thymus zygis</i> .....	72, 117		
<i>Tilesia baccata</i> .....	85		
<i>Trichophyton rubrum</i> .....	148, 179, 209		
<i>Trichotecium roseum</i> .....	221		
<i>Trypanosoma brucei</i> .....	256		
<i>Trypanosoma cruzi</i> .....	143, 172		

## Subjects

- (8Z,11Z,14Z)-8,11,14-heptadecatrien-2-ol ..... 123  
 (E)-Caryophyllene ..... 163  
<sup>13</sup>C NMR ..... 75, 123, 124  
 1-octen-3-ol ..... 184, 222, 237  
 2-pentyl-3,4,5,6-tetrahydropyridine ..... 74  
 2-undecanone ..... 72  
 3-methyl-2-alkanones ..... 65  
 3-methyl-2-tetradecanone ..... 65  
 6-hydroxydihydrothespirane ..... 86  
 6-methyl-5-hepten-2-yl acetate ..... 120  
 7-hydroxycalamenene ..... 208  
 above-ground parts ..... 64, 81, 96  
 absolute configuration ..... 68, 136  
 ABTS ..... 140, 154, 218, 262  
 accessions ..... 49, 119  
 Acute toxicity ..... 157  
 adulteration ..... 27, 220  
 AFLP ..... 97  
 alantolactone ..... 65, 240  
 alcohols ..... 19, 54, 63, 152, 175, 182, 261  
 alecrim-de-tabuleiro ..... 163  
 alkamide ..... 96  
 Alkylpyridines ..... 86  
 Allelochemicals ..... 25  
 Allelopathy ..... 25, 31, 59, 181, 212, 222  
 allyl isothiocyanate ..... 95  
 Alzheimer's ..... 32, 234  
 Amaryllidaceae ..... 90  
 Ambridge rose ..... 86  
 analgesic ... 51, 77, 118, 142, 157, 159, 162, 191, 259  
 anatomy ..... 156  
 Annonaceae ..... 53, 101, 198, 226  
 Antibacterial activity ..... 38, 193  
 anti-Candidal ..... 208  
 antifungal ..... 22, 38, 45, 48, 55, 84, 89, 91, 100, 102, 103, 117, 137, 142, 148, 155, 159, 178, 179, 187, 203, 208, 209, 210, 221, 230, 262, 265  
 Anti-hypernociceptive ..... 50  
 Anti-inflammatory ..... 33, 157, 249  
 antileishmanial ..... 172, 179  
 Antimycoplasmatic ..... 248  
 antinociceptive ..... 53, 77, 92, 118, 120, 163, 170  
 Antioxidant activity ..... 40, 47, 104, 146, 154, 166, 197, 250  
 antiproliferative ..... 153, 156  
 anti-snake ..... 225  
 anti-staphylococcal activity ..... 65, 240  
 Apiaceae ..... 49, 63, 73, 74, 155, 217, 218, 247  
 Apple ..... 168, 265  
 Araceae ..... 143  
 Arecaceae ..... 253  
 aridity index ..... 112  
 arnica ..... 142  
 aroma ... 18, 22, 41, 95, 121, 122, 128, 131, 132, 161, 167, 184, 185, 187, 200, 226, 234, 239, 245, 252, 253, 262  
 artichoke ..... 66, 132  
 artificial crushing finger device ..... 167  
 Asteraceae ... 45, 62, 64, 65, 66, 81, 85, 93, 96, 103, 142, 147, 152, 153, 156, 176, 180, 181, 183, 240, 257  
 ATD-GC ..... 261  
 Atlantic Rain Forest ..... 103  
 Avoseh NO ..... 237  
 BACE-1 ..... 32  
 Bacteria ..... 191, 264  
 Banwo ID ..... 237  
 Baobab ..... 202  
 bark ..... 124, 139, 175, 198, 201, 208, 234, 251  
 barley ..... 132  
 basil ..... 87, 121  
 behavior ..... 16, 17, 51, 116, 120, 125, 210, 263  
 bergamot ..... 125  
 BHA ..... 47, 197  
 BHT ..... 47  
 bioactivity ..... 117, 181, 195, 227, 230, 231  
 Bioassays ..... 16, 189  
 bioconversion ..... 171  
 biodiversity ..... 31, 204  
 bioherbicide ..... 58, 59, 114, 212  
 Biological activity ..... 22, 31, 172, 231  
 bio-pesticides ..... 247  
 biotransformation ..... 171, 187, 191, 242  
 Bio-valorization ..... 201  
 Blewit ..... 184  
 bornyl acetate ..... 56, 69, 108  
 Brassicaceae ..... 84, 95  
 brine shrimp ..... 31, 157, 159, 181  
 bryophytes ..... 192  
 bubble point ..... 235  
 Buchu ..... 39  
 by-products ..... 166, 194  
 Calamondin ..... 220  
 calamusenone ..... 119, 230  
 calamusenone oxidized analogous ..... 230  
 calcium channels ..... 149  
 Camphor ..... 163, 195  
 cancer ..... 22, 92, 99, 255  
 candeia ..... 107  
 Candidal ..... 208  
 candidosis ..... 138  
 Cape chamomile ..... 39  
 capillary electrophoresis ..... 251  
 caraway ..... 252  
 carbaryl ..... 165  
 Carenko® ..... 136  
 carotenoids ..... 205  
 carrot ..... 168  
 carvacrol ... 19, 33, 54, 55, 61, 71, 72, 92, 97, 104, 129, 145, 160, 179, 203, 215, 217, 218, 219, 221, 238, 259, 264  
 carvone ..... 57, 109, 204, 209, 217  
 caryophyllene oxide ..... 67, 70, 71, 80, 101, 119, 141, 144, 147, 173, 195, 236, 243, 244, 256  
 Caryophyllene oxide ..... 256  
 Cat ear ..... 180  
 CCC ..... 152  
 cedrol ..... 135, 186  
 cell-permeable inhibitors ..... 32  
 Cerrado ..... 103, 133, 176  
 chamomile ..... 39, 45, 62, 158  
 Chemometrics ..... 39  
 chemosystematics ..... 241  
 Chemotaxonomy ..... 73, 83, 225, 237  
 chemotype ..... 19, 30, 31, 49, 57, 73, 81, 97, 105, 108, 109, 129, 144, 158, 160, 170, 181, 204, 208, 209, 215, 219  
 chemotypification ..... 64  
 chicory ..... 132  
 chicory coffee ..... 132  
 chip technology ..... 251  
 Chirality ..... 17, 68  
 chitosan ..... 138  
 cholinesterase ..... 154  
 chondrocytes ..... 249  
 Chrism ..... 260  
 cineole ... 33, 56, 62, 70, 71, 75, 81, 100, 110, 122, 159, 216, 221, 224, 228, 246, 250, 256  
 cinnamon ..... 134, 145, 168, 251  
 Cipó imbé ..... 143  
 circadian activit ..... 219  
 cis-pinocamphone ..... 224  
 citotoxicity ..... 45, 157, 164  
 citral ..... 25, 57, 86, 118, 146, 149, 198, 209  
 citronella ..... 58, 162  
 citronellal ..... 58, 114, 162, 216  
 citronellol ..... 50, 51, 58, 59, 86, 98, 114  
 citrus ..... 121, 165, 177, 220, 232  
 Citrus ..... 78, 177, 220, 232, 233

- climate changes ..... 133  
climatic diversity ..... 98  
clove ..... 168, 200, 251  
Clusiaceae ..... 174, 244  
cluster analysis ..... 49, 62, 64, 73, 81, 99, 133, 161, 216  
Compositae ..... 62, 65  
conformational analysis ..... 240  
coniceine ..... 74  
coniine ..... 74  
conmaculatin ..... 74  
Conservation genetics ..... 207  
Contaminants ..... 27  
correlation ..... 39, 62, 75, 79, 99, 105, 112, 161, 197, 223, 228, 239, 244  
cosmetic ..... 21, 45, 79, 152, 167, 170, 204, 226, 250  
coumarin ..... 241  
coumarins ..... 232  
cow ..... 252  
COX-2 ..... 33  
Cristescu ..... 239  
cucumber ..... 141, 168  
cultivation ..... 39, 61, 192, 199  
culture media ..... 205, 215  
cumin ..... 242, 247  
Cupressaceae ..... 79  
cyclic monoterpenes ..... 52  
cyclocolorenone ..... 178, 248  
Cyclodextrins ..... 40  
Cymbeline ..... 86  
cytotoxic ..... 60, 147, 148, 150, 162, 164, 249  
cytotoxicity ..... 31, 33, 45, 148, 150, 153, 159, 162, 164, 172, 181, 256  
Deception ..... 16  
DEHP ..... 27  
dehydroaromadendrene ..... 230  
Dengue ..... 115, 116, 189  
Dering apparatus ..... 166  
Derivatives ..... 29  
descriptors ..... 49, 240  
dihydroeuparin ..... 66  
Dill ..... 131  
diplophyllin ..... 65, 240  
disease ..... 32, 67, 100, 116, 134, 163, 189, 200, 234, 247  
diurnal variations ..... 109  
DMDS ..... 16  
DMNT ..... 16  
DOSY ..... 125  
DPPH ..... 47, 102, 111, 128, 140, 154, 159, 166, 197, 218, 234, 259, 262  
dry plant ..... 63, 89, 122  
*E/Z*-isomers ..... 29  
*E*-Caryophyllene ..... 257  
Electromyography ..... 188  
enantioselective ..... 41, 136, 245  
environmental factors ..... 109, 112  
escova de garrafa ..... 110  
estoraque ..... 204  
eugenol ..... 100, 146, 198, 211, 227, 229, 251  
Euphorbiaceae ..... 99, 114, 183, 208  
extracellular peptidases ..... 210  
Extraction ..... 28, 62, 91, 185, 186, 255, 258  
extracts ..... 16, 25, 26, 27, 37, 42, 60, 77, 107, 108, 109, 122, 128, 130, 137, 139, 159, 166, 167, 183, 194, 197, 198, 208, 223, 225, 227, 242, 247, 250, 259  
Faba bean ..... 231  
Fabaceae ..... 196, 237  
Facanali R ..... 238  
Farnesols ..... 182  
fast-RP-HPLC ..... 232  
fatty acids ..... 91, 169, 194, 205  
fennel ..... 49, 116, 247, 255  
fertilization ..... 238  
fixed oil ..... 91  
flash aroma dispenser ..... 167  
flavonoids ..... 80, 150, 177, 180  
flavor ..... 26, 132, 181, 220, 226, 233, 239, 253  
Flavourings ..... 27  
Floral scent ..... 16, 261  
Flow cytometry ..... 38  
flowers ..... 16, 54, 80, 81, 86, 90, 109, 110, 127, 130, 144, 161, 167, 174, 190, 202, 213, 218, 219, 261  
Folin-Ciocalteu ..... 104, 250  
food spoilage ..... 48  
Food-borne illness ..... 227  
Foraging ..... 16  
formulations ..... 212, 226, 228  
fresh plant ..... 66, 89, 122, 153, 156  
Fresh-cut salads ..... 146  
fruits ..... 18, 37, 48, 49, 67, 73, 74, 78, 94, 101, 146, 154, 159, 168, 175, 194, 220, 236, 239, 243, 247, 250, 253, 258, 262, 265  
fumigation ..... 234, 265  
fungi ..... 48, 55, 89, 117, 137, 159, 171, 178, 179, 184, 199, 201, 203, 206, 209, 210, 221, 230, 264  
fungicidal ..... 38, 48, 137, 179, 230, 265  
fungicides ..... 17, 165, 223  
fungicides residues ..... 165  
furanoguai-1,4-diene ..... 123  
furocoumarins ..... 232  
Gamma Irradiation ..... 200  
Gene silencing ..... 37  
genetic differentiation ..... 97  
genetic diversity ..... 97, 207  
genotype ..... 95, 105, 161, 230  
Geraniaceae ..... 83  
geraniol ..... 19, 54, 58, 59, 85, 86, 98, 114, 151, 162, 171, 211, 231  
Germ tube ..... 38  
germacrene D ..... 53, 69, 70, 71, 79, 81, 85, 99, 101, 112, 126, 141, 144, 153, 155, 161, 164, 222, 236, 243, 244, 257  
glandular trichomes ..... 57, 80, 112  
Glucosinolate ..... 84  
Great Chrism ..... 260  
Greek Orthodox Chrism ..... 260  
greenhouse ..... 54, 212, 231, 238  
Guido F ..... 237  
Guttiferae ..... 244  
gypsywort ..... 67  
H<sub>2</sub>O<sub>2</sub> ..... 257  
halophytes ..... 146  
harvest ..... 62, 94, 109, 219, 238, 265  
hawthorn ..... 132  
headspace ..... 16, 90, 107, 122, 139, 161, 220, 239, 260, 261  
health ..... 15, 113, 116, 185, 200, 220, 227  
heat-sensitive compounds ..... 124  
helianthol A ..... 66  
hemolytic ..... 115, 162, 225  
HEp-2 cells ..... 60  
herbicide ..... 160  
hexadecanoic acid ..... 132, 144, 180, 206  
Himalayan range ..... 31  
histochemistry ..... 25, 80  
honeybee ..... 145, 150  
horseradish ..... 95  
hotrienol ..... 67  
Huaceae ..... 234  
human tumor cell lines ..... 153, 156  
Huyen DD ..... 236  
hydrogen peroxide ..... 257  
hydrosol ..... 245  
Hypericaceae ..... 244  
hypernociception ..... 50, 92  
hypernociceptive ..... 50  
Imaging ..... 20  
*in vitro* ..... 33, 47, 60, 65, 67, 78, 91, 107, 111, 134, 153, 156, 157, 160, 162, 172, 193, 194, 215, 227, 231, 264

- inclusion complexes..... 46  
indoles ..... 248  
Industrial by-products..... 194  
infections..... 38, 57, 65, 69, 103, 140, 193, 203, 210, 228, 230  
inhibition..... 25, 33, 42, 48, 77, 78, 100, 102, 103, 107, 114, 128, 137, 141, 149, 150, 151, 154, 156, 192, 193, 195, 209, 210, 212, 217, 225, 229, 234, 249, 251, 257, 259, 264  
inhibitory..... 25, 32, 33, 42, 45, 47, 48, 55, 69, 76, 103, 134, 140, 159, 162, 193, 194, 208, 209, 214, 221, 230, 243, 246, 251  
iNOS ..... 33, 249  
insect proof net ..... 263  
insecticidal .. 31, 110, 130, 137, 143, 181, 183, 187  
ionic liquid stationary phase ..... 182, 233  
irritant..... 19, 263  
isoalantolactone ..... 65, 240  
isobutanoate ..... 63, 105  
isopropyl N-methylantranilates..... 82  
Isothiocyanate..... 84  
isovalerate ..... 63, 105, 207  
ISSR ..... 207  
Jerusalem artichoke..... 66  
key odorants ..... 185  
Knee ..... 249  
lactones ..... 180, 187, 240, 241  
Lady Hillingdon ..... 86  
Lamiaceae..... 46, 55, 56, 67, 70, 80, 119, 120, 129, 144, 161, 172, 183, 195, 210, 211, 215, 227, 238  
larvicidal ..... 31, 115, 116, 181, 189, 247  
Lauraceae..... 226, 229, 236  
lavandin ..... 52  
leaf anatomy ..... 57  
leaf oil..... 75, 123, 124, 126, 135, 211, 213, 219, 236  
leaves..... 25, 47, 49, 52, 53, 57, 60, 61, 67, 70, 80, 82, 87, 88, 89, 93, 99, 107, 109, 110, 111, 115, 119, 121, 126, 127, 130, 131, 135, 144, 147, 152, 153, 156, 157, 159, 161, 162, 163, 164, 167, 170, 171, 172, 173, 174, 176, 179, 186, 188, 197, 198, 201, 208, 211, 213, 214, 223, 225, 226, 228, 230, 237, 243, 246, 250, 256, 259, 262  
leishmanicidal ..... 115, 162, 179  
leiva ..... 192  
lemon ..... 78, 168, 196  
Lemon balm ..... 210  
lichens..... 192  
light/dark test..... 82  
limonene..... 52, 57, 68, 69, 75, 77, 79, 85, 101, 103, 110, 114, 116, 120, 122, 125, 153, 155, 159, 160, 175, 176, 177, 184, 192, 196, 201, 209, 211, 213, 216, 218, 221, 223, 228, 236, 246, 250, 258, 260  
Limonene ..... 103, 177, 211  
limonene synthase ..... 52  
linalool..... 19, 33, 41, 46, 52, 54, 75, 87, 89, 98, 120, 121, 125, 127, 171, 184, 185, 211, 214, 231, 237, 250, 260  
Lippia ..... 39, 57, 97, 102, 140, 163, 179, 209, 219  
Lira ..... 206  
liverwort ..... 30, 241  
lovage ..... 132  
low pressure treatment ..... 113  
Malvaceae ..... 202  
MAPK..... 249  
marjoram..... 185, 238, 259  
Mass spectrometry..... 20  
mass transfer models..... 254  
maturation ..... 262  
meat..... 252  
Medicinal plants ..... 186, 199  
methyl 2-hydroxy-3-methylhexanoate ..... 90  
methyl eugenol..... 198  
methyl N-methylantranilates..... 82  
Microalgae ..... 205  
microencapsulated preparations..... 185  
microorganisms..... 17, 47, 69, 107, 178, 187, 192, 205, 226, 242  
microtubules ..... 25, 217, 229  
milk ..... 252  
mineral oil ..... 212  
mineral supplementation..... 199  
MIR..... 39, 228  
Model system ..... 168  
molluscicidal ..... 115  
monoterpenes..... 19, 25, 32, 37, 46, 52, 68, 72, 79, 80, 92, 97, 105, 106, 108, 109, 114, 115, 118, 126, 130, 133, 140, 153, 156, 160, 162, 175, 231, 233, 236, 247  
Monoterpenes..... 25, 50, 51, 118, 163  
Moroccan chamomile..... 62  
morphological ..... 30, 49, 57, 135, 161, 238  
morphology..... 57, 80, 112, 150, 202, 209  
mosquito ..... 116, 189, 247  
mosses ..... 16, 192, 241  
Multivariate statistical analysis..... 73, 81, 83, 239  
mung beans ..... 168  
mung sprouts..... 113  
mushrooms ..... 184  
Mutualism ..... 16  
Mycoplasmas..... 248  
Mycorrhiza ..... 199  
myristicin..... 100, 178, 247, 248  
Myrrh ..... 172  
Myrtaceae..... 71, 110, 111, 115, 170, 175, 216, 228, 253  
myrthenal ..... 217  
myrtle..... 75, 250  
nanjamonjagoke ..... 241  
Nard Chrism ..... 260  
native oils..... 167  
natural compounds ..... 187, 193, 217  
natural medicines ..... 31  
natural product..... 263  
Naturalness ..... 21  
Near Threatened ..... 207  
nematicidals..... 72  
nematicides ..... 183  
nematodes ..... 72, 183  
nematotoxic ..... 72  
neoplastic ..... 172  
NF-kB ..... 249  
NIR ..... 39, 228  
nitric oxide ..... 172, 257  
NMR..... 20, 63, 90, 116, 120, 123, 124, 125, 126, 135, 170, 205, 241  
*n*-nonane ..... 174, 244  
nociceptive..... 51, 118, 120, 259  
Nonphotochemical quenching ..... 37  
norisoprenoids ..... 187  
novel..... 132, 233, 235  
olfactory compounds..... 136  
optical isomer ..... 41, 245  
Optical isomer..... 41, 245  
ORAC ..... 40, 250  
orange peel..... 212  
Organic farming ..... 168  
orofacial ..... 51, 118  
Osteoarthritis ..... 249  
outer membrane protein ..... 251  
Oxalidaceae ..... 258  
packaging material ..... 131  
pain ..... 50, 51, 53, 92, 118, 120  
Pain ..... 163  
parsley ..... 247  
patchouli ..... 88  
patchoulol ..... 88  
pathogen ..... 251, 264, 265  
pathogenic microorganisms..... 76, 209, 246  
Pathogens ..... 168  
pathway ..... 19, 191, 223, 242, 249  
*p*-cymene..... 55, 70, 71, 102, 105, 120, 140, 160,

- 213, 215, 218, 219, 242, 264
- pear ..... 94
- pennyroyal ..... 254, 255
- pepper ..... 100, 159, 262
- peppermint ..... 86, 223, 233
- Peptidyl-tRNA hydrolase ..... 42
- perfumery ..... 90, 101, 127, 152, 167, 182, 204
- periaqueductal ..... 50, 53
- pest management ..... 130, 263
- pharmaceutical ..... 45, 102, 111, 193, 205, 216, 223, 226, 250
- phenylpropanoids ..... 19, 106, 247
- photoactivity ..... 117
- photosynthesis ..... 212
- Phthalates ..... 27
- Phthalic-acid esters ..... 27
- phytochemical ..... 53, 54, 81, 91, 101, 120, 133, 143, 163, 164, 190, 207, 213
- phytopathogenic fungi ..... 117
- Phytotoxic ..... 214
- phytotoxicity ..... 168, 217, 229, 231
- Pinaceae ..... 69
- pineapple ..... 192
- pinewood nematode ..... 72, 183
- Piperaceae ..... 47, 227
- Plant microstructure ..... 28
- plant protection ..... 137
- poliovirus ..... 60
- poppy ..... 168, 169
- post-harvest ..... 78, 87, 88, 173, 200, 264, 265
- postharvest diseases ..... 78
- potato ..... 168, 206
- Principal Component Analysis ..... 41, 121, 170, 185, 223, 245
- propolis ..... 22, 60, 150
- Protected Designation of Origin ..... 94, 192
- Protected Geographical Indication ..... 206
- Prototype ..... 177
- pulegone ..... 25, 109, 116, 184, 207, 223
- pumpkin ..... 194
- qPCR ..... 37
- quantification ..... 39, 51, 60, 124, 130, 165, 232
- quantitative descriptive analysis ..... 121, 239
- Raman optical activity ..... 68
- Ranunculaceae ..... 91
- rat mesenteric artery ..... 149, 151
- rats ..... 149, 151, 157
- red sacaca ..... 208
- repellent ..... 206, 263
- ripened seed capsules ..... 174
- Rocha pear ..... 94
- root ..... 59, 63, 64, 65, 95, 96, 100, 106, 108, 114, 126, 143, 153, 156, 168, 169, 186, 198, 199, 206, 214, 217, 229, 234, 240
- Rosaceae ..... 94
- rose ..... 86, 98, 128
- Rosemary ..... 76, 122, 221, 245, 255
- Rutaceae ..... 82
- sabinene ..... 19, 79, 85, 93, 105, 129, 135, 144, 148, 154, 160, 185, 201, 215, 236
- saffrole ..... 248
- Sage ..... 39
- Santalaceae ..... 182
- Santalols ..... 182
- satCMV ..... 141
- savory ..... 255, 264, 265
- sazonality ..... 133
- scent ..... 19, 129, 176, 200, 202, 241, 261
- seasonal variations ..... 108, 245
- seasons ..... 133, 238, 245
- secretory canals ..... 156
- seed germination ..... 25, 160, 214
- Seed treatment ..... 168
- seeds ..... 25, 47, 49, 59, 91, 113, 114, 148, 160, 168, 174, 194, 198, 212, 214, 231, 234, 252
- sesquiterpenes ..... 19, 29, 68, 73, 79, 97, 105, 106, 112, 115, 126, 133, 141, 144, 152, 153, 155, 162, 170, 174, 175, 178, 198, 201, 206, 208, 213, 222, 225, 233, 236, 240, 241, 248
- SFE ..... 137, 255
- shelf-life ..... 78, 265
- Shelf-life ..... 168
- Shiikuwasha ..... 220
- simulation ..... 235
- Sleeping sickness ..... 256
- snake venom ..... 225
- snowdrop ..... 90
- soil biota ..... 199
- Solanaceae ..... 18, 37, 54
- solid phase microextraction ..... 95, 128, 131, 139, 202, 220, 258
- soybean ..... 114, 212
- spastic muscle ..... 188
- spices ..... 93, 185, 200, 234, 259, 261
- spider mite ..... 143, 206
- SPME ..... 16, 95, 107, 131, 132, 139, 167, 175, 185, 200, 220, 239, 258
- $\beta$ -caryophyllene ..... 213
- stems ..... 49, 70, 80, 99, 135, 144, 153, 156, 174, 186, 198, 214, 219
- stereochemistry ..... 20, 68, 136
- strawberry ..... 48, 168
- stripping ..... 235
- stroke ..... 188
- structure elucidation ..... 20, 123
- Structure-odour relation ..... 29
- sugar beet ..... 132
- sulcatyl acetate ..... 120
- supercritical fluid extraction ..... 28, 91, 137, 254, 255
- surface response ..... 177
- survival ..... 20, 145
- Sustainable ..... 199
- sweet potato ..... 206
- sweet potato weevil ..... 206
- symbiosis ..... 199
- Synergism ..... 38
- synergy ..... 264
- synthesis ..... 29, 63, 77, 82, 84, 90, 136, 187, 191, 215, 249
- tail suspension test ..... 82
- Taste ..... 15, 146
- Tea tree ..... 221, 228
- TEAC ..... 140, 250
- Terpene biosynthesis ..... 19
- terpene synthase ..... 19
- Terpene synthases ..... 37
- terpenes ..... 19, 107, 133, 137, 177, 191, 223, 252
- terpenoids ..... 17, 20, 30, 52, 68, 80, 136, 137, 180, 236, 237, 242, 244, 261
- Therapeutic tool ..... 138
- Threatened Species ..... 207
- thyme ..... 19, 117, 127, 185, 242, 251, 255, 264
- thymol ..... 19, 55, 72, 78, 85, 97, 102, 129, 140, 160, 179, 185, 190, 203, 211, 215, 218, 219, 259, 264
- TLC ..... 56, 117, 152, 230, 234, 241
- Tomato ..... 18, 239
- toxic ..... 19, 32, 33, 72, 145, 162, 172, 225, 230, 263
- toxicity ..... 74, 105, 115, 137, 153, 157, 159, 162, 164, 189, 203, 230, 263
- TPS ..... 19
- traditional ..... 45, 47, 53, 99, 100, 132, 152, 172, 192, 194, 198, 199, 205, 255, 258
- transgenic plant ..... 52
- trypanocidal ..... 256
- tsetse flies ..... 256
- tubers ..... 66
- Umbelliferae ..... 77, 243
- vanillates ..... 77
- variability ..... 49, 62, 74, 75, 76, 79, 83, 98, 105, 108, 109, 114, 126, 144, 170, 198, 202, 205, 209, 218, 239, 250
- variation ..... 39, 71, 85, 97, 112, 140, 181, 207, 219, 260

vasorelaxant.....	149, 151
vector control .....	116, 189
vegetative organs.....	156
Verbenaceae.....	57, 140, 163, 209, 213
Vibrational circular dichroism .....	68
Vibrational spectroscopy.....	39
virus .....	60, 116, 141, 263
Volatile.....	22, 30, 31, 64, 185, 190, 200, 222, 231, 239, 255, 258, 259
Volatile chemicals .....	31
Volatile components.....	30, 222
Volatiles.....	16, 17, 18, 20, 95, 128, 129, 131, 147, 161, 184
vulvovaginal candidosis .....	138
weed .....	59, 74, 114, 180, 229, 231
weed management .....	229
wheat .....	25, 168
whitefly.....	263
wiped film reboiler.....	177
wood .....	110, 139, 183, 198, 201
Wormwood.....	105
yarrow .....	45, 96
$\alpha$ -pinene.....	33, 45, 59, 64, 66, 70, 71, 75, 79, 80, 99, 110, 126, 135, 141, 154, 155, 192, 196, 201, 216, 219, 222, 236, 243, 264
$\alpha$ -Santalol.....	29
$\beta$ -caryophyllene.....	80, 85, 105, 141, 142, 144, 198, 236
$\beta$ -cyclodextrin.....	46
$\beta$ -myrcene.....	62, 79, 105, 145, 191
$\beta$ -phellandrene.....	69, 196, 243
$\beta$ -pinene.....	59, 64, 69, 79, 99, 126, 142, 196, 222, 224
$\beta$ -secretase .....	32
$\gamma$ -terpinene .....	220



## Authors

Aazza S	140, 218
Abazi S	147
Abed L	75
Adamiec	185
Adjalian E	198
Adomou A	198
Afonso Pires SM	134, 145
Agnaniet H	234
Ahibo AC	123, 126
Ahibo CA	124
Aires A	194
Akhlaghi H	77
Al-ani I	106
Alcobia D	159
Alitonou GA	198
Almeida CA	210
Almeida EDP	46
Almeida JA	194
Almeida JRGS	51
Almeida LFR	223
Almeida	46, 51, 133, 194, 210, 217, 223, 230, 259, 284, 288
Alsina I	161
Alsina Ina	128
Alvarenga AA	226
Alvarez SL	201
Álvarez-Iglesias	231
Alves AAR	165
Alves FC	170
Alves LPL	162
Alves MF	87, 173
Alves PB	87, 88, 119, 120, 121, 171, 173, 179, 210
Alves RBN	176
Alves RS	92, 227
Alves	111, 163, 165, 170, 176, 188, 189, 227, 230, 259, 284, 288
Alviano CS	208, 210
Alviano DS	208, 209, 210
Alviano	230, 284
Amancio CA	209
Anass E	62
Andrade EHA	70, 71, 85, 99, 100, 101, 115
Andrade G	257
Andrade J	143
Andrade MA	142
Andrade	47, 70, 71, 85, 99, 100, 101, 115, 142, 143, 219, 284
André AC	170
Andressa MS	175, 258
Anguille JJ	234
Ankle A	234
Antoniolli AR	53, 118
Antunes MDC	146
Arabacý O	61
Arantes S	157, 159
Araujo AAS	46, 50, 149, 151
Araújo	164
Arenas Velásquez AM	256
Arrigoni-Blank MF	87, 88, 119, 121, 173, 179, 230, 284
Arriola-Guevara E	177
Asada S	52
Asakawa Y	30, 241, 242
Ascensão L	80
Atoji H	86
Avlessi F	198
Avoseh NO	237
Azevedo MMB	208, 209
Bahcevandziev K	129
Baldovini N	26
Ballottari M	37
Bankova V	22
Banwo ID	237
Baranska M	39
Barata AM	49
Barbizan DS	204
Barbosa P	72, 183
Barreto AS	118, 151
Barreto RSS	51, 118, 120
Barreto	51, 118, 120, 151, 163, 189, 284, 285, 288
Barroso JG	49, 62, 80, 110, 129, 140, 142, 143, 183, 184, 191, 192, 216, 218, 255
Bassi R	37
Batista LR	142
Batista RS	57
Baydar H	61
Bazzali O	135
Begnini IM	178, 248
Beleboni RO	179
Belhassen E	26
Bella Cruz A	178
Bellucci I	253
Bencsik T	251
Benevides PC	204
Benites J	196
Beric T	221
Bernardes GG	226
Bernardo-Gil	250
Bernáth J	105
Bettencourt E	49
Bezerra DP	92
Bezić N	141
Bicchi C	182
Bighelli A	75, 123, 124, 126, 135
Bikanga R	234
Bitencourt TA	179
Bizzo HR	165, 176, 208
Blagojević PD	64, 81, 96, 240
Blank A	173
Blank AF	87, 88, 119, 121, 179, 209, 210
Blázquez MA	104
Boaro CSF	223
Bobeda CRR	172, 175, 258
Boido E	253
Boira H	104, 108, 109
Bombo AB	153, 156
Bonaccorsi I	232
Bonato CM	58, 59
Bonato	58, 59, 114, 212, 284
Bongui JB	234
Bonjardim LR	51, 53, 118, 149
Borg-Karlson A-K	17
Bosilcov A	54
Bossi S	37
Böszörményi A	251
Botelho VT	257
Boti JB	123, 124, 126
Bouzabata A	75
Boylan F	82
Bozza M	172
Braga TV	107
Bragança H	201
Branquinho AR	146
Bravo F	196
Bravo LR	47
Brevard H	26
Brito AF	88
Brito RG	50, 51
Buchbauer G	139, 158
Bufalo J	223
Bufalo MC	60
Cabral C	45
Cabrera D	253
Cachet T	27
Cagliero C	182
Calvo-Irabién LM	97, 112
Camargo	225

Cammareri M.....	37	Delgado FMG.....	48
Campos-Takaki GM.....	210	Delgado M.....	116
Candeias F.....	159	Dellacassa E.....	253
Candeias MF.....	157	Dellacassa.....	225, 285
Cândido.....	188	Denić M.....	65, 67
Canhoto J.....	33	Di Dato F.....	37
Canul-Puc K.....	112	Dias C.....	193, 194
Cardoso MG.....	110, 140, 142, 143, 227	Dias CN.....	115, 162
Cardoso RSD.....	53	Dias LS.....	183
Cardoso.....	163, 188, 285	Díaz-Tielas C.....	217, 229
Carlos IZ.....	257	Djordjevic.....	180, 285
Carneiro SMP.....	115, 162	Dobravalskyte.....	166, 167
Carolino M.....	192	Dognini J.....	178, 248
Carrau F.....	253	Domingos C.....	192
Carreira LMM.....	70, 71, 85, 99, 101	Domingos VD.....	85
Carvalho EEN.....	85	Domingues FC.....	38
Carvalho FAA.....	115, 162	Đorđević A.....	244
Carvalho NCC.....	70, 71, 99, 102	Đorđević M.....	96
Casanova J.....	75, 123, 124, 126, 135	Đorđević A.....	243
Castanheira P.....	32	Đorđević MR.....	66
Castola V.....	75, 126	Đorđević ND.....	73, 74
Cavalcante MQ.....	152	Doria.....	164, 284
Cavaleiro C.....	32, 33, 45, 58, 59, 114, 117, 138, 148, 155, 203, 249, 256, 257	Duarte A.....	38
Cavaleiro.....	5, 11, 49, 231, 285	Duarte RC.....	200
Cebola.....	250, 285	Dudai N.....	25
Chaimovitch D.....	25	Dugo P.....	232, 233
Chandre F.....	263	Dunkić V.....	141
Chaves FCM.....	208	Dzib G.....	112
Chizzola R.....	93, 144	Eguaras MJ.....	134, 145
Churio MS.....	134	EHA.....	102
Cicarelli RMB.....	256	Eike R.....	234
Coelho JAP.....	254	El Amrani A.....	62
Coelho JP.....	255	Ennigrou.....	262
Cordeiro I.....	103	Espirito Santo C.....	205
Cordero C.....	182	Estevam.....	46, 189, 259
Cordova CMM.....	248	Esteves T.....	201
Coser RN.....	152	Fachin AL.....	179
Costa AS.....	121	Falconieri D.....	91
Costa DCM.....	209	Faleiro L.....	146
Costa EV.....	53, 120	Faleiro M.....	218
Costa IM.....	204	Fareleira JMNA.....	254, 255
Costa JR.....	205	Faria JMS.....	72, 216
Costa LCB.....	57	Farias Junior PA.....	179
Costa P.....	40	Fariña.....	253
Costa.....	4, 10, 40, 53, 57, 120, 121, 135, 145, 156, 164, 182, 204, 205, 209, 239, 285	Farneti.....	239, 287
Costas-Gil A.....	217, 229	Faro C.....	32
Cotoraci.....	260	Feitosa VLC.....	50
Cristescu.....	239	Felsó P.....	251
Cristino AF.....	254	Fernandes ESS.....	155
Cruz C.....	192	Fernandes RPM.....	179
Cruz EMO.....	179	Fernandes-Ferreira M.....	174, 247
Cruz MT.....	33, 45, 148, 203	Fernandez NJ.....	134
Cruz RCB.....	178	Ferrarese-Filho O.....	58
Cruz-Morais J.....	157, 159	Ferraz PC.....	152
Cujic N.....	221	Ferreira MASS.....	48, 76
Cunha PS.....	149, 151	Ferreira R.....	176
da Glória BA.....	153, 156	Ferreira S.....	38
da Silva GF.....	46	Ferrer MAB.....	160
da Silva JKR.....	85, 99, 100, 102	Ferrer-Ortega MM.....	97
da Silva.....	230	Figueiredo AC.....	49, 62, 72, 80, 94, 110, 129, 140, 142, 143, 183, 184, 191, 192, 206, 207, 215, 216, 218
Dai DN.....	236	Figueiredo RO.....	204
Damiani N.....	134	Filho OF.....	114
Dantas-Barros AM.....	107	Filippi J-J.....	26
Danthu P.....	202	Finkousova A.....	78
de Almeida LFR.....	133	Fonseca.....	164
De Armas Y.....	111	Fontes SS.....	87
de Carvalho JE.....	153, 156	Fraga BP.....	149, 151
de Figueiredo M.....	98	Franca SC.....	179
de Medeiros IA.....	151	Franco CRP.....	171
de Oliveira ASS.....	153, 156	Franco CS.....	102
de Vasconcelos MCBM.....	193, 194	Frankova A.....	4, 78, 113, 168, 169, 286
Degenhardt J.....	19	Frei F.....	133
Dekić M.....	83, 84	Freitas-Silva O.....	172
Deletre E.....	263	Fritz R.....	134
Delgado FG.....	89	Głowniak K.....	190

Gago C	146, 218	Kamkaen	246, 286
Gago G	140	Karapandzova M	69, 79
Gallucci SM	204	Karban J	137
Galoburda R	95, 131, 161	Kawaraya A	86
García-Fajardo JA	177	Kern K	58
García-Rellán D	104, 108, 109	Kern KA	114
Garibaldi A	264, 265	Kindlovits S	105
Garrido PM	145	Kiril B	205
Gaspar C	138	Kitao M	41, 245
Gaspar EMM	200, 206	Klee HJ	18
Gediga J	187	Kloucek P	78, 113, 168
Geissler M	158	Kloucek	78, 113, 168, 169, 286
Gende LB	134, 145	Koca U	91
Gładkowski W	187	Kochmańska J	190
Gliszczyńska A	187	Kocsis B	251
Głowniak K	30	Koehorst RR	199
Gochev V	158	Kozyra M	190
Godányi R	251	Krause S	19
Godjevac D	125, 221	Kremer D	141
Godoy RLO	172, 175, 258	Kruma Z	95, 128, 131, 161
Gomes AC	89	Kuchtova P	168
Gomes MS	110	Kukaniev MA	224
Gomes Neto SC	88	Kulevanova S	69, 79
Gonçalves MJ	33, 45, 148, 203	Kundakovic T	55
Gonçalves S	40	Kuriata-Adamusiak R	136
Gouvea ACSM	175	Lacerda ICA	227
Graça SFN	48	Lakusic B	55
Graña E	217, 229	Larsen MK	252
Grangeiro-Junior S	46	Laubscher CP	199
Grãos M	32	Lazareviæ J	244
Grasso	233	Lazarevic	96, 180
Grego JAB	94	Lazari D	147
Guedes AP	174	Lebibi J	234
Guerreiro MC	186	Lejonklev J	252
Guido F	237	Lemberkovics É	251
Guimarães AG	92	Leon LL	172
Gullino ML	264, 265	Leerma MJ	47, 111
Gurgel ESC	214	Liberto E	182
Gutiérrez-González P	177	Liddle P	27
Hamada Y	68	Lima AAN	46
Hanganu D	56, 122	Lima AS	129
Hanganu	260	Lima DS	162
Harren	239	Lima FWJ	107
Hasegawa T	29	Lima PSS	53, 118
Hassan AS	237	Lima RN	173
Hayashi T	41, 245	Lima	119, 163, 201, 287
Henriques MCMO	172	Llorens-Molina JA	108, 109
Hermida L	145	Lochynski S	136
Herrero JM	47, 111	Lopes JF	200, 206
Higuchi M	241	Lopes MC	45, 148, 249
Hodaj E	147	Lopes VR	49
Höferl M	158	Lopes	94, 140, 230, 287
Horváth Gy	251	López J	196
Hoskovec	169	López S	109
Huyen DD	236	Lopez-Reyes JG	264, 265
Ilic M	154	Lorenzo	225
Ishii-Iwamoto EL	58, 59, 114	Lucas FCA	70, 71, 85, 99, 101, 214
Izumi H	29, 68	Lucas SC	94
Jamal Eddine J	62	Lucca-Júnior W	50, 51
Jesus AS	119, 173	Lucchese	163
Jesus HCR	87, 171, 179	Ludwiczuk A	30
Jezler CN	57	Luz JMQ	238
Jianbin	261	Luz SFM	100
jing	261	Machado AMR	175, 258
Jirovetz L	139, 158	Maffei M	37
Jones	222, 230, 286	Magalhães NO	256, 257
Jorge E	47, 111	Magalhães PM	152
Joulain D	21	Maia JGS	71, 85, 99, 100, 101, 102, 162
Jovanović O	154, 243	Maia, JGS	70
Jovanović VS	154	Maia	70, 71, 85, 99, 100, 101, 102, 115, 162, 219, 287
Judas F	249	Makani Th	234
Julião M	146	Maluf WR	143
Júnior	46, 50, 51, 53, 92, 118, 120, 149, 163	Marchioro M	151
Kadifkova-Panovska T	69	Marghitas LAI	150
Kaftandzieva A	69	Markovic S	55
Kalinova	169	Marongiu B	91, 203
Kamatou GPP	228		

Marques MOM	133, 223, 238	Nobre BP	254, 255
Marques RM	58, 59, 114	Noma Y	242
Martin T	263	Noudogbessi JP	198
Martínez N	253	Novak J	127
Martinez	138, 225	Novo MT	116, 247
Martinez-de-Oliveira J	138	Nunes IS	200, 206
Martínez-Natarén DA	97	Nunes MJC	76
Martínez-Peñalver A	217, 229	Nunes RS	151
Martins CHG	257	Nurcahyanti ADR	106
Martins GZ	256, 257	Occhipinti A	37
Martins MBG	256	Ogunsanwo AA	237
Martins MR	157, 159	Ogunwande IA	236, 237
Masango	235, 287	Ohta	220, 288
Mashimbye MJ	211, 213	Ohya N <sup>1</sup> ; Hasegawa T	68
Mateussi	212	Olah NK	56, 122
Matias D	195	Olah	260
Matos DC	257	Oliveira ARMF	57
Matos OC	117	Oliveira CM	110, 143
Matos P	254, 255	Oliveira DEC	186
Matsumura Y	41, 245	Oliveira MA	92
Maxia A	203	Oliveira RA	57
Mbot E	234	Oliveira RC	238
McFeeters H	42	Oros	260
McFeeters RL	42	Ortuño C	104
Medina AF	59	Ouattara ZA	124
Medinilla M	47	Ozcelik B	91
Meierhenrich U	26	Pacheco S	175, 258
Meireles ENM	100	Palavra AF	254, 255
Melo JO	179	Paliæ I	244
Melo NSR	92	Palić I	243
Mendes AF	249	Pall E	150
Mendes MD	215	Palmeira-de-Oliveira R	138
Mendonça MS	214	Panagopoulos T	146
Meneghetti EK	178, 248	Parra-Tabla V	97
Menezes PP	46	Paudel P	31
Menut C	198, 202, 216, 234, 263	Paudel	31, 181
Menut	166	Paula ACCF	226
Miguel MG	140, 146, 218	Paula ACCFF	227
Milenkovic M	55	Paura L	161
Miltojevic AB	82	Pavela R	137
Miltojevic AB	90	Pedro L	192, 218
Minteguiaga	225	Pedro LG	49, 62, 80, 110, 129, 140, 142, 143, 183, 184, 191, 216
Miranda CASF	110	Pedrol	231
Mitić V	154	Pedroso L	76
Mladenović MZ	63, 64, 73	Pelissari GP	257
Moiteiro C	116, 195, 196, 201	Pereira M	157
Molnár P	251	Pereira	157, 250, 288
Mondello L	232, 233	Petersen MA	252
Monteagudo U	47, 111	Petroviæ G	244
Monteiro OS	70, 71, 99, 102	Petrović G	154
Monteiro	70, 71, 99, 102, 207	Pham VT	135
Montes AL	186	Phan KL	135
Moraes DFC	115, 162	Piccoli RH	227
Moré M	16	Pingsheng	261
Moreira FV	149, 151	Pintado CMB	48
Moreira RRD	256, 257	Pinto S	184
Moreno PRH	103	Piras A	91, 203
Morère A	234	Porcedda S	91
Morgovan C	122	Porrini MP	145
Morgovan	260	Portella RO	133
Mortensen G	252	Porto ALM	171
Mos L	122	Porto C	204
Mota L	80	Poudel	181
Mota M	72	Póvoa O	207
Mota P	205	Prado DS	50, 51, 53
Muizniece-Brasava S	131	Prado	50, 51, 53, 163
Munteanu MF	56	Prata APN	53
Murakami H	41, 245	Prelle A	264
Nakagawa M	245	Queiroga CL	152
Nascimento SS	46	Queiroz JA	38, 138
Natividade MMP	227	Queiroz VT	110
Nelson DL	107, 110	Quintans JS	118, 120
Nepomuceno ALA	257	Quintans JSS	50, 51, 53, 92
Neto AP	204	Quintans Júnior LJ	118
Nguyen TH	135	Quintans	46, 50, 51, 53, 92, 118, 120, 149, 163, 164, 288
Niculæ M	150		
Nii K	241		

Quintans-Júnior LJ	46, 50, 51, 53, 120, 149
Quitans-Júnior LJ	92
Radoias G	54
Radulović N	65, 67, 81, 83, 84
Radulovic N	96
Radulović NS	63, 64, 66, 73, 90
Radulovic NS	74, 77, 82
Radulović	240
Raggi L	103
Ragonese C	233
Raguso RA	16
Rakouth B	202
Ramis G	111
Ramos AR	100
Ramos GR	47
Randelović V	83
Randjelovic PJ	82
Rasoamanana E	202
Razanamaro O	202
Rebelo RA	178, 248
Rehder VLG	153, 156
Reigosa MJ	217, 229
Reigosa	217, 229, 231
Reis B	58, 114
Reis	58, 114, 184, 212, 254, 255, 288
Rellán DG	160
Resende JMV	142
Resende O	186
Rezende CM	165, 170, 176
Riazi Ali	218
Ribeiro AF	71, 85, 101, 102
Ribeiro EAN	151
Ribeiro	71, 85, 101, 102, 151, 219
Ricciardi	225
Ricciardi	225
Rijo P	195
Ríos D	196
Rivero CF	109
Rocha DK	116
Rodilla JM	89
Rodrigues AG	138
Rodrigues AM	191
Rodrigues AS	165
Rodrigues DF	256
Rodrigues KAF	115, 162
Rodrigues LMA	142
Rodrigues LS	207
Rodrigues	160, 188, 189, 285, 288
Rodríguez MV	195
Rojas M	196
Rojas R	201
Romano A	40
Romano	40, 250
Roma-Rodrigues C	191
Rosa A	91
Rosa EA	193, 194
Rosas EC	172
Rossini C	145
Rostoks N	128
Rubiolo P	182
Rufino AT	249
Ruiz ALTG	153, 156
Rušćić M	141
Russo M	232
Saavedra MJ	193, 194
Sá-Correia I	191
Sajfrtova M	137
Sakamoto A	68
Sakurai K	86
Salamone A	160
Sales J F	186
Salgueiro L	32, 33, 45, 58, 59, 114, 117, 138, 148, 155, 203, 249, 256, 257
Salgueiro	5, 9, 231, 288
Sanchez J	216
Sánchez I	196
Sánchez-Moreiras AM	217, 229
Sancho MV	160
Sandasi M	39, 228
Sandru CD	150
Santana CJ	200, 206
Santana HCD	176
Santana MT	51
Santo	110, 116, 189, 204
Santos AG	257
Santos DA	88, 120
Santos JUM	214
Santos LE	256, 257
Santos MI	76
Santos MRV	50, 51, 118, 120, 149, 151
Santos P	194
Santos PL	50
Santos PM	191
Santos PS	88
Santos	50, 51, 57, 76, 88, 118, 120, 121, 145, 149, 151, 163, 170, 188, 189, 191, 194, 214, 226, 227, 256, 257, 259, 287, 289
Santos-Junior HM	170
Sárosi S	105
Sartoratto A	152
Sasaki	220
Satyal P	31, 224
Satyal	4, 11, 31, 181, 289
Saucedo Y	47, 111
Savikin K	221
Scher R	179
Schimmel J	19
Schmiderer C	127
Schmidt E	139, 158
Sciarrone D	233
Sela F	79
Sena I	216
Serafini MR	46, 259
Setzer WN	31, 42, 224
Setzer	31, 42, 181, 222
Sforcin JM	60
Sgorbini B	182
Sgrot F	248
Sharopov FS	224
Silva DC	57
Silva EA	226
Silva F	38
Silva FG	186
Silva GH	171
Silva LA	48, 89
Silva LF	143
Silva MA	176
Silva MAAP	121
Silva SM	238
Silva TB	53
Silva	4, 10, 38, 46, 48, 53, 57, 85, 89, 99, 100, 102, 111, 114, 115, 121, 143, 162, 164, 171, 172, 192, 212, 227, 285, 286, 289
Simó EF	47, 111
Simões MF	195
Simonović S	243
Siqueira EP	107
S-Manzanera MCA	127
Smid J	78, 113, 168
Smid	4, 78, 113, 168, 169, 289
Soares J	238
Sobrinho AFS	172
Socaci S	56, 122
Socaciu	260
Son LC	236
Sousa DP	50
Sousa I	76
Sousa MJ	184
Sousa RMOF	247
Sousa RMOFS	174
Souto	163
Souza APS	214
Souza JA	140, 142
Souza140	142, 163, 164, 189, 258, 259, 285, 289

Sovova H	28, 137	Vignale B	253
Spadaro D	264, 265	Viljoen AM	39, 228
Sparinska A	128	Villavicencio ALCH	200
Spinu M	150	Vlazny	169
Sponza S	93, 127	Vuko E	141
Sporer F	106	Wanner J	139, 158
Stan L	150	Wattanachaiyingcharoen	246, 290
Stankovic S	221	Wawrzeńczyk C	187
Stanković S	81	Wedelia	103
Stefkov G	69, 79	Weisbjerg MR	252
Stevens CM	98	Widmer V	234
Stevic T	221	Wink M	106
Stojanoviæ G	244	Winkel C	15
Stojanović G	154, 243	Withwell MI	98
Stojanovic NM	77, 82	Wojtowicz E	132
Stojanovic	4, 77, 82, 180, 289	Wojtowicz	132, 185, 290
Stojanović-Radić Z	65, 67, 83, 84	Woltering	239
Strub D	136	Woods	181
Suárez-Jacobo A	177	Xavier-Filh	188
Sukilovic B	55	Yahada	220
Svatoš A	20	Yamada H	29
Svensson G	16	Yamamoto	220, 290
Szczepanik M	130	Yang	197
Szumny A	130	Yapi TA	123, 126
Tadtong	246, 289	Ye	197
Takátsy A	251	Yoshimoto	220
Talou T	95	Youcefi F	218
Talou	95, 166, 167, 289	Young MCM	103
Tankeu YS	228	Zacchino SA	195, 201
Tefiani C	218	Zámboiné-Németh E	105
Teixeira ML	140, 143	Zawirska-Wojtasiak R	132
Termolino P	37	Zawirska-Wojtasiak	132, 185, 290
Thai TH	236	Zawitowska B	130
Thang TD	236	Zdunic G	125
Theka T	211, 213	Zenghui	261
Thompson RM	42, 224	Zeni ALB	248
Tiamiyu AZ	237	Zhang	197, 263, 290
Tieman DM	18	Zhao	17, 166, 167, 290
Tinoco MT	89, 159	Zimmermann AC	178
Tizziani T	248	Zlatković B	243
Tofana	260	Zlatkovic DB	77
Tokita N	68	Zlatkovic	77, 180, 290
Toma CC	122	Zoghbi MGB	214
Toma	260	Zuzarte M	33, 148, 203
Tomi F	75, 123, 124, 126, 135		
Tomi K	41, 245		
Tomson L	95		
Torquillo HS	172, 175, 258		
Torre G	232		
Torres I	196		
Torres	177, 196, 225, 289		
Torres-Martínez LG	177		
Tortajada HB	160		
Tovar IM	127		
Trajkovska-Dokik E	69		
Tran HT	135		
Tran MH	135		
Trindade H	94, 129, 215, 216		
Tsuro M	52		
Turiel NA	99, 101		
Turowska-Tyrk I	136		
Utsunomiya N	245		
Vacas S	108, 109		
Vajs V	125		
Valente F	76		
Valente J	148		
Van Den Berger C	207		
van der Heyden Y	47		
Vander Heyden I	111		
Venskutonis	166		
Verdeguer M	104		
Verissimo CSC	76		
Vermaak I	228		
Vicente NF	226		
Vicet L	111		
Videira R	32		
Vieira P	183		



## Participants List







## Participants List

Surname	Name	Institution	Country	E-mail
Abdelaziz	Elamrani	Faculty of Science Ain chock	Morocco	aelamrani77@yahoo.fr
Aguiar Andrade	Eloisa Helena	Universidade Federal do Pará	Brazil	eloisandrade@ufpa.br
Almeida	Catia	Instituto de Microbiologia Paulo de Góes /Universidade Federal do Rio de Janeiro	Brazil	catiamancio@yahoo.com.br
Alvarez	Lorena	University of Vigo	Spain	lorena.alvarez@uvigo.es
Alves	Jorge	Ervitas Catitas	Portugal	j.p.alves@zonmail.pt
Alviano	Daniela	Instituto de Microbiologia Paulo de Góes /Universidade Federal do Rio de Janeiro	Brazil	danielalviano@micro.ufrj.br
Antunes	Maria Dulce	Universidade do Algarve	Portugal	mantunes@ualg.pt
Appezato-da-Glória	Beatriz	ESALQ, University of São Paulo	Brazil	bagloria@usp.br
Arrigoni-Blank	Maria de Fátima	Federal University of Sergipe	Brazil	fatima.blank@gmail.com
Asada	Satoshi	Meijo University	Japan	113561001@ccalumni.meijo-u.ac.jp
Asakawa	Yoshinori	Tokushima Bunri University	Japan	asakawa@ph.bunri-u.ac.jp
Ascensão	Lia	Centro de Biotecnologia Vegetal	Portugal	lia.ascensao@fc.ul.pt
Asma	Ennigrou EP Ennigrou	FST	Tunisia	asma.ennigrou@gmail.com
Azevedo	Mariana	Instituto de Microbiologia Paulo de Góes /Universidade Federal do Rio de Janeiro	Brazil	marimbarros@gmail.com
Azevedo Doria	Grace Anne		Brazil	gracedoria@hotmail.com
Azevedo Turiel	Nathalie	UFPA	Brazil	nathalie@ufpa.br
Bahcevandziev	Kiril	CERNAS/ESAC	Portugal	kiril@esac.pt
Bankova	Vassya	Bulgarian Academy of Sciences	Bulgaria	bankova@prgchm.bas.bg
Barbosa	Pedro	Évora University	Portugal	pmbarbosa@ymail.com
Barreto Alves	Péricles	Federal University of Sergipe	Brazil	periclesbalves@gmail.com
Barroso	José G.	Faculdade de Ciências da Universidade de Lisboa	Portugal	jgbarroso@fc.ul.pt
Baydar	Hasan	Süleyman Demirel University Agricultural Faculty	Turkey	hasanbaydar@sdu.edu.tr
Bélanger	André	Horticulture R&D Centre	Canada	andre.belanger@agr.gc.ca
Bernáth	Jenő	Corvinus University of Budapest	Hungary	jeno.bernath@uni-corvinus.hu
Bicchi	Carlo	University of Turin	Italy	carlo.bicchi@unito.it
Bighelli	Ange	University of Corsica	France	ange.bighelli@univ-corse.fr
Bizzo	Humberto	Embrapa Food Technology	Brazil	bizzo@ctaa.embrapa.br
Blagojevic	Polina	Faculty of Science and Mathematics, University of Nis	Serbia	blagojevicpolina@gmail.com
Blank	Arie Fitzgerald	Universidade Federal de Sergipe	Brazil	arie.blank@gmail.com
Bobeda	Cláudio	Instituto Federal de Educação, Ciência e Tecnologia do Rio de Janeiro – Unidade de Nilópolis	Brazil	claudio.bobeda@ifrrj.edu.br
Bonato	Carlos Moacir	Universidade Estadual de	Brazil	cmbonato@hotmail.com

Surname	Name	Institution	Country	E-mail
		Maringá		
Borg-Karlson	Anna-Karin	KTH, Royal Institute of Technology	Sweden	akbk@kth.se
Bosilcov	Alin	Brüder Unterweger GmbH	Austria	alin.bosilcov@unterweger-oils.com
Bouzabata	Amel	CNRS-UMR 6134	France	amelbouz2009@gmail.com
Buchbauer	Gerhard	University of Vienna	Austria	gerhard.buchbauer@univie.ac.at
Bufalo	Michelle Cristiane	Instituto de Biociencias - UNESP	Brazil	michelle_bufalo@yahoo.com.br
Búfalo	Jennifer	Instituto de Biociências - UNESP	Brazil	jenniferbufalo@yahoo.com.br
Cabral	Célia	Faculdade de Farmácia /Universidade de Coimbra	Portugal	cmdscabral@gmail.com
Cachet	Thierry	IOFI	Belgium	tcachet@iofiorg.org
Calvo	Luz Maria	Centro de Investigación Científica de Yucatan	Mexico	lumali@cicy.mx
Cardoso	Maria das Graças	Universidade Federal de Lavras	Brazil	mcardoso@dqj.ufla.br
Carneiro	Ana Paula		Portugal	apccarneiro@sapo.pt
Cavaleiro	Carlos	Faculdade de Farmácia, Universidade de Coimbra	Portugal	cavaleir@ff.uc.pt
Cebola	Maria-João	Instituto Superior Técnico	Portugal	m.joao.cebola@gmail.com
Celine	Delasalle	Albert Vieille SAS	France	cdelasalle@albertvieille.com
César Duarte	Renato	IPEN-CNEN/SP - USP & FCT-UNL	Portugal	renatoduarte@yahoo.com.br
Chaintreau	Alain	Firmenich SA	Switzerland	rosella.gill@firmenich.ch
Chizzola	Remigius	Veterinary Medicine University Vienna	Austria	Remigius.Chizzola@vetmeduni.ac.at
Choukri	Tefiani	Tlemcen university	Algeria	choukri13@gmail.com
Clemente Filha Ferreira de Paula	Ana Cardoso	Instituto Federal de Educação Ciencia e Tecnologia - IFMG/ Bambuí	Brazil	ana.paula@ifmg.edu.br
Clery	Robin	Givaudan	Switzerland	robin.clery@givaudan.com
Coelho	José	IST	Portugal	jcoelho@deq.isel.ipl.pt
Colaço Rodrigues	Ana Margarida	Faculdade de Ciências da Universidade de Lisboa	Portugal	amargaridacrodrigues@gmail.com
Costa	Larissa	Universidade Estadual de Santa Cruz	Brazil	larissacbc@hotmail.com
Costa	Patrícia	University of Algarve	Portugal	a26166@ualg.pt
Costas-Gil	Aitana	University of Vigo	Spain	aitana@uvigo.es
Da Silva	Joyce Kelly	Universidade Federal do Pará	Brazil	joycekellys@ufpa.br
de Figueiredo	Maria		South Africa	Maria.Figueiredo@kzndae.gov.za
de Souza Siqueira Barreto	Rosana	Universidade Federal de Sergipe	Brazil	rosanasfisis@hotmail.com
Degenhardt	Jörg	Martin Luther University Halle	Germany	joerg.degenhardt@pharmazie.uni-halle.de
Dekić	Milan	State University of Novi Pazar	Serbia	mdekic@np.ac.rs
Delgado	Fernanda	CERNAS/ IPCB-ESACB	Portugal	fdelgado@ipcb.pt
Dellacassa	Eduardo	Universidad de la República del Uruguay	Uruguay	edellac@fq.edu.uy
Denić	Marija	Faculty of Science and Mathematics, University of Niš	Serbia	denijum@yahoo.com
Dias	Clarice Noleto	Universidade Federal do Maranhão	Brazil	clia_noleto@yahoo.com.br
Djordjevic	Nevenka	Faculty of Science and Mathematics, University of Nis	Serbia	nevenka_83@yahoo.com
Đorđević	Miljana	Faculty of Science and Mathematics, University of Niš	Serbia	miljanadjordjevic11@gmail.com
Dudai	Nativ	Agricultural Research Organization	Israel	nativdud@gmail.com
Dugo	Paola	University of Messina	Italy	pdugo@unime.it

Surname	Name	Institution	Country	E-mail
Dunkić	Valerija	Faculty of Science	Croatia	dunkic@pmfst.hr
Emilie	Deletre	Cirad	France	emilie.deletre@cirad.fr
Erich	Schmidt	University of Vienna	Austria	info@artandfragrance.de
Eva	benOuaghran	Givaudan Deutschland GmbH	Germany	sandra.wunder@givaudan.com
Falconieri	Daniilo	Universita' Di Cagliari	Italy	daniilo.falconieri@tiscali.it
Faria	Jorge	Faculdade de Ciências da Universidade de Lisboa	Portugal	jmfaria@fc.ul.pt
Farneti	Brian	Wageningen UR	Italy	brian.farneti@gmail.com
Fernandes	Carlos	Planalto Dourado, Lda	Portugal	planlato.dourado@yahoo.fr
Ferreira-Maria	Madalena		Portugal	madalenam@sapo.pt
Figueiredo	Ana Cristina da Silva	Faculdade de Ciências da Universidade de Lisboa	Portugal	acsf@fc.ul.pt
Figueiredo	Roseane Oliveira de	NATURA	Brazil	roseanefigueiredo@natura.net
Frankova	Adela	Czech University of Life Sciences	Czech Republic	frankovaa@af.czu.cz
Franz	Chlodwig	Institute for Applied Botany and Pharmacognosy	Austria	chlodwig.franz@vetmeduni.ac.at
French	Lee	Pepsico R&D	United States	lee.french@pepsico.com
Frischke	Karen	Paul Kaders GmbH	Germany	karen.frischke@kaders.de
Galoburda	Ruta	Latvia University of Agriculture	Latvia	ruta.galoburda@llu.lv
García Rellán	David	Instituto Agroforestal Mediterráneo. Universidad Politécnica de Valencia	Spain	dagarrel@upvnet.upv.es
Gerhard	Full	Drom Fragrances	Germany	gfu@drom.com
Gliszczynska	Anna Maria	Wroclaw University of Environmental and Life Sciences	Poland	anna.gliszczynska@wp.pl
Gonçalves	Sandra	University of Algarve	Portugal	smgoncalves@ualg.pt
Gouveia	Hélia		Portugal	hrmg@live.com.pt
Guimarães Silva	Fabiano	IF Goiano	Brazil	fabianocefetrv@yahoo.com.br
Hasegawa	Toshio	Saitama University	Japan	toshihas@mail.saitama-u.ac.jp
Helene	Mercat	L OREAL	France	hmercat@rd.loreal.com
Hodaj	Entela	Agricultural University of Tirana	Albania	entelahodaj@hotmail.com
Horváth	Györgyi	University of Pécs	Hungary	gyorgyi.horvath@aok.pte.hu
Huguette	Agnaniet	Université des Sciences et Techniques de Masuku (USTM)	Gabon	ahuguette2001@yahoo.fr
Ilic	Marija	Faculty of Science and Mathematics, University of Nis	Serbia	marija.fertico@gmail.com
Jan	Demyttenaere	European Flavour Association (EFFA)	Belgium	jdemyttenaere@effa.eu
Jean-Jacques	Filippi	Université de Nice	France	Jean-Jacques.Filippi@unice.fr
Jones	Chelsea	University of Alabama in Huntsville	United States	cdj0006@uah.edu
Joulain	Daniel	SCBZ Conseil	France	dajoulain@wanadoo.fr
Kamatou	Guy	Tshwane University of Technology	South Africa	kamatoug@tut.ac.za
Kamkaen	Narisa	Suan Dusit Rajabhat Universtiy	Thailand	nkamkaen@gmail.com
Karapandzova	Marija	Faculty of Pharmacy	Macedonia	marijakarapandzova@yahoo.com
Karl-Heinz	Kubeczka	University of Wuerzburg	Germany	kubeczka@t-online.de
Karlsen	Jan	University of Oslo	Norway	jan.karlsen@farmasi.uio.no
Kitao	Makiko	Kinki University	Japan	1133650004n@nara.kindai.ac.jp
Kloucek	Pavel	CULS Prague	Czech Republic	kloucek@af.czu.cz
Koehorst	Robin	Cape Peninsula University of Technology	South Africa	koehorstr@cput.ac.za
Kozyra	Malgorzata	Medical University of Lublin	Poland	mkozyra@pharmacognosy.org
Kruma	Zanda	Latvia University of Agriculture	Latvia	zanda.kruma@llu.lv

Surname	Name	Institution	Country	E-mail
Kuriata-Adamusiak	Renata	Wroclaw University of Technology, Faculty of Chemistry	Poland	renata.kuriata@pwr.wroc.pl
Lambert	Ronald	Cranfield University	United Kingdom	rjwlambert@cranfield.ac.uk
Lazarević	Jelena	University of Niš, Faculty of Medical Sciences	Serbia	jelena217@yahoo.com
Lejonklev	Johan	Aarhus university	Denmark	JohanS.Lejonklev@agrsci.dk
Leng	Pingsheng	Beijing University of Agriculture	China	lengpsh@tom.com
Lima	Ana Sofia Borges	Faculdade de Ciências da Universidade de Lisboa	Portugal	aslima@fc.ul.pt
Llorens-Molina	Juan Antonio	Universidad Politécnica de Valencia	Spain	juallom2@qim.upv.es
Lochynski	Stanislaw	Wroclaw College of Physiotherapy	Poland	stanislaw.lochynski@pwr.wroc.pl
Lopes	Violeta Rolim	INRB, I.P.	Portugal	violeta.lopes@inrb.pt
Lopez-Reyes	Jorge Giovanni	University of Torino	Italy	giovanny.lopez@unito.it
Lucas	Flávia	Universidade do Estado do Pará	Brazil	copaldoc@yahoo.com.br
Lucas	Susana	Faculdade de Ciências da Universidade de Lisboa	Portugal	susanalucas1@gmail.com
Ludwiczuk	Agnieszka	Medical University of Lublin	Poland	aludwiczuk@pharmacognosy.org
Maia	José Guilherme	Universidade Federal do Pará	Brazil	gmaia@ufpa.br
Marchant	Melanie	IOFI	Belgium	mmarchant@iofiorg.org
Markovic	Stevan	Faculty of Pharmacy, Belgrade	Serbia	brankasukilovic@yahoo.com
Marongiu	Bruno	Universita' Di Cagliari	Italy	maronb@unica.it
Martínez-Peñalver Mas	Ana	University of Vigo	Spain	anampe3@uvigo.es
Martins	M. Rosário	Universidade de Évora	Portugal	mrm@uevora.pt
Marx Young	Maria Cláudia	Instituto de Botânica	Brazil	mcmyoung@yahoo.com.br
Masango	Phinn	Self	Austria	mutape2002@yahoo.co.uk
Matos	Olívia	INRB-Instituto Nacional de Recursos Biológicos	Portugal	oliviamatos@sapo.pt
Mendes	Marta D.	Faculdade de Ciências da Universidade de Lisboa	Portugal	mdmendes@fc.ul.pt
Méndez Tovar	Inés	ITACYL	Spain	mentovin@itacyl.es
Mendoza	Beatriz	Faculdade de Ciências da Universidade de Lisboa	Portugal	bia_mendoza@hotmail.com
Menut	Chantal	Faculté de Pharmacie	France	chantal.menut@univ-montp2.fr
Miltojević	Ana	the Faculty of Science and Mathematics	Serbia	anamiltojevic@yahoo.com
Mladenović	Marko	Faculty of Science and Mathematics	Serbia	markohem87@gmail.com
Moiteiro	Cristina	FCUL	Portugal	cmmoiteiro@fc.ul.pt
Moreno	Paulo Roberto H.	Instituto de Química - USP	Brazil	prmoreno@iq.usp.br
Mota	Luisa	FCUL	Portugal	lbmota@gmail.com
Munteanu	Melania Florina	West University Vasile Goldis Arad	Romania	anaionescuro@yahoo.com
Mur	Leon	Centre of Expertise for Plant compounds	Netherlands	leon.mur@plantenstoffen.nl
Nelson	David	Universidade Federal de Minas Gerais	Brazil	dleenelson@gmail.com
Németh-Zámboriné	Éva	Corvinus University of Budapest	Hungary	eva.nemeth@uni-corvinus.hu
Neves	Pedro		Portugal	poisne@hotmail.com
Nicolas	Baldovini	Université de Nice	France	baldovin@unice.fr
Niculae	Mihaela	University of Agricultural Sciences and Veterinary Medicine Cluj-Napoca	Romania	nicmisi23@yahoo.com
Nii	Kaeko		Japan	niiik@ph.bunri-u.ac.jp
Nunes	Inês Santos	Faculdade de Ciências e	Portugal	inez.sn@gmail.com

Surname	Name	Institution	Country	E-mail
		Tecnologia da Universidade Nova de Lisboa		
Nurcahyanti	Agustina	Institute of Pharmacy and Molecular Biotechnology, University of Heidelberg	Germany	agustina.nurcahyanti@gmx.de
Occhipinti	Andrea	University of Torino (Italy)	Italy	andrea.occhipinti@unito.it
Ogunwande	Isiaka Ajani	Lagos State University, Ojo	Nigeria	isiaka.ogunwande@lasunigeria.org
Ohta	Hideaki	Nakamura Gakuen University	Japan	hohta@nakamura-u.ac.jp
Olah	Neli Kinga	Vasile Goldis Western University, Arad	Romania	olahdr@aol.com
Oliveira Ferreira de Sousa	Rose Marie	Universidade do Minho, Escola de Ciências	Portugal	rosemarie@bio.uminho.pt
Pedro	Luis	Faculdade de Ciências da Universidade de Lisboa	Portugal	lmpedro@fc.ul.pt
Pereira	Paula	ULHT	Portugal	paulapereira1@sapo.pt
Philippe	Marruedo	L'OREAL	France	pmarruedo@rd.loreal.com
Pinto	Eduarda	Faculdade de Ciências da Universidade de Lisboa	Portugal	eduarda.1992@hotmail.com
Pinto	Sara	Instituto Politécnico de Bragança	Portugal	sarapinto109@hotmail.com
Pires	Sância	CIMO-ESAB-IPB	Portugal	spires@ipb.pt
Pirola	Camillo	Milestone Srl	Italy	marketing@milestonesrl.com
Proença da Cunha	António	University of Coimbra	Portugal	pdacunha@ci.uc.pt
Protzen	Maren	Paul Kaders GmbH	Germany	maren.protzen@kaders.de
Queiroga	Carmen Lucia	CPQBA-UNICAMP	Brazil	queiroga@cpqba.unicamp.br
Queiroz Luz	José Magno	Federal University of Uberlandia	Brazil	jmagno@umuarama.ufu.br
Quintans	Jullyana	Federal University of Sergipe	Brazil	jullyanas@yahoo.com.br
Quintans	Lucindo Quintans	Federal University of Sergipe	Brazil	lucindojr@gmail.com
Radoias	Georges	Brüder Unterweger GmbH	Austria	georges.radoias@unterweger-oils.com
Radulovic	Niko	Faculty of Science and Mathematics, University of Nis	Serbia	nikoradulovic@yahoo.com
Raggi	Ludmila	Instituto de Botânica	Brazil	ludraggi@gmail.com
Raguso	Robert	Cornell University	United States	rar229@cornell.edu
Rebelo	Ricardo	Universidade Regional de Blumenau	Brazil	ricardorebelo@furb.br
Rehder	Vera Lucia Garcia	CPQBA-UNICAMP	Brazil	rehder@cpqba.unicamp.br
Reis	Bruno	Universidade Estadual de Maringá	Brazil	brunoblew@hotmail.com
Rezende	Claudia	Federal University of Rio de Janeiro	Brazil	crezende@iq.ufrj.br
Rodrigues	Leandra			liarodrigues@isa.utl.pt
Rodrigues	Sheyla Alves	Universidade Tiradentes	Brazil	shelrodriques@hotmail.com
Rolim de Almeida	Luiz Fernando	Univ. Estadual Paulista/UNESP	Brazil	luizfernando@ibb.unesp.br
Rubiolo	Patrizia	University of Turin	Italy	patrizia.rubiolo@unito.it
Rufino	Ana	Centre of Neurosciences and Cell Biology	Portugal	ana.t.rufino@gmail.com
Russo	Marina	University of Messina	Italy	mrusso@unime.it
Sajfrtova	Marie	Institute of Chemical Process Fundamentals of the ASCR, v. v. i.	Czech Republic	sajfrtova@icpf.cas.cz
Sakurai	Kazutoshi	Takasago International Corporation	Japan	kazutoshi_sakurai@takasago.com
Sales Barreto	André	Federal University of Sergipe	Brazil	asbfisio@hotmail.com
Salgueiro	Lígia	Faculdade de Farmácia, Universidade de Coimbra	Portugal	ligia@ff.uc.pt

Surname	Name	Institution	Country	E-mail
Sandasi	Maxleene	Tshwane University of Technology	South Africa	sandasim@tut.ac.za
Santos	Deisylaine Maria	Universidade Federal de Sergipe	Brazil	deisysantos@hotmail.com
Santos	Isabel	Instituto Superior de Agronomia	Portugal	missantos@gmail.com
Santos	Marcio	Universidade Federal de Sergipe	Brazil	marciorvsantos@bol.com.br
Satyal	Prabodh	University of Alabama in Huntsville	United States	ps0013@uah.edu
Saucedo Hernández	Yanelis		Cuba	ysaucedo@uclv.edu.cu
Savikin	Katarina		Serbia	ksavikin@mocbilja.rs
Scheffer	Hans		The Netherlands	jjc.scheffer@hetnet.nl
Scher	Ricardo	Universidade Federal de Sergipe	Brazil	scher@ufs.br
Schilcher	Heinz	Formerly: Free University Berlin	Germany	schilcher_h@hotmail.com
Sena	Ines	Faculdade de Ciências da Universidade de Lisboa	Portugal	icsena@fc.ul.pt
Serafini	Mairim Russo		Brazil	maiserafini@hotmail.com
Silva	Filomena	University of Beira Interior	Portugal	filomena@fcsaude.ubi.pt
Silva	Lúcia	UBI	Portugal	mlas@ubi.pt
Simões	M. Fátima	Faculdade de Farmácia, Universidade de Lisboa	Portugal	fasimoes@ff.ul.pt
Smid	Kuba	Czech University of Life Sciences	Czech Republic	smidj@af.czu.cz
Sousa	Américo Azevedo	Universidade Federal de Sergipe	Brazil	americosousa_bio@yahoo.com.br
Souza	Josefina Aparecida	Federal University of Lavras - UFLA	Brazil	josefina@dqi.ufla.br
Sovova	Helena	Institute of Chemical Process Fundamentals	Czech Republic	sovova@icpf.cas.cz
Sponza	Silvia	University of Veterinary Medicine	Austria	silvia.sponza@vetmeduni.ac.at
Stojanoviæ	Gordana	Faculty of Science and Mathematics, University of Niš	Serbia	gocast@pmf.ni.ac.rs
Stojanovic	Nikola	Faculty of Medicine, University of Nis	Serbia	nikola.st90@yahoo.com
Sukilovic	Branka	Faculty of Pharmacy, Belgrade	Serbia	brankasukilovic@yahoo.com
Svatos	Ales	MPI for Chemical Ecology	Germany	svatos@ice.mpg.de
Tadtong	Sarin	Faculty of Pharmacy, Srinakharinwirot University	Thailand	sarin@swu.ac.th
Theka	Takalani	University of Cape Town	South Africa	takalani.theka@gmail.com
Thierry	Talou	Université de Toulouse, INP-ENESIACET	France	talou@cict.fr
Thompson	Rachel	University of Alabama in Huntsville	United States	rmt0002@uah.edu
Tieman	Denise	University of Florida	United States	dtieman@ufl.edu
Tolentino	Vanessa	Faculdade de Ciências da Universidade de Lisboa	Portugal	vanessa.s.tolentino@hotmail.com
Toma	Claudia-Crina	Western University "Vasile Goldis" of Arad	Romania	claudiatoma2004@yahoo.com
Tomi	Félix	University of Corsica	France	felix.tomi@univ-corse.fr
Tomi	Kenichi	Kinki University	Japan	tomi@nara.kindai.ac.jp
Tomson	Lolita	Latvia University of Agriculture	Latvia	lolita.tomson@llu.lv
Torcato	Cristina		Portugal	angel_cris1993@hotmail.com
Torquillo	Helena	IFRJ/Campus Nilópolis	Brazil	hpal@ig.com.br
Torres-Martínez	Lilia Guadalupe	Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de Jalisco, A.C. (CIATEJ)	Mexico	lilia_gtm@hotmail.com
Trindade	Helena	Faculdade de Ciências da Universidade de Lisboa	Portugal	htrindade@fc.ul.pt

Surname	Name	Institution	Country	E-mail
Tsuro	Masato	Meijo University	Japan	tsuro@meijo-u.ac.jp
Valente	Júlia	Centro de Estudos Farmacêuticos - Faculdade de Farmácia da Universidade de Coimbra	Portugal	jurita8@yahoo.com.br
Vasconcelos	Maria do Carmo		Portugal	mcarmov@utad.pt
Verdeguer Sancho	Mercedes	Instituto Agroforestal Mediterráneo Universidad Politécnica de Valencia	Spain	merversa@doctor.upv.es
Videira	Rita	Faculty of Pharmacy, University of Coimbra	Portugal	cavaleir@ff.uc.pt
Viljoen	Alvaro	Tshwane University of Technology	South Africa	ViljoenAM@tut.ac.za
Wanner	Juergen	Kurt Kitzing GmbH	Germany	juergen.wanner@kurtkitzing.de
Watthanachaiyingcharoen	Rith	Faculty of Pharmacy, Srinakharinwirot University	Thailand	stadtong@yahoo.com
Winkel	Chris	Givaudan	Netherlands	chris.winkel@givaudan.com
Wojtowicz	Elzbieta	Institute of Agricultural and Food Biotechnology	Poland	ewojt@man.poznan.pl
Wolfgang	Tigges	Givaudan Deutschland GmbH	Germany	sandra.wunder@givaudan.com
Yamamoto	Kenta	Nakamura Gakuen University	Japan	s10d101@nakamura-u.ac.jp
Yoshiaki	Noma	Tokushima Bunri University	Japan	ynoma@tokushima.bunri-u.ac.jp
Yoshinori	Asakawa	Tokushima Bunri University	Japan	asakawa@ph.bunri-u.ac.jp
Zawirska-Wojtasiak	Renata	Poznan University of Life Sciences	Poland	renazaw@up.poznan.pl
Zawitowska	Beata	Nicolaus Copernicus University	Poland	beatamalina87@wp.pl
Zdunic	Gordana		Serbia	gzdunic@mocbilja.rs
Zhang	Xianzhong	Zhejiang University	China	auhxyz@126.com
Zhao	Tianming	Institut National Polytechnique de Toulouse	France	steven1214.zhao@gmail.com
Zlatkovic	Dragan	Faculty of Science and Mathematics	Serbia	dragan.zlatkovic@gmail.com
Zuzarte	Mónica		Portugal	mzuzarte@hotmail.com





## Sponsors

The 43rd International Symposium on Essential Oils was generously sponsored by:



CBV | Centro de Biotecnologia Vegetal  
PBRG | Plant Biotechnology Research Group



INSTITUTE FOR BIOTECHNOLOGY AND BIOENGINEERING



FUNDAÇÃO  
CALOUSTE  
GULBENKIAN

ESTÉTICA VIVA



Flavour and  
Fragrance Journal –  
read the free  
sample issue here.



MILESTONE  
HELPING  
CHEMISTS



Taylor & Francis  
Taylor & Francis Group



I.L.C. - Instrumentos de Laboratório e Científicos, Lda.

FCT

under CBV / IBB  
plurianual funding and  
research contract  
PTDC/AGR-CFL/117026/2010

Fundação para a Ciência e a Tecnologia  
MINISTÉRIO DA EDUCAÇÃO E CIÊNCIA

The generous obligingness of the following entities is gratefully acknowledged:



UNIVERSIDADE  
DE LISBOA

